

3-2010

# Completed Genome Sequence of the Anaerobic Iron-Oxidizing Bacterium *Acidovorax ebreus* Strain TPSY

Kathryne G. Byrne-Bailey  
*University of California, Berkeley*

Karrie A. Weber  
*University of Nebraska-Lincoln, kweber@unl.edu*

Antinea H. Chair  
*University of California, Berkeley*

Saumyaditya Bose  
*University of California, Berkeley*

Traci Knox  
*University of California, Berkeley*

*See next page for additional authors*

Follow this and additional works at: <http://digitalcommons.unl.edu/bioscifacpub>

 Part of the [Life Sciences Commons](#)

---

Byrne-Bailey, Kathryne G.; Weber, Karrie A.; Chair, Antinea H.; Bose, Saumyaditya; Knox, Traci; Spanbauer, Trisha L.; Chertkov, Olga; and Coates, John D., "Completed Genome Sequence of the Anaerobic Iron-Oxidizing Bacterium *Acidovorax ebreus* Strain TPSY" (2010). *Faculty Publications in the Biological Sciences*. 193.  
<http://digitalcommons.unl.edu/bioscifacpub/193>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications in the Biological Sciences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Kathryne G. Byrne-Bailey, Karrie A. Weber, Antinea H. Chair, Saumyaditya Bose, Traci Knox, Trisha L. Spanbauer, Olga Chertkov, and John D. Coates

## Completed Genome Sequence of the Anaerobic Iron-Oxidizing Bacterium *Acidovorax ebreus* Strain TPSY<sup>∇</sup>

Kathryne G. Byrne-Bailey,<sup>1†</sup> Karrie A. Weber,<sup>1,3†</sup> Antinea H. Chair,<sup>1</sup> Saumyaditya Bose,<sup>1</sup> Traci Knox,<sup>1</sup> Trisha L. Spanbauer,<sup>3</sup> Olga Chertkov,<sup>2</sup> and John D. Coates<sup>1\*</sup>

Department of Plant and Microbiology, University of California, Berkeley, California 94720<sup>1</sup>; Los Alamos National Laboratory, Los Alamos, New Mexico 87545<sup>2</sup>; and School of Biological Sciences, University of Nebraska, Lincoln, Nebraska 68588<sup>3</sup>

Received 4 November 2009/Accepted 25 November 2009

***Acidovorax ebreus* strain TPSY is the first anaerobic nitrate-dependent Fe(II) oxidizer for which there is a completed genome sequence. Preliminary protein annotation revealed an organism optimized for survival in a complex environmental system. Here, we briefly report the completed and annotated genome sequence of strain TPSY.**

Microorganisms from diverse anoxic environments are capable of nitrate-dependent Fe(II) oxidation at circumneutral pH (4, 11, 17, 18, 20, 21). Despite their geochemical importance (22), little is known of the underlying biochemical and genetic mechanisms. Genome sequencing of several nitrate-dependent Fe(II) oxidizers will provide insight into this process. By comparing Fe(II) oxidation mechanisms in various organisms, we hope to identify both the conserved and disparate aspects of the metabolism. The genome of *Acidovorax ebreus* strain TPSY is the first of these to be sequenced.

Strain TPSY is a motile, Gram-negative facultative anaerobe isolated from groundwater collected from the U.S. Department of Energy Integrated Field Research Challenge site at Oak Ridge, TN. Growth experiments performed as previously described (21) revealed TPSY's incapacity for lithoautotrophic growth, which was supported by a lack of genes in the genome encoding any known CO<sub>2</sub> fixation pathways. TPSY did grow mixotrophically with Fe(II) as the electron donor and a 0.1 mM acetate carbon source. 16S rRNA gene sequence analysis placed TPSY in the class *Betaproteobacteria* with 99.8% similarity to *Acidovorax* sp. strain JS42 in the family *Comamonadaceae*.

The completed genome consisted of a single circular chromosome of 3,796,573 bp with an average 66.8% G+C content. A total of 3,479 protein-encoding genes were predicted, and 34 (0.98%) had no similarity to public database sequences. Sequencing performed at the Department of Energy Joint Genome Institute (JGI) used Sanger sequencing and 454 pyrosequencing to a depth of 20× coverage. All JGI library construction and sequencing techniques can be found at <http://www.jgi.doe.gov/>. Sequence assembly, quality assessment, and annotation were performed using the software Phred/Phrap/Consed ([www.phrap.com](http://www.phrap.com)) (6–8), Dupfinisher (10), CRITICA (2), GLIMMER, and GENERATION (5) and the JGI Integrated Microbial Genomes site (12). The

completed genome sequence contained 33,341 reads and had an average of ninefold coverage per base and an error rate of <1 in 100,000.

TPSY was named in part for its meandering motility, and its genome confirmed the twitching phenotype with the presence of *pilT*, *pilU*, and a complete set of flagellar and chemotaxis genes. The ability of TPSY to oxidize simple alcohols and acids with oxygen or nitrate respiration was confirmed by the genome. In addition, biosynthetic pathways for all amino acids except tyrosine and phenylalanine were present. No homologues of chorismate mutase (EC 5.4.99.5), an enzyme required for tyrosine and phenylalanine anabolism, were identified. The genome contained both intact Embden-Meyerhof-Parnas and Entner-Doudoroff pathways, in addition to a pentose phosphate pathway and a trichloroacetic acid cycle.

In support of its facultative anaerobicity, a complete set of genes for denitrification and three different terminal oxidases (cytochrome *aa*<sub>3</sub>, *cb*<sub>3</sub>, and cytochrome *d* quinol oxidase) were present. The *cb*<sub>3</sub> and cytochrome *d* oxidases, with their high oxygen affinity, putatively enable survival in microaerobic environments (14).

TPSY had sequences encoding 30 transposases, 11 integrases, and 11 phage/prophage-related genes. A region of particular interest putatively conferred resistance to lead, arsenate, and mercury: *pbrRATARTBC*, *arsRDAB*, and *merRPCADE*. Evidence suggests horizontal transfer and insertion of this region, as it was flanked on the 5' end by λ prophage-related genes and the 3' end encoded a putative Tn21 transposase. Phenotypic studies by the method of Wang et al. (19) revealed MICs of 16 μM phenylmercuric acetate and 250 μM MgCl<sub>2</sub>. TPSY was also capable of growth in the presence of arsenate (10 mM) but did not use it as an electron acceptor.

Related to phage infection, one CRISPR (clustered, regularly interspaced, short palindromic repeats) region (3, 16) was predicted. The core proteins, the *cas1* and *cas2* genes, and a *csn1* gene formed the CRISPR subtype Nmeni, which is associated with vertebrate pathogens and commensals (9). However, the lack of typical pathogenic type I or III secretion systems such as the *hec* cluster of *Dickeya chrysanthemi* (15) or the *inv/spa* system of *Salmonella enterica* serovar Typhimurium

\* Corresponding author. Mailing address: Department of Plant and Microbiology, University of California, Berkeley, CA 94720. Phone: (510) 643-8455. Fax: (510) 642-4995. E-mail: [jcoates@berkeley.edu](mailto:jcoates@berkeley.edu).

† The first two authors contributed equally to this work and are listed alphabetically.

<sup>∇</sup> Published ahead of print on 18 December 2009.

(13) indicated that TPSY would probably not exhibit a pathogenic lifestyle.

**Nucleotide sequence accession number.** The genome sequence of *Acidovorax ebreus* strain TPSY (formerly *Diaphorobacter* sp. strain TPSY) reported in this paper has been deposited in the GenBank database under accession number NC\_011992.

We thank Lynne Goodwin for her management of the sequencing project at the JGI.

This research was supported by Office of Science (BER), U.S. Department of Energy, grant DE-FG02-07ER64390 to J.D.C. at UCB and by the Nebraska Tobacco Settlement Biomedical Research Development Fund to K.A.W. at the University of Nebraska, Lincoln.

The sequencing work on behalf of J.D.C. was performed through the Community Sequencing Project 2008 for the JGI under the auspices of the U.S. Department of Energy Office of Science, Biological and Environmental Research Program, and by the University of California Lawrence Berkeley National Laboratory under contract DE-AC02-05CH11231, Lawrence Livermore National Laboratory under contract DE-AC52-07NA27344, and Los Alamos National Laboratory under contract DE AC02-06NA25396.

#### REFERENCES

- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**:403–410.
- Badger, J. H., and G. J. Olsen. 1999. CRITICA: coding region identification tool invoking comparative analysis. *Mol. Biol. Evol.* **16**:512–524.
- Barrangou, R., C. Fremaux, H. Deveau, M. Richards, P. Boyaval, S. Moineau, D. A. Romero, and P. Horvath. 2007. CRISPR provides acquired resistance against viruses in prokaryotes. *Science* **315**:1709–1712.
- Chaudhuri, S. K., J. G. Lack, and J. D. Coates. 2001. Biogenic magnetite formation through anaerobic biooxidation of Fe(II). *Appl. Environ. Microbiol.* **67**:2844–2848.
- Delcher, A. L., D. Harmon, S. Kasif, O. White, and S. L. Salzberg. 1999. Improved microbial gene identification with GLIMMER. *Nucleic Acids Res.* **27**:4636–4641.
- Ewing, B., and P. Green. 1998. Base-calling of automated sequencer traces using *Phred*. II. Error probabilities. *Genome Res.* **8**:186–194.
- Ewing, B., L. Hillier, M. C. Wendl, and P. Green. 1998. Base-calling of automated sequencer traces using *Phred*. I. Accuracy assessment. *Genome Res.* **8**:175–185.
- Gordon, D., C. Abajian, and P. Green. 1998. *Consed*: a graphical tool for sequence finishing. *Genome Res.* **8**:195–202.
- Haft, D. H., J. Selengut, E. F. Mongodin, and K. E. Nelson. 2005. A guild of 45 CRISPR-associated (Cas) protein families and multiple CRISPR/Cas subtypes exist in prokaryotic genomes. *PLoS Comput. Biol.* **1**:e60.
- Han, C. S., and P. Chain. 2006. Finishing repeat regions automatically with Dupfinisher, p. 141–146. *In* H. R. Arabnia and H. Valafar (ed.), *Proc. 2006 Int. Conf. Bioinformatics Comput. Biol.* CSREA Press, Las Vegas, NV.
- Lack, J. G., S. K. Chaudhuri, R. Chakraborty, L. A. Achenbach, and J. D. Coates. 2002. Anaerobic biooxidation of Fe(II) by *Dechlorosoma suillum*. *Microb. Ecol.* **43**:424–431.
- Markowitz, V. M., E. Szeto, K. Palaniappan, Y. Grechkin, K. Chu, I. M. Chen, I. Dubchak, I. Anderson, A. Lykidis, K. Mavromatis, N. N. Ivanova, and N. C. Kyrpides. 2008. The Integrated Microbial Genomes (IMG) system in 2007: data content and analysis tool extensions. *Nucleic Acids Res.* **36**:D528–D533.
- Nies, D. H. 1995. The cobalt, zinc, and cadmium efflux system CzcABC from *Alcaligenes eutrophus* functions as a cation-proton antiporter in *Escherichia coli*. *J. Bacteriol.* **177**:2707–2712.
- Preisig, O., D. Anthamatten, and H. Henneke. 1993. Genes for a microaerobically induced oxidase complex in *Bradyrhizobium japonicum* are essential for a nitrogen-fixing endosymbiosis. *Proc. Natl. Acad. Sci. U. S. A.* **90**:3309–3313.
- Rojas, C. M., J. H. Ham, W. L. Deng, J. J. Doyle, and A. Collmer. 2002. HecA, a member of a class of adhesins produced by diverse pathogenic bacteria, contributes to the attachment, aggregation, epidermal cell killing, and virulence phenotypes of *Erwinia chrysanthemi* EC16 on *Nicotiana glauca* seedlings. *Proc. Natl. Acad. Sci. U. S. A.* **99**:13142–13147.
- Sorek, R., V. Kunin, and P. Hugenholz. 2008. CRISPR—a widespread system that provides acquired resistance against phages in bacteria and archaea. *Nat. Rev. Microbiol.* **6**:181–186.
- Straub, K. L., M. Benz, B. Schink, and F. Widdel. 1996. Anaerobic, nitrate-dependent microbial oxidation of ferrous iron. *Appl. Environ. Microbiol.* **62**:1458–1460.
- Straub, K. L., and B. E. E. Buchholz-Cleven. 1998. Enumeration and detection of anaerobic ferrous iron-oxidizing, nitrate-reducing bacteria from diverse European sediments. *Appl. Environ. Microbiol.* **64**:4846–4856.
- Wang, Y., M. Moore, H. S. Levinson, S. Silver, C. Walsh, and I. Mahler. 1989. Nucleotide sequence of a chromosomal mercury resistance determinant from a *Bacillus* sp. with broad-spectrum mercury resistance. *J. Bacteriol.* **171**:83–92.
- Weber, K. A., D. B. Hedrick, A. D. Peacock, J. C. Thrash, D. C. White, L. A. Achenbach, and J. D. Coates. 2009. Physiological and taxonomic description of the novel autotrophic, metal oxidizing bacterium, *Pseudogulbenkiania* sp. strain 2002. *Appl. Microbiol. Biotechnol.* **83**:555–565.
- Weber, K. A., J. Pollock, K. A. Cole, S. M. O'Connor, L. A. Achenbach, and J. D. Coates. 2006. Anaerobic nitrate-dependent iron(II) bio-oxidation by a novel lithoautotrophic betaproteobacterium, strain 2002. *Appl. Environ. Microbiol.* **72**:686–694.
- Weber, K. A., M. M. Urrutia, P. F. Churchill, R. K. Kukkadapu, and E. E. Roden. 2006. Anaerobic redox cycling of iron by freshwater sediment microorganisms. *Environ. Microbiol.* **8**:100–113.