

2012

Draft Genome Sequence of the Anaerobic, Nitrate-Dependent, Fe(II)-Oxidizing Bacterium *Pseudogulbenkiania ferrooxidans* Strain 2002

Kathryne G. Byrne-Bailey
University of California, Berkeley

Karrie A. Weber
University of Nebraska-Lincoln, kweber@unl.edu

John D. Coates
University of California, Berkeley, jdcoates@berkeley.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/bioscifacpub>

 Part of the [Life Sciences Commons](#)

Byrne-Bailey, Kathryne G.; Weber, Karrie A.; and Coates, John D., "Draft Genome Sequence of the Anaerobic, Nitrate-Dependent, Fe(II)-Oxidizing Bacterium *Pseudogulbenkiania ferrooxidans* Strain 2002" (2012). *Faculty Publications in the Biological Sciences*. 194.
<http://digitalcommons.unl.edu/bioscifacpub/194>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications in the Biological Sciences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Draft Genome Sequence of the Anaerobic, Nitrate-Dependent, Fe(II)-Oxidizing Bacterium *Pseudogulbenkiania ferrooxidans* Strain 2002

Kathryne G. Byrne-Bailey,^a Karrie A. Weber,^{a,b} and John D. Coates^a

Department of Plant and Microbial Biology, University of California, Berkeley, California, USA,^a and School of Biological Sciences, University of Nebraska, Lincoln, Nebraska, USA^b

***Pseudogulbenkiania ferrooxidans* strain 2002 was isolated as a lithoautotrophic, Fe(II)-oxidizing, nitrate-reducing bacterium. Phylogenetically, it is in a clade within the family *Neisseriaceae* in the order *Nessieriales* of the class *Betaproteobacteria*. It is anticipated that comparative genomic analysis of this strain with other nitrate-dependent, Fe(II)-oxidizing bacteria will aid in the elucidation of the genetics and biochemistry underlying this critically important geochemical metabolism.**

Pseudogulbenkiania ferrooxidans strain 2002 (formerly *Lutiella nitroferum* strain 2002) is a non-spore-forming, Gram-negative, motile rod belonging to the family *Neisseriaceae* in the order *Nessieriales* of the class *Betaproteobacteria* (13). Strain 2002 was isolated as part of a study of nitrate-dependent Fe(II) oxidation in freshwater lake sediments and characterized as the first mesophilic, nitrate-dependent, lithoautotrophic Fe(II) oxidizer (13). The 16S rRNA gene sequence had 99.3% similarity to the type species *Pseudogulbenkiania subflava* (10), whose genomic sequence is not publically available, and 99.8% similarity to *Pseudogulbenkiania* sp. strain NH8B, whose complete genomic sequence is available (8). The latter nitrate-reducing strain was isolated from agricultural paddy rice and soybean fields in Japan (12). It is anticipated that comparative genomics of *P. ferrooxidans* strain 2002 with other nitrate-dependent, Fe(II)-oxidizing bacteria will aid in the elucidation of the genetics underlying this globally important metabolism.

The draft genome of strain 2002 consisted of 4.2 Mb in 20 contigs with an average G+C content of 64.66%. A total of 3,927 protein-encoding genes were predicted. Sequencing performed at the Joint Genome Institute (JGI) included Sanger sequencing and 454 pyrosequencing to a depth of 20× coverage. All JGI library construction and sequencing techniques can be found at <http://www.jgi.doe.gov/>. Sequence assembly, quality assessment, and annotation were performed using the Phred/Phrap/Consed (CodonCode Corporation, Dedham, MA) (4, 5), Dupfinisher (6), PRODIGAL (7), and GENERATION software and the JGI Integrated Microbial Genomes site (11).

Early comparative genomics revealed little synteny between strain 2002 and the genome of *Acidovorax ebreus* strain TPSY, the first nitrate-dependent iron oxidizer for which there is a publically available genome sequence (2; K. G. Byrne-Bailey et al., unpublished data). In addition, preliminary annotation did not reveal a known carbon fixation pathway that would allow strain 2002 to survive by a lithoautotrophic lifestyle.

Type *c* cytochromes have been implicated in the mechanism of nitrate-dependent Fe(II) oxidation (3), with further biophysical evidence for strain 2002 (13) and *A. ebreus* strain TPSY (Byrne-Bailey et al., unpublished). Automated annotation of the genome of strain 2002 identified 41 genes predicted to encode *c*-type cytochromes, only 1 of which was a diheme, with no predicted proteins

containing multiple heme domains. This is equivalent to both the predicted number of cytochrome proteins in *A. ebreus* strain TPSY and other organisms known to metabolize iron (Byrne-Bailey et al., unpublished). However, manual curation of the genome identified a number of low-similarity homologues to *pio* genes and MtrA proteins which have defined multiple heme domains, none of which were clustered in strain 2002. Iron oxidation in the phototroph *Rhodospirillum rubrum* TIE-1 was shown to involve a number of proteins encoded by the *pio* genes (9), whereas iron reduction by *Shewanella* spp. involves the Mtr proteins (1). It is hoped that further investigation of these and other genes will aid in the understanding of nitrate-dependent iron oxidation in neutrophilic bacteria.

Nucleotide sequence accession number. The genomic sequence of *P. ferrooxidans* strain 2002 reported here has been deposited in the GenBank database under accession number NZ_ACIS01000000.

ACKNOWLEDGMENTS

We thank Lynne Goodwin for her management of the sequencing project at the JGI.

The sequencing work was performed through Community Sequencing Project 2008 for the JGI under the auspices of the U.S. Department of Energy Office of Science Biological and Environmental Research Program and by the University of California Lawrence Berkeley National Laboratory under contract DE-AC02-05CH11231, the Lawrence Livermore National Laboratory under contract DE-AC52-07NA27344, and the Los Alamos National Laboratory under contract DE AC02-06NA25396. Research in the lab of J.D.C. on nitrate-dependent Fe(II) oxidation is funded by the Energy Biosciences Institute, Berkeley, CA.

REFERENCES

1. Beliaev AS, Saffarini DA. 1998. *Shewanella putrefaciens mtrB* encodes an outer membrane protein required for Fe(III) and Mn(IV) reduction. *J. Bacteriol.* 180:6292–6297.
2. Byrne-Bailey KG, et al. 2010. Completed genome sequence of the anaerobic

Received 10 February 2012 Accepted 20 February 2012

Address correspondence to John D. Coates, jcoates@berkeley.edu.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.00214-12

- obic iron-oxidizing bacterium *Acidovorax ebreus* strain TPSY. *J. Bacteriol.* **192**:1475–1476.
3. Chaudhuri SK, Lack JG, Coates JD. 2001. Biogenic magnetite formation through anaerobic biooxidation of Fe(II). *Appl. Environ. Microbiol.* **67**: 2844–2848.
 4. Ewing B, Green P. 1998. Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res.* **8**:186–194.
 5. Ewing B, Hillier L, Wendl MC, Green P. 1998. Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res.* **8**:175–185.
 6. Han CS, Chain P. 2006. Finishing repeat regions automatically with Dupfinisher, p 141–146. *In* Arabnia HR, Valafar H (ed), *Proceeding of the 2006 International Conference on Bioinformatics & Computational Biology*. CSREA Press, Las Vegas, NV.
 7. Hyatt D, et al. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**:119.
 8. Ishii S, et al. 2011. Complete genome sequence of the denitrifying and N(2)O-reducing bacterium *Pseudogulbenkiania* sp. strain NH8B. *J. Bacteriol.* **193**:6395–6396.
 9. Jiao Y, Newman DK. 2007. The *pio* operon is essential for phototrophic Fe(II) oxidation in *Rhodopseudomonas palustris* TIE-1. *J. Bacteriol.* **189**: 1765–1773.
 10. Lin MC, Chou JH, Arun AB, Young CC, Chen WM. 2008. *Pseudogulbenkiania subflava* gen. nov., sp. nov., isolated from a cold spring. *Int J. Syst Evol Microbiol.* **58**:2384–2388.
 11. Markowitz VM, et al. 2008. The integrated microbial genomes (IMG) system in 2007: data content and analysis tool extensions. *Nucleic Acids Res.* **36**:D528–D533.
 12. Tago K, Ishii S, Nishizawa T, Otsuka S, Senoo K. 2011. Phylogenetic and functional diversity of denitrifying bacteria isolated from various rice paddy and rice-soybean rotation fields. *Microbes Environ.* **26**: 30–35.
 13. Weber KA, et al. 2009. Physiological and taxonomic description of the novel autotrophic, metal oxidizing bacterium, *Pseudogulbenkiania* sp. strain 2002. *Appl. Microbiol. Biotechnol.* **83**:555–565.