

2011

# Hidden SUSY at the LHC: the light higgsino-world scenario and the role of a lepton collider

Howard Baer

*University of Oklahoma*, baer@ou.edu

Vernon Barger

*University of Wisconsin*, barger@pheno.wisc.edu

Peisi Huang

*University of Wisconsin*, peisi.huang@unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsfacpub>

---

Baer, Howard; Barger, Vernon; and Huang, Peisi, "Hidden SUSY at the LHC: the light higgsino-world scenario and the role of a lepton collider" (2011). *Faculty Publications, Department of Physics and Astronomy*. 197.  
<http://digitalcommons.unl.edu/physicsfacpub/197>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications, Department of Physics and Astronomy by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

RECEIVED: August 4, 2011

REVISED: October 10, 2011

ACCEPTED: October 21, 2011

PUBLISHED: November 8, 2011

# Hidden SUSY at the LHC: the light higgsino-world scenario and the role of a lepton collider

Howard Baer,<sup>a</sup> Vernon Barger<sup>b</sup> and Peisi Huang<sup>b</sup>

<sup>a</sup>*Dept. of Physics and Astronomy, University of Oklahoma,  
Norman, OK 73019, U.S.A.*

<sup>b</sup>*Dept. of Physics, University of Wisconsin,  
Madison, WI 53706, U.S.A.*

*E-mail:* [baer@nhn.ou.edu](mailto:baer@nhn.ou.edu), [barger@pheno.wisc.edu](mailto:barger@pheno.wisc.edu), [phuang7@wisc.edu](mailto:phuang7@wisc.edu)

**ABSTRACT:** While the SUSY flavor, CP and gravitino problems seem to favor a very heavy spectrum of matter scalars, fine-tuning in the electroweak sector prefers low values of superpotential mass  $\mu$ . In the limit of low  $\mu$ , the two lightest neutralinos and light chargino are higgsino-like. The light charginos and neutralinos may have large production cross sections at LHC, but since they are nearly mass degenerate, there is only small energy release in three-body sparticle decays. Possible dilepton and trilepton signatures are difficult to observe after mild cuts due to the very soft  $p_T$  spectrum of the final state isolated leptons. Thus, the higgsino-world scenario can easily elude standard SUSY searches at the LHC. It should motivate experimental searches to focus on dimuon and trimuon production at the very lowest  $p_T(\mu)$  values possible. If the neutralino relic abundance is enhanced via non-standard cosmological dark matter production, then there exist excellent prospects for direct or indirect detection of higgsino-like WIMPs. While the higgsino-world scenario may easily hide from LHC SUSY searches, a linear  $e^+e^-$  collider or a muon collider operating in the  $\sqrt{s} \sim 0.5 - 1$  TeV range would be able to easily access the chargino and neutralino pair production reactions.

**KEYWORDS:** Supersymmetry Phenomenology

**ARXIV EPRINT:** [1107.5581](https://arxiv.org/abs/1107.5581)

---

**Contents**

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Higgsino-world parameter space and mass spectra</b>	<b>4</b>
<b>3</b>	<b>Neutralino relic density and direct detection rates</b>	<b>8</b>
<b>4</b>	<b>Higgsino-world scenario at the LHC</b>	<b>10</b>
4.1	Sparticle production at LHC7	10
4.2	Branching fractions and collider signatures	11
4.3	Collimated dilepton +MET signature from $\tilde{Z}_1\tilde{Z}_2$ production	12
<b>5</b>	<b>Prospects for ILC or a muon collider</b>	<b>15</b>
<b>6</b>	<b>Summary and conclusions</b>	<b>17</b>

---

**1 Introduction**

The well-known instability of the scalar sector of the Standard Model (SM) to quadratic divergences is elegantly solved by the introduction of supersymmetry (SUSY) (the Minimal Supersymmetric Standard Model commonly used today was introduced by [1]; for reviews of SUSY phenomenology, see [3]; [2, 4–6]). In the case of the Minimal Supersymmetric Standard Model (MSSM) with soft SUSY breaking (SSB) terms, the divergences in the scalar sector are rendered to merely logarithmic. Interplay between the electroweak sector and the SUSY partners suggests the superpartner masses should exist at or around the TeV scale to avoid re-introduction of fine-tuning.

While the MSSM may be very appealing, it does suffer several pathologies. Unfettered soft SUSY breaking terms lead to large rates for flavor-changing neutral current processes and  $CP$  violation [7]. Inclusion of grand unified theories with SUSY may lead to unacceptably high rates for proton decay [8]. And in gravity-mediated SUSY breaking models (SUGRA), gravitino production followed by late-time gravitino decays in the early universe are in conflict with the successful picture of Big Bang nucleosynthesis (BBN) unless re-heat temperatures after inflation are limited to  $T_R \lesssim 10^5$  GeV [9, 10]. The latter bound is in conflict with appealing baryogenesis models such as thermal [11–20] (or non-thermal [21–24]) leptogenesis, which require  $T_R \gtrsim 2 \times 10^9$  GeV ( $10^6$  GeV).

A common solution to the above four problems is to push the SUSY matter scalars into the multi-TeV regime [25–27]. The heavy scalars thus suppress loop-induced flavor and  $CP$  violating processes, and suppress proton decay rates. If the multi-TeV scalars derive from a SUGRA model with a simple form for the Kähler potential, then the gravitino mass

$m_{3/2}$  is also expected to exist in the multi-TeV range. By pushing  $m_{3/2}$  into the 10-50 TeV range, the gravitino lifetime can be reduced to  $\tau_{3/2} \lesssim 1$  second, so the gravitino decays shortly before BBN begins, this solving the gravitino problem [10, 28].

At first glance, multi-TeV gravitino and scalar masses seem in conflict with SUSY electroweak fine-tuning. The possible SUSY electroweak fine-tuning arises from minimization of the scalar potential after electroweak symmetry breaking. Here, the tree-level electroweak breaking conditions are familiarly written as [29]

$$B\mu = \frac{(m_{H_u}^2 + m_{H_d}^2 + 2\mu^2) \sin 2\beta}{2}, \quad \text{and} \quad (1.1)$$

$$\mu^2 = \frac{m_{H_d}^2 - m_{H_u}^2 \tan^2 \beta}{(\tan^2 \beta - 1)} - \frac{M_Z^2}{2}, \quad (1.2)$$

where  $B$  is the bilinear SSB term and  $m_{H_u}^2$  and  $m_{H_d}^2$  are the up and down Higgs SSB masses evaluated at the weak scale,  $\mu$  is the superpotential Higgs mass term and  $\tan \beta$  is the ratio of Higgs field vevs:  $\tan \beta = \frac{v_u}{v_d}$ .

A measure of fine-tuning

$$\Delta_i \equiv \left| \frac{\partial \log M_Z^2}{\partial \log a_i} \right| \quad (1.3)$$

was advocated in ref. [30]. More sophisticated measures were advocated in Ref's [31], while in ref. [32], the  $\mu$  parameter itself is taken as a measure of fine-tuning: the latter paper requires  $|\mu| \lesssim 1$  TeV to avoid too much fine-tuning. This measure motivates the well-known hyperbolic branch/ focus point (HB/FP) region of minimal supergravity (mSUGRA or CMSSM) as allowing for heavy scalars with low  $\mu$  value and low fine-tuning [32–34]. A virtue of the HB/FP region is that multi-TeV scalars can co-exist with apparent low levels of electroweak fine-tuning.

In this paper, we will consider supersymmetric models with large, multi-TeV scalar masses, but with low, sub-TeV superpotential  $\mu$  term. We consider the case with intermediate range gaugino masses. This scenario, with

$$|\mu| \ll m_{gauginos} \ll m_{scalars}, \quad (1.4)$$

has been dubbed “higgsino-world” by Kane [35, 36], and leads to a sparticle mass spectrum with a light higgsino-like chargino  $\widetilde{W}_1$  and two light higgsino-like neutralinos  $\widetilde{Z}_1$  and  $\widetilde{Z}_2$ . In models with gaugino mass unification at  $M_{GUT}$ , then the state  $\widetilde{Z}_3$  will be mainly bino-like, while  $\widetilde{Z}_4$  and  $\widetilde{W}_2$  will be wino-like.

While the higgsino-world scenario seems highly appealing due to its ability to reconcile multi-TeV scalars and gravitinos with low electroweak fine-tuning, it has perhaps fallen out of favor for two reasons. First, higgsino-world leads to a very low thermal relic density of neutralinos, not at all in accord with measurements from WMAP and other experiments which require [37]

$$\Omega_{DM} h^2 = 0.1123 \pm 0.0035 \quad \text{at 68\% CL.} \quad (1.5)$$

Second, higgsino-world scenarios are not easily realized in the paradigm mSUGRA/CMSSM framework, since elevating scalar masses into the multi-TeV region for a given value of GUT

scale gaugino mass  $m_{1/2}$  pushes one beyond the HB/FP region into a portion of parameter space where radiative EWSB is not realized under the assumption of universal scalar masses  $m_0$  at  $M_{GUT}$ .

Pertaining to the dark matter issue, a number of recent works have emphasized that the standard picture of a thermal SUSY WIMP as dark matter is subject to very high fine-tuning [38–40]. Furthermore, non-standard cosmologies have many desirable features, and may even be favored by string theoretic constructions. For instance, Kane *et al.* have shown [41] that at least one moduli field in string theory should maintain a mass at or around the 10 TeV scale. Such moduli fields can be produced via coherent oscillations in the early universe, and decay into WIMPs, thereby augmenting the WIMP abundance [42], or they can decay into SM particles, thus generating entropy and diluting the WIMP abundance. Gelmini *et al.* have shown in this case that SUSY models with *any* value — either too high or too low — of thermal WIMP abundance may give rise to the measured CDM abundance via the enhancement or diminution due to scalar field (moduli) decays [43, 44]. In particular, for higgsino-world with too low a thermal WIMP abundance, the light higgsino abundance can be enhanced by moduli decays, leading to the correct abundance of higgsino-like WIMPs.

Alternatively, in SUSY models wherein the strong  $CP$  problem is solved by the introduction of Peccei-Quinn-Weinberg-Wilczek [45–48] invisible axion [49–52], one must introduce an axion supermultiplet, which contains an  $R$ -parity even spin-0 saxion  $s(x)$ , along with an  $R$ -parity odd spin- $\frac{1}{2}$  axino  $\tilde{a}(x)$ , in addition to the light pseudoscalar axion field  $a(x)$ . In models such as these, with a TeV-scale axino and a higgsino-like neutralino as LSP, the  $\tilde{Z}_1$  abundance can be augmented by axino production and subsequent re-annihilation at temperatures above BBN but below neutralino freeze-out [53]. Depending on the various PQMSSM model parameters, the CDM consists of an axion/neutralino admixture, where either the axion or the neutralino can dominate the abundance [54].

A third modification of the thermal WIMP abundance may also occur: a model with a supposed underabundance of neutralino dark matter may enjoy enhancement of the relic DM abundance due to thermal gravitino production [55–57] followed by cascade decays to the LSP state [58–61].

Pertaining to the issue of higgsino-world being difficult to realize in the paradigm mSUGRA model, we note that it is easily realized in models with non-universal GUT scale Higgs masses (NUHM) [62–65]. In fact, in GUT models such as SO(10), the matter supermultiplets live in the 16-dimensional spinor representation, while Higgs superfields live in 10 or other dimensional multiplets. In such models, there is little reason to expect matter-Higgs SSB universality at  $M_{GUT}$ .

For the above reasons, we feel that it may be opportune to reconsider the higgsino-world scenario, and whether such a scenario would be visible to LHC SUSY searches. Toward this end, we discuss in section 2 the higgsino-world parameter space and expected mass spectra. In section 3, we present calculations of the standard thermal neutralino abundance in the higgsino-world scenario, and discuss its direct and indirect detection in the case where non-standard cosmological processes augment the relic higgsino abundance. In section 4, we evaluate the dominant sparticle production cross sections for the lighter matter states at

the LHC and calculate their branching fractions. While higgsino production cross sections occur at possibly observable levels, the compressed spectra lead to sparticle decays with very low energy release, and very soft detectable particles. To the best of our knowledge, higgsino-world SUSY can effectively elude standard SUSY searches for jets plus missing  $E_T$  ( $MET$ ), and also for isolated multi-leptons+ $MET$  at LHC7 (LHC at  $\sqrt{s} = 7$  TeV) with  $\sim 10 \text{ fb}^{-1}$  of integrated luminosity. In section 5, we discuss higgsino-world signatures at a TeV scale lepton collider such as ILC or a muon collider (MC). In section 6, we present our final discussion and conclusions.

## 2 Higgsino-world parameter space and mass spectra

We will adopt the Isajet 7.81 program for SUSY particle mass spectrum generation [71, 72]. To generate spectra in higgsino-world scenario, we will adopt the Isasugra non-universal Higgs mass parameter space (NUHM2):

$$m_0, m_{1/2}, A_0, \tan\beta, \mu, m_A. \tag{2.1}$$

In the above parameter space,  $m_0$ ,  $m_{1/2}$  and  $A_0$  are the usual GUT scale parameters, although here  $m_0$  is reserved only for matter scalars, and not Higgs scalar soft masses. The two additional parameters  $\mu$  and  $m_A$  are stipulated at the weak scale, and are used to solve for the weak scale values of  $m_{H_u}^2$  and  $m_{H_d}^2$ . These latter parameters are run from the weak to GUT scale, and their GUT scale values are determined by enforcing the input weak scale values of  $\mu$  and  $m_A$ . We will take  $m_0 \sim m_{\tilde{G}}$  to be in the multi-TeV range, so that we obtain a decoupling solution to the SUSY flavor, CP,  $p$ -decay and gravitino problems. Thus, the parameters  $m_0$  and also  $A_0$  and  $m_A$  will be largely irrelevant for our analysis. The main parameter space dependence will arise from just varying  $\mu$  and  $m_{1/2}$ . Since we are interested in the light higgsino-world scenario, with  $\mu \ll M_i$  (where  $M_i$  are the weak scale gaugino masses), the parameter  $\tan\beta$ , which induces gaugino-higgsino mixing, will also not be terribly relevant.

The higgsino-world input parameters and mass spectra for two sample benchmark points with  $\mu = 150$  and  $300$  GeV are listed in table 1. We also take  $m_0 = 5000$  GeV,  $m_{1/2} = 800$  GeV,  $A_0 = 0$ ,  $\tan\beta = 10$  and  $m_A = 800$  GeV. The spectra are also shown in figure 1 for the higgsino-world case where  $\mu = 150$  GeV (HW150).

From table 1 or figure 1, we see that the three states  $\tilde{W}_1$ ,  $\tilde{Z}_1$  and  $\tilde{Z}_2$  all have masses clustered around the value of  $\mu = 150$  or  $300$  GeV. These states are dominantly higgsino-like. The weak scale gaugino masses  $M_1 \sim 352$  GeV and  $M_2 \sim 638$  GeV for the two cases, so that  $\tilde{Z}_3$  is bino-like and  $\tilde{W}_2$  and  $\tilde{Z}_4$  are wino-like. The squarks and sleptons all are decoupled, with masses in the multi-TeV range, since  $m_0 = 5$  TeV. The  $\tilde{Z}_2 - \tilde{Z}_1$  mass gap is just 16.2 GeV and 28.2 GeV, respectively, for the two cases. We also show the higgsino fraction of the lightest neutralino:  $v_H = \sqrt{v_1^{(1)2} + v_2^{(1)2}}$  where  $v_1^{(i)}$  is the higgsino  $\tilde{h}_u^0$  content and  $v_2^{(i)}$  is the higgsino  $\tilde{h}_d^0$  content of neutralino  $\tilde{Z}_i$  in the notation of ref. [29]. Here, the value of  $v_H$  is 0.98 for HW150 and 0.9 for HW300. Increasing  $\mu$  to 500 GeV, using the same choice of other model parameters, decreases  $v_H \sim 0.21$ , so that in this case the lightest


parameter	HW150	HW300
$m_0$	5000	5000
$m_{1/2}$	800	800
$A_0$	0	0
$\tan \beta$	10	10
$\mu$	150	300
$m_A$	800	800
$m_{\tilde{g}}$	2004.9	2004.2
$m_{\tilde{u}_L}$	5171.5	5171.4
$m_{\tilde{t}_1}$	3240.2	3243.8
$m_{\tilde{b}_1}$	4267.8	4269.4
$m_{\tilde{e}_R}$	4869.4	4870.1
$m_{\tilde{W}_2}$	672.7	675.4
$m_{\tilde{W}_1}$	156.3	310.5
$m_{\tilde{Z}_4}$	688.2	691.0
$m_{\tilde{Z}_3}$	356.3	366.9
$m_{\tilde{Z}_2}$	158.9	311.4
$m_{\tilde{Z}_1}$	142.7	283.2
$m_h$	120.1	120.1
$\sigma(LHC7)$	1055 fb	63.5 fb
$\Omega_{\tilde{Z}_1}^{std} h^2$	0.008	0.03
$BF(b \rightarrow s\gamma)$	$3.5 \times 10^{-4}$	$3.5 \times 10^{-4}$
$\sigma^{SI}(\tilde{Z}_1 p)$ (pb)	$1.0 \times 10^{-8}$	$3.1 \times 10^{-8}$
$\langle \sigma v \rangle _{v \rightarrow 0}$ (cm <sup>3</sup> /sec)	$0.28 \times 10^{-24}$	$0.09 \times 10^{-24}$
$v_H^{(1)}$	0.98	0.90

**Table 1.** Input parameters and masses in GeV units for two higgsino-world scenario benchmark points HW150 and HW300, with  $\mu = 150$  and  $300$  GeV, respectively.


neutralino is no longer dominantly higgsino-like, but rather of mixed higgsino-bino variety. We also see from table 1 that the standard thermal neutralino abundance is  $\Omega_{\tilde{Z}_1}^{std} h^2 \sim 0.008$  and  $0.03$ , respectively, *i.e.* well below the WMAP-measured CDM abundance.

In figure 2, we show color-coded contours of the higgsino fraction  $v_H$  of the lightest neutralino  $\tilde{Z}_1$  in the  $\mu$  vs.  $m_{1/2}$  parameter space plane for  $m_0 = 5$  TeV,  $A_0 = 0$  and  $\tan \beta = 10$ . The green, yellow and especially red regions contain a lightest neutralino with large higgsino fraction  $v_H \gtrsim 0.5$ . This region essentially defines the higgsino-world scenario parameter space, which is found at low  $|\mu|$  and large  $m_{1/2}$ . As one enters the blue-shaded region, the  $\tilde{Z}_1$  becomes increasingly bino-like. The unshaded region at low  $\mu$  is excluded by LEP2 limits on the lightest chargino:  $m_{\tilde{W}_1} \lesssim 103.5$  GeV.

In figure 3, we show a). the color-coded mass contours of the lightest neutralino  $\tilde{Z}_1$ , and b). the  $m_{\tilde{Z}_2} - m_{\tilde{Z}_1}$  mass gap (which is always very close to the value of the  $m_{\tilde{W}_1} - m_{\tilde{Z}_1}$  mass gap). In the higgsino-world scenario with low  $\mu$  and large  $m_{1/2}$ , we find that  $m_{\tilde{Z}_1}$




**Figure 1.** Sparticle mass spectra for a light higgsino world scenario with  $\mu = 150$  GeV,  $m_A = 800$  GeV,  $m_0 = 5000$  GeV,  $m_{1/2} = 600$  GeV,  $A_0 = 0$  and  $\tan\beta = 10$ . We take  $m_t = 173.3$  GeV.



**Figure 2.** Color-coded contours of higgsino content  $v_H$  of the  $\tilde{Z}_1$  in the  $\mu$  vs.  $m_{1/2}$  plane with  $m_A = 800$  GeV,  $m_0 = 5000$  GeV,  $A_0 = 0$  and  $\tan\beta = 10$ .


can drop as low as  $\sim 90$  GeV, where the lower limit comes from the LEP2 constraint on chargino masses. Meanwhile, the mass gap  $m_{\tilde{Z}_2} - m_{\tilde{Z}_1}$  drops as low as  $\sim 10$  GeV in the extreme higgsino-world region. Thus, for extreme higgsino-world parameters, we always





**Figure 3.** In a)., we plot contours of  $m_{\tilde{Z}_1}$  while in b). we plot contours of  $m_{\tilde{Z}_2} - m_{\tilde{Z}_1}$  in the  $\mu$  vs.  $m_{1/2}$  plane with  $m_A = 800$  GeV,  $m_0 = 5000$  GeV,  $A_0 = 0$  and  $\tan\beta = 10$ .

expect the  $\tilde{Z}_2$  states to decay via three-body modes or loop-suppressed two-body decays such as  $\tilde{Z}_2 \rightarrow \tilde{Z}_1\gamma$  [86]. Two-body decays such as  $\tilde{Z}_2 \rightarrow \tilde{Z}_1 Z$  or  $\tilde{Z}_1 h$  will always be closed in the higgsino-world scenario.




**Figure 4.** Regions of thermal neutralino abundance in the  $\mu$  vs.  $m_{1/2}$  plane with  $m_A = 800$  GeV,  $m_0 = 5000$  GeV,  $A_0 = 0$  and  $\tan\beta = 10$ .

### 3 Neutralino relic density and direct detection rates

In the light higgsino-world scenario, if the lightest neutralino is dominantly higgsino-like, then it will sustain large  $\tilde{Z}_1\tilde{Z}_1$  annihilation cross sections into vector boson states  $ZZ$  and  $W^+W^-$ . This will result in a standard thermal neutralino abundance typically well below WMAP-measured values of  $\Omega_{CDM}h^2 \sim 0.11$ . In figure 4, we show color-coded contours of the standard thermal neutralino abundance  $\log_{10}\Omega_{\tilde{Z}_1}^{std}h^2$ . We see that indeed in the low  $\mu$  region  $\Omega_{\tilde{Z}_1}^{std}h^2$  is at the  $10^{-2.5} - 10^{-1}$  range, where  $10^{-1}$  occurs for mixed higgsino-bino states. Thus, a non-standard cosmology is likely needed to explain the CDM abundance in the higgsino-world scenario. Indeed, many “non-standard” scenarios can be highly motivated by other physics considerations (the presence of TeV-scale moduli in string theory, the axion solution to the strong CP problem  $\dots$ ), and so may well be more appealing than simple thermal production of WIMPs.

If the higgsino relic abundance is enhanced (say, by moduli decays, or by axino production and decay, or by gravitino production and decay) beyond standard expectations, then a higgsino-like WIMP may well make up the bulk of dark matter. In this case, we present in figure 5 color-coded contours of spin-independent neutralino-proton scattering cross section in units of  $10^{-9}$  pb in the  $\mu$  vs.  $m_{1/2}$  plane. The red and yellow shaded regions have  $\sigma_{SI}(p\tilde{Z}_1) \gtrsim 30 \times 10^{-9}$  pb, while green-shaded regions have  $\sigma_{SI}(p\tilde{Z}_1) \gtrsim 20 \times 10^{-9}$  pb. The Xenon-100 experiment [87] — for  $m_{WIMP} \sim 100 - 200$  GeV — excludes  $\sigma_{SI}(\tilde{Z}_1p) \gtrsim 10^{-8}$  pb, so already a large portion of higgsino-world parameter space is excluded if higgsino-like WIMPs make up all the CDM. It is shown in ref. [54] that in the case of the Peccei-Quinn




**Figure 5.** Contours of  $\sigma^{SI}(\tilde{Z}_1 p)$  in units of  $10^{-9}$  pb in the  $\mu$  vs.  $m_{1/2}$  plane with  $m_A = 800$  GeV,  $m_0 = 5000$  GeV,  $A_0 = 0$  and  $\tan\beta = 10$ .

augmented MSSM, where an axion-axino-saxion supermultiplet is required to solve the strong  $CP$ -problem, that for some ranges of PQMSSM parameters, higgsino-like WIMPs could make up virtually all the DM abundance, while for other PQMSSM parameters, (low re-heat temperature  $T_R$  or if cosmologically produced axinos decay to neutralinos before neutralino freeze-out), then the CDM abundance may be axion-dominated, while the higgsino abundance maintains its standard relic density. In this latter case, the assumed WIMP abundance would have to be scaled down by a factor of 10-100, and so the higgsino-world scenario would then escape Xenon-100 null-search constraints.

If higgsino-like WIMPs comprise the bulk of dark matter, then it may also be possible to detect them via searches for galactic halo WIMP annihilation into final states containing positrons, anti-protons, gamma-rays [88, 89] or anti-deuterons [90]. In these cases, the WIMP annihilation rate is always proportional to thermally averaged WIMP annihilation cross section times relative velocity, in the limit where  $v \rightarrow 0$  (in the galactic halo):  $\langle\sigma v\rangle|_{v\rightarrow 0}$ . The exact detection rates will also depend on various astrophysical quantities, and details of the detection devices and their backgrounds. Here, we merely present color-coded contours of  $\langle\sigma v\rangle|_{v\rightarrow 0}$  in units of  $10^{-24} \text{cm}^3/\text{sec}$ . The red-, yellow- and green-shaded regions will typically lead to observable levels of gamma-ray or antimatter detection rates, if higgsino-like WIMPs dominate the CDM relic density.<sup>1</sup> However, in scenarios like

<sup>1</sup>Recently the Fermi LAT collaboration has studied gamma ray emissions from a variety of Milky Way satellite galaxies, and has placed new limits on  $\langle\sigma v\rangle|_{v\rightarrow 0}$ , ref. [91]. For WIMPs of mass  $\sim 200$  GeV, they exclude  $\langle\sigma v\rangle|_{v\rightarrow 0} \gtrsim 10^{-25} \text{cm}^3/\text{s}$  in the case of dominant WIMP annihilation into  $WW$ . In our case, higgsino-like neutralinos dominantly annihilate into  $WW$  or  $ZZ$  for  $m_{\text{higgsino}} < m_t$ , and into  $t\bar{t}$  for  $m_{\text{higgsino}} \gtrsim m_t$ . Thus, the deep higgsino region (red-to-green shaded) of figure 6 would now be excluded in the case where higgsinos comprised the bulk of dark matter.



**Figure 6.** Contours of  $\langle \sigma v \rangle|_{v \rightarrow 0}$  in units of  $10^{-24} \text{ cm}^3/\text{sec}$  in the  $\mu$  vs.  $m_{1/2}$  plane with  $m_A = 800 \text{ GeV}$ ,  $m_0 = 5000 \text{ GeV}$ ,  $A_0 = 0$  and  $\tan \beta = 10$ .


the PQMSSM with mixed axion-WIMP dark matter, but with axion domination, these rates will be suppressed due to the low halo abundance of higgsino-like WIMPs. In other PQMSSM cases where the axino  $\tilde{a}$  is the LSP ([92–95]; for a recent review of axion/axino dark matter, see [96]), then the higgsino-like WIMPs would all have decayed to relic axinos, and no direct or indirect detection signals would be seen (although detection of relic axions would still be possible).

## 4 Higgsino-world scenario at the LHC

### 4.1 Sparticle production at LHC7

In the HW scenario, squarks and sleptons are assumed decoupled from collider physics. In the limit of large scalar masses, the reach of LHC7 [73] with  $2 \text{ fb}^{-1}$  for gluino pair production is to  $m_{1/2} \sim 250 \text{ GeV}$  (corresponding to  $m_{\tilde{g}} \sim 700 \text{ GeV}$ ), while the reach of LHC14 with  $100 \text{ fb}^{-1}$  is to  $m_{1/2} \sim 650 \text{ GeV}$  (corresponding to  $m_{\tilde{g}} \sim 1400 \text{ GeV}$ ) [74–81]. Thus, for most of HW parameter space, gluino pair production will be below LHC sensitivity. We then expect chargino/neutralino pair production to be the most promising SUSY cross sections at LHC.

In figure 7, we show the dominant sparticle pair production cross sections in  $fb$  for LHC7 from the higgsino-world scenario versus  $\mu$  for other model parameters as in table 1. We adopt the computer code Prospino so that the results are valid at NLO in QCD [82]. For low values of  $\mu \lesssim 300 \text{ GeV}$  (the deep higgsino region), we see that  $\widetilde{W}_1^\pm \widetilde{Z}_1$  and  $\widetilde{W}_1^\pm \widetilde{Z}_2$  are



**Figure 7.** Dominant chargino and neutralino production cross sections versus  $\mu$  at LHC7 for light higgsino-world SUSY scenario. Other parameters are fixed as in figure 1.

dominant, followed closely by  $\widetilde{W}_1^+ \widetilde{W}_1^-$  and  $\widetilde{Z}_1 \widetilde{Z}_2$  production. As  $\mu$  moves beyond 300 GeV, the lighter charginos and neutralinos become mixed gaugino-higgsino states, and some of the cross sections drop rapidly and become comparable to production rates for some of the heavier charginos and neutralinos.<sup>2</sup> Other potentially visible cross sections such as for  $\widetilde{Z}_2 \widetilde{Z}_2$  are several orders of magnitude below these. The sum total of the reactions shown in figure 7 agrees well with output for all SUSY reactions as generated by Isajet as shown in table 1, so these are indeed the dominant production reactions.

#### 4.2 Branching fractions and collider signatures

The sparticle branching fractions can be read off from the Isajet decay table for sparticle cascade decays [83–85]. For light charginos  $\widetilde{W}_1$ , we find

- $\widetilde{W}_1^- \rightarrow \ell \bar{\nu}_\ell \widetilde{Z}_1$  at 11.1% for each species  $\ell = e, \mu$  or  $\tau$ ,
- $\widetilde{W}_1^- \rightarrow d \bar{u} \widetilde{Z}_1$  at 33.3% ,
- $\widetilde{W}_1^- \rightarrow s \bar{c} \widetilde{Z}_1$  at 33.3%

since the three-body chargino decays are dominated by the  $W^*$  propagator.

For  $\widetilde{Z}_2$ , we find typically

- $\widetilde{Z}_2 \rightarrow \ell^+ \ell^- \widetilde{Z}_1$  at 3.5% for each species  $\ell = e, \mu$  or  $\tau$ ,

<sup>2</sup>Reactions such as  $\widetilde{W}_1 \widetilde{Z}_3$  production may offer some hope for sparticle detection at LHC since  $\widetilde{Z}_3 \rightarrow \widetilde{W}_1 W$  or  $\widetilde{Z}_i Z$  or  $\widetilde{Z}_1 h$ . Since the decay products of  $\widetilde{W}_1$  are largely invisible, these signal reactions would suffer enormous backgrounds from direct  $W$  or  $Z$  production and vector boson pair production.

- $\tilde{Z}_2 \rightarrow \nu_\ell \bar{\nu}_\ell \tilde{Z}_1$  at 21.5% (summed over all neutrino species) and
- $\tilde{Z}_2 \rightarrow q\bar{q}\tilde{Z}_1$  at 68% summed over all quark species.

In addition, the decay  $\tilde{Z}_2 \rightarrow \tilde{Z}_1\gamma$  occurs at an enhanced rate: 0.8% (0.2%) for HW150 (HW300) [86].

By combining production cross sections with branching fractions, we find that  $\tilde{W}_1\tilde{Z}_1$  production will lead to either soft jets+MET (likely buried under QCD background(BG)) or a soft isolated lepton+MET (likely buried under BG from direct  $W$ -boson production). Thus, we do not expect this reaction to lead to observable signatures.

The reaction  $\tilde{W}_1^+\tilde{W}_1^-$  will lead to either 1) soft jets+MET, 2) soft isolated lepton plus jets+MET or 3) soft dilepton pair+MET. We expect each of these also to be buried beneath SM backgrounds from QCD or vector boson pair production.

The reaction  $\tilde{Z}_1\tilde{Z}_2$  production can lead to 1) soft jets+MET or 2) soft, low invariant mass dilepton pairs+MET. The first of these is likely buried beneath QCD background. The second of these has a chance at observability since the  $m(\ell^+\ell^-)$  will be bounded by  $m_{\tilde{Z}_2} - m_{\tilde{Z}_1}$  and thus lead to a distinctive mass edge upon a continuum background arising from  $WW$  or  $Z\gamma^*$  production, where  $\gamma^* \rightarrow \ell^+\ell^-$ . If the dilepton pair is at high  $p_T$ , due to a highly boosted  $\tilde{Z}_2$ , then we expect the dilepton pair to be highly collimated in opening angle, and to appear rather distinctively compared to known backgrounds.

The reaction  $\tilde{W}_1\tilde{Z}_2$  production will lead to either 1) soft jets+MET, 2) soft jets plus collimated soft dilepton+MET or 3) soft trileptons+MET. The first of these is likely buried beneath QCD background. The second is possibly observable, and should be present if the cleaner  $\tilde{Z}_1\tilde{Z}_2 \rightarrow \ell^+\ell^- + MET$  is found. The third case yields the venerable clean trilepton signature which has been evaluated for the Tevatron [66–70] and LHC [76, 97, 98]. While  $W^*\gamma^*$  and  $W^*Z^* \rightarrow 3\ell$  backgrounds proved most daunting for the Tevatron, at LHC the dominant background comes from  $t\bar{t}$  production [98].

### 4.3 Collimated dilepton +MET signature from $\tilde{Z}_1\tilde{Z}_2$ production

We first investigate the  $pp \rightarrow \tilde{Z}_1\tilde{Z}_2 \rightarrow \ell^+\ell^- + MET$  signal against the following SM backgrounds:

- $W^+W^-$  production (including  $WW \rightarrow \tau^+\tau^-$ ),
- $t\bar{t}$  production,
- $\gamma^* \rightarrow \ell^+\ell^-$  (Drell-Yan) production,
- $Z + jets$  with  $Z \rightarrow \tau^+\tau^-$  (tau pair) production,
- $\gamma^*Z$  production, where  $Z \rightarrow \nu_\ell\bar{\nu}_\ell$  and  $\gamma^* \rightarrow \ell^+\ell^-$  or  $\tau^+\tau^-$ .

We generate sparticle production and decay events at parton level using Isajet [71, 72] in Les Houches Event (LHE) format, and then feed the LHE files into Pythia [99] for initial/final state radiation, hadronization and underlying event. All backgrounds are generated with Pythia except  $\gamma^*Z^*$  which is generated by Madgraph/MadEvent [100]. The

collider events are then fed into the PGS toy detector simulation program [101]. Jets are found using an anti- $k_T$  jet finding algorithm with cone size  $\Delta R = 0.5$ . Leptons are classified as isolated if they contain less than 5 GeV hadronic activity in a cone of  $\Delta R = 0.2$  about the lepton direction.

Since the leptons from  $\tilde{Z}_1\tilde{Z}_2$  production are expected to be quite soft, we will focus initially upon the case of dimuon production, since muons can be identified more easily than electrons at very low  $p_T$ . Signal and background cross sections before and after cuts are listed in table 2. We first require:

- two opposite-sign muons: one with  $p_T(\mu_1) > 15$  GeV and  $|\eta(\mu_1)| < 0.9$  (central region), while the other has  $p_T(\mu_2) > 5$  GeV with  $|\eta(\mu_2)| < 2.4$ .

To reduce the large background from Drell-Yan dimuon production, we next impose

- $MET > 25$  GeV,

since  $MET$  in the DY case only arises from particles lost along the beam-line or cracks, or from energy mis-measurement, mainly from hadron radiation. There is also a large background from  $t\bar{t}$  production of dimuons and a background from single top in association with a W-boson which has a hard b-jet, but this always comes along with two hard b-jets from the  $t \rightarrow bW$  decays. Thus, we also require the number of jets

- $n(jets) = 0$ ,

where jets are identified as a cluster of hadrons with  $p_T(jet) > 15$  GeV,  $|\eta(jet)| < 2.4$ . At this stage, the largest background comes from  $W^+W^-$  production, which yields a continuum distribution in dimuon invariant mass  $m(\mu^+\mu^-)$ , whilst the signal dimuons are restricted to  $m(\mu^+\mu^-) < m_{\tilde{Z}_2} - m_{\tilde{Z}_1}$  which is just 16.2 GeV for HW150. Thus, we require

- $m(\mu^+\mu^-) < 20$  GeV .


Signal and BG after these cuts are listed for HW150 in table 2.

From table 2, we see that the dimuon signal comes about 30% from  $\tilde{Z}_1\tilde{Z}_2$  production, and about 70% from  $\tilde{W}_1\tilde{Z}_2$  production. In the latter case, the  $\tilde{W}_1$  usually decays to  $q\bar{q}'\tilde{Z}_1$  but with very low energy release, which sometimes escapes the “no-jet” cut. Meanwhile, the dominant remaining background comes from tau-pair, Drell-Yan and  $W^+W^-$  production. The remaining signal is 0.43 fb, while the summed SM background is  $\sim 11.9$  fb. The  $5\sigma$  discovery cross section for LHC7, assuming  $10 \text{ fb}^{-1}$  of integrated luminosity is 5.45 fb, so that the case of HW150 is far below this limit.

The inclusive muon  $p_T$  distribution before cuts is shown in figure 8 for the HW150 benchmark. Here, we see that the spectrum from HW150 benchmark is very soft, with the bulk of the distribution below 15 GeV. Thus, few of the signal events escape even the first cut listed above on  $p_T(\mu) > 15$  GeV. The signal rates for HW150 after cuts only corresponds to four events in  $10 \text{ fb}^{-1}$  of integrated luminosity, while SM background lies at the  $\sim 120$  event level.

process	$\sigma$ (fb)	$\sigma$ (after cuts, fb)
$\widetilde{W}_1 \widetilde{Z}_2$	313	0.3
$\widetilde{Z}_1 \widetilde{Z}_2$	192	0.13
$\gamma^* \rightarrow \mu^+ \mu^-$ (DY)	$1.1 \times 10^6$	4
$W^+ W^- \rightarrow \mu^+ \mu^-$	235.5	2.3
$\gamma^* Z \rightarrow \mu^+ \mu^- \nu_i \bar{\nu}_i$	6.8	0.3
$\gamma^*, Z \rightarrow \tau^+ \tau^- \rightarrow \mu^+ \mu^-$	$1.5 \times 10^4$	5
$t\bar{t} \rightarrow \mu^+ \mu^-$	$8.9 \times 10^4$	$< 0.3$

**Table 2.** Signal and BG cross sections in fb before and after cuts at LHC7. The signal rates are for higgsino-world benchmark point HW150. Each background process requires  $p_T(\mu) > 5$  GeV.




**Figure 8.** Distribution in  $p_T(\mu)$  from  $pp \rightarrow \widetilde{Z}_1 \widetilde{Z}_2 \rightarrow \mu^+ \mu^- + E_T^{\text{miss}}$  events at LHC from higgsino-world benchmark point HW150.

Likewise, trilepton signatures from  $\widetilde{W}_1^\pm \widetilde{Z}_2$  production yield a very soft isolated lepton spectrum, and are also difficult to extract at an observable level.<sup>3</sup> The search for jets+ $MET$  from higgsino pair production also yields a very soft jet and MET spectrum, and is difficult to extract from prodigious SM backgrounds.

<sup>3</sup>The reach of LHC14 for clean trileptons from  $\widetilde{W}_1 \widetilde{Z}_2$  production using  $100 \text{ fb}^{-1}$  of integrated luminosity has been calculated in ref. [98] in the Focus Point region of the mSUGRA model, where the thermal abundance  $\Omega_{\widetilde{Z}_1} h^2$  is required to be 0.11. This region contains  $\widetilde{Z}_1$  of mixed bino-higgsino variety with a larger  $m_{\widetilde{W}_1} - m_{\widetilde{Z}_1}$  mass gap than in the higgsino-world case, and corresponds to the light blue region of our figure 4. The LHC14 clean trilepton reach extended to  $m_{\widetilde{W}_1} \sim 260$  GeV, corresponding to  $m_{1/2} \sim 500$  GeV. Thus, in figure 4, we would expect the LHC14 reach via clean trileptons with  $100 \text{ fb}^{-1}$  to cover the light blue region for  $m_{1/2} \lesssim 0.5$  TeV.






**Figure 9.** Contours of  $2m_{\tilde{W}_1}$  in the  $\mu$  versus  $m_{1/2}$  parameter plane for other SUSY parameters as in HW150 scenario.

Thus, higgsino-world SUSY seems capable of eluding standard SUSY searches via isolated multi-leptons. The key feature of the HW scenario is that possibly observable levels of dimuon and trimuon production can occur, but at very low  $p_T$  levels. Our studies then motivate our experimental colleagues to push for dimuon and trimuon analyses at the very lowest  $p_T(\mu)$  levels in order to extract a possible signal.

## 5 Prospects for ILC or a muon collider

We have seen that the LHC has essentially no reach for the HW SUSY scenario due to a very soft spectrum of observable sparticle decay products. However, we have seen that the HW scenario mainly occurs for  $\mu \lesssim 250$  GeV (for  $m_{1/2} \lesssim 1$  TeV), which also corresponds to  $m_{\tilde{W}_1} \lesssim 250$  GeV. Contours of  $2m_{\tilde{W}_1}$  are shown in figure 9, where we see that the region with  $2m_{\tilde{W}_1} \lesssim 500$  GeV covers almost all of HW parameter space (compare against figure 2). For this mass range, chargino pair production, and also  $\tilde{Z}_1\tilde{Z}_2$  production, should be within range of the proposed international Linear Collider (ILC), which is proposed to operate initially at an energy  $\sqrt{s} = 500$  GeV. Chargino pair production would also be accessible to higher energy  $e^+e^-$  colliders like CLIC, or a muon collider (MC) operating in the TeV regime.


In figure 10, we show the cross sections for  $e^+e^- \rightarrow \tilde{W}_1^+\tilde{W}_1^-$  and  $e^+e^- \rightarrow \tilde{Z}_1\tilde{Z}_2$  using SUSY parameters as in the HW150 benchmark, but with  $\mu$  varying from 100-250 GeV. The variation in  $\mu$  causes  $m_{\tilde{W}_1}$  to vary, and in fact  $m_{\tilde{W}_1} \sim \mu$ , so that our results are plotted versus the more physical  $m_{\tilde{W}_1}$  value. We take  $\sqrt{s} = 500$  GeV. We see that over



**Figure 10.** Cross sections for chargino pair production and neutralino pair production versus  $m_{\tilde{W}_1}$  at a  $\sqrt{s} = 500$  GeV ILC or MC collider. We take SUSY parameters as in figure 7, and vary  $\mu$  to give variation in  $m_{\tilde{W}_1}$ .

most of HW parameter space, the chargino pair production cross section is in the several hundred fb range, until  $m_{\tilde{W}_1}$  approaches the kinematic limit for pair production. Chargino pair production will be signaled at ILC or MC by 1) soft multijet +  $\cancel{E}$  production, 2) soft isolated lepton plus jets +  $\cancel{E}$  production and 3) dilepton +  $\cancel{E}$  production, depending on whether the charginos decay leptonically or hadronically. These signatures should be easily visible against SM backgrounds such as  $WW$  production via distributions such as “missing mass”:  $\eta_h = \sqrt{\cancel{E}^2 - p^2}$  [109, 110]. In addition, SM backgrounds such as dilepton or dijet production from the  $\gamma\gamma$  initial state will contain energy depositions all in the same plane, while the SUSY signal will contain acoplanar events. Thus, the HW scenario should be easily visible at ILC, or a higher energy muon collider, even though it is difficult to see at LHC.

A distinctive feature of the HW scenario is that the  $\tilde{W}_1$ ,  $\tilde{Z}_1$  and  $\tilde{Z}_2$  are all mainly higgsino-like, whereas in models such as mSUGRA, these states are almost always gaugino-like. In ref. [109, 110], it is shown that for wino-like  $\tilde{W}_1$  and  $\tilde{Z}_2$ , the  $\tilde{W}_1^+\tilde{W}_1^-$  and  $\tilde{Z}_1\tilde{Z}_2$  production cross sections are steeply increasing functions of the electron beam polarization  $P_L(e^-)$  (where  $P_L(e^-) \sim -1$  corresponds to pure right-polarized  $e^-$ ,  $P_L(e^-) = +1$  corresponds to pure left-polarized  $e^-$ , and  $P_L(e^-) = 0$  corresponds to unpolarized  $e^-$  beams). In figure 11, we plot the  $e^+e^- \rightarrow \tilde{W}_1^+\tilde{W}_1^-$  and  $\tilde{Z}_1\tilde{Z}_2$  cross sections versus  $P_L(e^-)$  for the HW150 benchmark. In the HW scenario,  $\tilde{W}_1^+\tilde{W}_1^-$  production only increases by a factor of  $\sim 3.5$  as  $P_L(e^-)$  varies from -1 to +1, whereas in mSUGRA it typically increases by factors of about 100 [109, 110]. In addition, the  $\tilde{Z}_1\tilde{Z}_2$  cross section for HW150 is nearly



**Figure 11.** Cross sections for chargino pair production and neutralino pair production versus  $P_L(e^-)$  at a  $\sqrt{s} = 500$  GeV ILC collider. We take SUSY parameters as in HW1, with  $\mu = 150$  GeV.

flat versus  $P_L(e^-)$ , while in mSUGRA, it is typically increasing by factors of 20-30. Thus, variability of the SUSY production cross sections versus beam polarization will quickly allow one to extract much of the gaugino/higgsino content of the charginos/neutralinos which are accessible to an ILC with adjustable beam polarization.

## 6 Summary and conclusions

The higgsino-world SUSY scenario with multi-TeV scalars,  $\mu \lesssim 250$  GeV and intermediate scale gauginos is very appealing in that it can reconcile a decoupling solution to the SUSY flavor, CP,  $p$ -decay and gravitino problems with apparently low levels of naturalness or electroweak fine-tuning. The scenario is characterized by a mass hierarchy  $|\mu| \ll m_{1/2} \ll m_0$ , where  $m_0$  is the GUT scale mass of matter scalars. The HW scenario is most easily realized in models with non-universal Higgs masses, where the weak scale values of  $\mu$  and  $m_A$  are taken as free parameters. In the HW scenario, the  $\widetilde{W}_1$ ,  $\widetilde{Z}_1$  and  $\widetilde{Z}_2$  states are all light with mass  $\lesssim 250$  GeV, and dominantly higgsino-like. The remaining sparticles may well be heavy and inaccessible to LHC searches.

The standard thermal abundance of higgsino-like  $\widetilde{Z}_1$  particles is well below WMAP-measured values. However, in appealing cosmological scenarios such as those containing TeV-scale scalar fields such as moduli, or in scenarios with mixed axion- $\widetilde{Z}_1$  cold dark matter, the neutralino abundance can be easily pushed up into the measured range. If this is so, then there are excellent prospects for direct or indirect detection of higgsino-like relic WIMPs, and we expect experiments such as Xenon-100 or Xenon-1-ton to fully explore this possibility. Alternatively, in DM models such as the mixed  $a\widetilde{Z}_1$  scenario [54], it is

also possible to tune PQ parameters such that the WIMP abundance remains tiny, while the bulk of CDM is comprised of axions. Thus, the HW scenario will not be completely excludable by direct or indirect WIMP search experiments if no signals for WIMPs are seen.

At the LHC, gluino and squark production may be suppressed by large values of  $m_{\tilde{g}}$  and especially  $m_{\tilde{q}}$ . The  $\widetilde{W}_1\widetilde{Z}_1$ ,  $\widetilde{W}_1\widetilde{Z}_2$ ,  $\widetilde{Z}_1\widetilde{Z}_2$  and  $\widetilde{W}_1^+\widetilde{W}_1^-$  production reactions are then dominant, but are difficult to detect at LHC due to the small  $\widetilde{W}_1 - \widetilde{Z}_1$  and  $\widetilde{Z}_2 - \widetilde{Z}_1$  mass gaps, which lead to very soft visible particle production. The reaction  $pp \rightarrow \widetilde{Z}_1\widetilde{Z}_2$  may lead to tightly collimated OS/SF dilepton pairs, although calculations of signal and background after simple cuts indicate these occur at unobservable levels. Trileptons from  $\widetilde{W}_1\widetilde{Z}_2$  production are also difficult to see due to the soft spectrum of isolated leptons. Our studies should motivate our experimental colleagues to push for di- and tri-muon analyses at the very lowest levels of  $p_T(\mu)$  which are possible.

A linear  $e^+e^-$  collider such as ILC or a  $\mu^+\mu^-$  collider operating with  $\sqrt{s} \sim 0.5 - 1$  TeV should be able to make a thorough search for the HW scenario. If HW SUSY is discovered at ILC, then it should be possible to extract the gaugino/higgsino content of the  $\widetilde{W}_1$ ,  $\widetilde{Z}_2$  and  $\widetilde{Z}_1$  states using various kinematic and angular distributions along with beam polarization. Thus, the HW scenario provides a concrete realization of a SUSY construct which may well remain hidden from LHC and dark matter searches, but which is fully testable at a TeV-scale lepton collider.

## Acknowledgments

This work was supported in part by the U.S. Department of Energy under grant No. DE-FG02-04ER41305 and DE-FG02-95ER40896. VB thanks the KITP-Santa Barbara for its hospitality.

## References

- [1] S. Dimopoulos and H. Georgi, *Softly broken supersymmetry and SU(5)*, *Nucl. Phys.* **B 193** (1981) 150 [[INSPIRE](#)].
- [2] N. Sakai, *Naturalness in supersymmetric guts*, *Z. Phys.* **C 11** (1981) 153 [[INSPIRE](#)].
- [3] X. Tata, *Weak scale supersymmetry: from superfields to scattering events*, Cambridge University Press, Cambridge U.K. (2006).
- [4] M. Drees, R. Godbole and P. Roy, *Theory and phenomenology of sparticles*, World Scientific, Singapore (2004).
- [5] P. Binétruy, *Supersymmetry*, Oxford University Press, Oxford U.K. (2006).
- [6] S.P. Martin, *A Supersymmetry primer*, [hep-ph/9709356](#) [[INSPIRE](#)].
- [7] F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini, *A complete analysis of FCNC and CP constraints in general SUSY extensions of the standard model*, *Nucl. Phys.* **B 477** (1996) 321 [[hep-ph/9604387](#)] [[INSPIRE](#)].
- [8] H. Murayama and A. Pierce, *Not even decoupling can save minimal supersymmetric SU(5)*, *Phys. Rev.* **D 65** (2002) 055009 [[hep-ph/0108104](#)] [[INSPIRE](#)].

- [9] S. Weinberg, *Cosmological constraints on the scale of supersymmetry breaking*, *Phys. Rev. Lett.* **48** (1982) 1303 [INSPIRE].
- [10] M. Khlopov and A.D. Linde, *Is it easy to save the gravitino?*, *Phys. Lett. B* **138** (1984) 265 [INSPIRE].
- [11] M. Fukugita and T. Yanagida, *Baryogenesis without grand unification*, *Phys. Lett. B* **174** (1986) 45 [INSPIRE].
- [12] M. Luty, *Baryogenesis via leptogenesis*, *Phys. Rev. D* **45** (1992) 455 [INSPIRE].
- [13] W. Buchmüller and M. Plümacher, *Baryon asymmetry and neutrino mixing*, *Phys. Lett. B* **389** (1996) 73 [hep-ph/9608308] [INSPIRE].
- [14] W. Buchmüller and M. Plümacher, *Neutrino masses and the baryon asymmetry*, *Int. J. Mod. Phys. A* **15** (2000) 5047 [hep-ph/0007176] [INSPIRE].
- [15] R. Barbieri, P. Creminelli, A. Strumia and N. Tetradis, *Baryogenesis through leptogenesis*, *Nucl. Phys. B* **575** (2000) 61 [hep-ph/9911315] [INSPIRE].
- [16] G. Giudice, A. Notari, M. Raidal, A. Riotto and A. Strumia, *Towards a complete theory of thermal leptogenesis in the SM and MSSM*, *Nucl. Phys. B* **685** (2004) 89 [hep-ph/0310123] [INSPIRE].
- [17] W. Buchmüller, R. Peccei and T. Yanagida, *Leptogenesis as the origin of matter*, *Ann. Rev. Nucl. Part. Sci.* **55** (2005) 311 [hep-ph/0502169] [INSPIRE].
- [18] W. Buchmüller, P. Di Bari and M. Plümacher, *Cosmic microwave background, matter-antimatter asymmetry and neutrino masses*, *Nucl. Phys. B* **643** (2002) 367 [Erratum *ibid.* **B 793** (2008) 362] [hep-ph/0205349] [INSPIRE].
- [19] W. Buchmüller, P. Di Bari and M. Plümacher, *Leptogenesis for pedestrians*, *Annals Phys.* **315** (2005) 305 [hep-ph/0401240] [INSPIRE].
- [20] W. Buchmüller, P. Di Bari and M. Plümacher, *Some aspects of thermal leptogenesis*, *New J. Phys.* **6** (2004) 105.
- [21] G. Lazarides and Q. Shafi, *Origin of matter in the inflationary cosmology*, *Phys. Lett. B* **258** (1991) 305 [INSPIRE].
- [22] K. Kumekawa, T. Moroi and T. Yanagida, *Flat potential for inflaton with a discrete  $R$  invariance in supergravity*, *Prog. Theor. Phys.* **92** (1994) 437 [hep-ph/9405337] [INSPIRE].
- [23] T. Asaka, K. Hamaguchi, M. Kawasaki and T. Yanagida, *Leptogenesis in inflaton decay*, *Phys. Lett. B* **464** (1999) 12 [hep-ph/9906366] [INSPIRE].
- [24] G. Giudice, M. Peloso, A. Riotto and I. Tkachev, *Production of massive fermions at preheating and leptogenesis*, *JHEP* **08** (1999) 014 [hep-ph/9905242] [INSPIRE].
- [25] M. Dine, A. Kagan and S. Samuel, *Naturalness in supersymmetry, or raising the supersymmetry breaking scale*, *Phys. Lett. B* **243** (1990) 250 [INSPIRE].
- [26] A.G. Cohen, D. Kaplan and A. Nelson, *The more minimal supersymmetric standard model*, *Phys. Lett. B* **388** (1996) 588 [hep-ph/9607394] [INSPIRE].
- [27] H. Baer, S. Kraml, A. Lessa, S. Sekmen and X. Tata, *Effective supersymmetry at the LHC*, *JHEP* **10** (2010) 018 [arXiv:1007.3897] [INSPIRE].
- [28] M. Kawasaki, K. Kohri, T. Moroi and A. Yotsuyanagi, *Big-bang nucleosynthesis and gravitino*, *Phys. Rev. D* **78** (2008) 065011 [arXiv:0804.3745] [INSPIRE].

- [29] H. Baer and X. Tata, *Weak scale supersymmetry: from superfields to scattering events*, Cambridge University Press, Cambridge U.K. (2006).
- [30] R. Barbieri and G. Giudice, *Upper bounds on supersymmetric particle masses*, *Nucl. Phys. B* **306** (1988) 63 [INSPIRE].
- [31] G.W. Anderson and D.J. Castano, *Measures of fine tuning*, *Phys. Lett. B* **347** (1995) 300 [hep-ph/9409419] [INSPIRE].
- [32] K.L. Chan, U. Chattopadhyay and P. Nath, *Naturalness, weak scale supersymmetry and the prospect for the observation of supersymmetry at the Tevatron and at the CERN LHC*, *Phys. Rev. D* **58** (1998) 096004 [hep-ph/9710473] [INSPIRE].
- [33] J.L. Feng, K.T. Matchev and T. Moroi, *Multi-TeV scalars are natural in minimal supergravity*, *Phys. Rev. Lett.* **84** (2000) 2322 [hep-ph/9908309] [INSPIRE].
- [34] J.L. Feng, K.T. Matchev and T. Moroi, *Focus points and naturalness in supersymmetry*, *Phys. Rev. D* **61** (2000) 075005 [hep-ph/9909334] [INSPIRE].
- [35] G.L. Kane, *A higgsino-LSP world*, in *Perspectives on supersymmetry*, G.L. Kane ed., World Scientific, Singapore (1998).
- [36] G.L. Kane and J.D. Wells, *Higgsino cold dark matter motivated by collider data*, *Phys. Rev. Lett.* **76** (1996) 4458 [hep-ph/9603336] [INSPIRE].
- [37] WMAP collaboration, E. Komatsu et al., *Seven-year Wilkinson Microwave Anisotropy Probe (WMAP) observations: cosmological interpretation*, *Astrophys. J. Suppl.* **192** (2011) 18 [arXiv:1001.4538] [INSPIRE].
- [38] J.R. Ellis and K.A. Olive, *How finely tuned is supersymmetric dark matter?*, *Phys. Lett. B* **514** (2001) 114 [hep-ph/0105004] [INSPIRE].
- [39] H. Baer and A.D. Box, *Fine-tuning favors mixed axion/axino cold dark matter over neutralinos in the minimal supergravity model*, *Eur. Phys. J. C* **68** (2010) 523 [arXiv:0910.0333] [INSPIRE].
- [40] H. Baer, A.D. Box and H. Summy, *Neutralino versus axion/axino cold dark matter in the 19 parameter SUGRA model*, *JHEP* **10** (2010) 023 [arXiv:1005.2215] [INSPIRE].
- [41] B.S. Acharya, G. Kane, S. Watson and P. Kumar, *A non-thermal WIMP miracle*, *Phys. Rev. D* **80** (2009) 083529 [arXiv:0908.2430] [INSPIRE].
- [42] T. Moroi and L. Randall, *Wino cold dark matter from anomaly mediated SUSY breaking*, *Nucl. Phys. B* **570** (2000) 455 [hep-ph/9906527] [INSPIRE].
- [43] G.B. Gelmini and P. Gondolo, *Neutralino with the right cold dark matter abundance in (almost) any supersymmetric model*, *Phys. Rev. D* **74** (2006) 023510 [hep-ph/0602230] [INSPIRE].
- [44] G. Gelmini, P. Gondolo, A. Soldatenko and C.E. Yaguna, *The effect of a late decaying scalar on the neutralino relic density*, *Phys. Rev. D* **74** (2006) 083514 [hep-ph/0605016] [INSPIRE].
- [45] R. Peccei and H.R. Quinn, *CP conservation in the presence of instantons*, *Phys. Rev. Lett.* **38** (1977) 1440 [INSPIRE].
- [46] R. Peccei and H.R. Quinn, *Constraints imposed by CP conservation in the presence of instantons*, *Phys. Rev. D* **16** (1977) 1791 [INSPIRE].
- [47] S. Weinberg, *A new light boson?*, *Phys. Rev. Lett.* **40** (1978) 223 [INSPIRE].

- [48] F. Wilczek, *Problem of strong  $p$  and  $t$  invariance in the presence of instantons*, *Phys. Rev. Lett.* **40** (1978) 279 [INSPIRE].
- [49] J.E. Kim, *Weak interaction singlet and strong CP invariance*, *Phys. Rev. Lett.* **43** (1979) 103 [INSPIRE].
- [50] M.A. Shifman, A. Vainshtein and V.I. Zakharov, *Can confinement ensure natural CP invariance of strong interactions?*, *Nucl. Phys.* **B 166** (1980) 493 [INSPIRE].
- [51] M. Dine, W. Fischler and M. Srednicki, *A simple solution to the strong CP problem with a harmless axion*, *Phys. Lett.* **B 104** (1981) 199 [INSPIRE].
- [52] A.P. Zhitnitskii, *On possible suppression of the axion hadron interactions* (In Russian), *Sov. J. Nucl. Phys.* **31** (1980) 260 [*Yad. Fiz.* **31** (1980) 497] [INSPIRE].
- [53] K.-Y. Choi, J.E. Kim, H.M. Lee and O. Seto, *Neutralino dark matter from heavy axino decay*, *Phys. Rev.* **D 77** (2008) 123501 [arXiv:0801.0491] [INSPIRE].
- [54] H. Baer, A. Lessa, S. Rajagopalan and W. Sreethawong, *Mixed axion/neutralino cold dark matter in supersymmetric models*, *JCAP* **06** (2011) 031 [arXiv:1103.5413] [INSPIRE].
- [55] M. Bolz, A. Brandenburg and W. Buchmüller, *Thermal production of gravitinos*, *Nucl. Phys.* **B 606** (2001) 518 [Erratum *ibid.* **B 790** (2008) 336-337] [hep-ph/0012052] [INSPIRE].
- [56] J. Pradler and F.D. Steffen, *Thermal gravitino production and collider tests of leptogenesis*, *Phys. Rev.* **D 75** (2007) 023509 [hep-ph/0608344] [INSPIRE].
- [57] V.S. Rychkov and A. Strumia, *Thermal production of gravitinos*, *Phys. Rev.* **D 75** (2007) 075011 [hep-ph/0701104] [INSPIRE].
- [58] K. Kohri, M. Yamaguchi and J. Yokoyama, *Neutralino dark matter from heavy gravitino decay*, *Phys. Rev.* **D 72** (2005) 083510 [hep-ph/0502211] [INSPIRE].
- [59] T. Asaka, S. Nakamura and M. Yamaguchi, *Gravitinos from heavy scalar decay*, *Phys. Rev.* **D 74** (2006) 023520 [hep-ph/0604132] [INSPIRE].
- [60] M. Endo, F. Takahashi and T. Yanagida, *Inflaton decay in supergravity*, *Phys. Rev.* **D 76** (2007) 083509 [arXiv:0706.0986] [INSPIRE].
- [61] H. Baer, R. Dermisek, S. Rajagopalan and H. Summy, *Neutralino, axion and axino cold dark matter in minimal, hypercharged and gaugino AMSB*, *JCAP* **07** (2010) 014 [arXiv:1004.3297] [INSPIRE].
- [62] J.R. Ellis, K.A. Olive and Y. Santoso, *The MSSM parameter space with nonuniversal Higgs masses*, *Phys. Lett.* **B 539** (2002) 107 [hep-ph/0204192] [INSPIRE].
- [63] J.R. Ellis, T. Falk, K.A. Olive and Y. Santoso, *Exploration of the MSSM with nonuniversal Higgs masses*, *Nucl. Phys.* **B 652** (2003) 259 [hep-ph/0210205] [INSPIRE].
- [64] H. Baer, A. Mustafayev, S. Profumo, A. Belyaev and X. Tata, *Neutralino cold dark matter in a one parameter extension of the minimal supergravity model*, *Phys. Rev.* **D 71** (2005) 095008 [hep-ph/0412059] [INSPIRE].
- [65] H. Baer, A. Mustafayev, S. Profumo, A. Belyaev and X. Tata, *Direct, indirect and collider detection of neutralino dark matter in SUSY models with non-universal Higgs masses*, *JHEP* **07** (2005) 065 [hep-ph/0504001] [INSPIRE].
- [66] H. Baer and X. Tata, *Probing charginos and neutralinos beyond the reach of LEP at the Tevatron collider*, *Phys. Rev.* **D 47** (1993) 2739 [INSPIRE].

- [67] H. Baer, M. Drees, F. Paige, P. Quintana and X. Tata, *Trilepton signal for supersymmetry at the Fermilab Tevatron revisited*, *Phys. Rev. D* **61** (2000) 095007 [[hep-ph/9906233](#)] [[INSPIRE](#)].
- [68] V.D. Barger and C. Kao, *Trilepton signature of minimal supergravity at the upgraded Tevatron*, *Phys. Rev. D* **60** (1999) 115015 [[hep-ph/9811489](#)] [[INSPIRE](#)].
- [69] K.T. Matchev and D.M. Pierce, *New backgrounds in trilepton, dilepton and dilepton plus  $\tau$  jet SUSY signals at the Tevatron*, *Phys. Lett. B* **467** (1999) 225 [[hep-ph/9907505](#)] [[INSPIRE](#)].
- [70] SUGRA WORKING GROUP collaboration, V. Barger et al., *Report of the SUGRA working group for Run II of the Tevatron*, [hep-ph/0003154](#) [[INSPIRE](#)].
- [71] F.E. Paige, S.D. Protopopescu, H. Baer and X. Tata, *ISAJET 7.69: a Monte Carlo event generator for  $pp$ ,  $\bar{p}p$  and  $e^+e^-$  reactions*, [hep-ph/0312045](#) [[INSPIRE](#)].
- [72] H. Baer, J. Ferrandis, S. Kraml and W. Porod, *On the treatment of threshold effects in SUSY spectrum computations*, *Phys. Rev. D* **73** (2006) 015010 [[hep-ph/0511123](#)] [[INSPIRE](#)].
- [73] H. Baer, V. Barger, A. Lessa and X. Tata, *Capability of LHC to discover supersymmetry with  $\sqrt{s} = 7$  TeV and  $1 \text{ fb}^{-1}$* , *JHEP* **06** (2010) 102 [[arXiv:1004.3594](#)] [[INSPIRE](#)].
- [74] H. Baer, X. Tata and J. Woodside, *Multi-lepton signals from supersymmetry at hadron super colliders*, *Phys. Rev. D* **45** (1992) 142 [[INSPIRE](#)].
- [75] H. Baer, C.-h. Chen, F. Paige and X. Tata, *Signals for minimal supergravity at the CERN Large Hadron Collider: multi-jet plus missing energy channel*, *Phys. Rev. D* **52** (1995) 2746 [[hep-ph/9503271](#)] [[INSPIRE](#)].
- [76] H. Baer, C.-h. Chen, F. Paige and X. Tata, *Signals for minimal supergravity at the CERN Large Hadron Collider. 2: multi-lepton channels*, *Phys. Rev. D* **53** (1996) 6241 [[hep-ph/9512383](#)] [[INSPIRE](#)].
- [77] H. Baer, C.-H. Chen, M. Drees, F. Paige and X. Tata, *Probing minimal supergravity at the CERN LHC for large  $\tan\beta$* , *Phys. Rev. D* **59** (1999) 055014 [[hep-ph/9809223](#)] [[INSPIRE](#)].
- [78] H. Baer, C. Balázs, A. Belyaev, T. Krupovnickas and X. Tata, *Updated reach of the CERN LHC and constraints from relic density,  $b \rightarrow s\gamma$  and  $a(\mu)$  in the mSUGRA model*, *JHEP* **06** (2003) 054 [[hep-ph/0304303](#)] [[INSPIRE](#)].
- [79] S. Abdullin and F. Charles, *Search for SUSY in (leptons +) jets +  $E_{\text{miss}}(T)$  final states*, *Nucl. Phys. B* **547** (1999) 60 [[hep-ph/9811402](#)] [[INSPIRE](#)].
- [80] CMS collaboration, S. Abdullin et al., *Discovery potential for supersymmetry in CMS*, *J. Phys. G* **28** (2002) 469 [[hep-ph/9806366](#)] [[INSPIRE](#)].
- [81] B. Allanach, J. Hetherington, M.A. Parker and B. Webber, *Naturalness reach of the Large Hadron Collider in minimal supergravity*, *JHEP* **08** (2000) 017 [[hep-ph/0005186](#)] [[INSPIRE](#)].
- [82] W. Beenakker, R. Hopker and M. Spira, *PROSPINO: a program for the production of supersymmetric particles in next-to-leading order QCD*, [hep-ph/9611232](#) [[INSPIRE](#)].
- [83] H. Baer, J.R. Ellis, G. Gelmini, D.V. Nanopoulos and X. Tata, *Squark decays into gauginos at the  $p\bar{p}$  collider*, *Phys. Lett. B* **161** (1985) 175 [[INSPIRE](#)].
- [84] G. Gamberini, *Heavy gluino and squark decays at  $p\bar{p}$  collider*, *Z. Phys. C* **30** (1986) 605 [[INSPIRE](#)].



- [85] H. Baer, V.D. Barger, D. Karatas and X. Tata, *Detecting gluinos at hadron supercolliders*, *Phys. Rev. D* **36** (1987) 96 [INSPIRE].
- [86] H. Baer and T. Krupovnickas, *Radiative neutralino decay in supersymmetric models*, *JHEP* **09** (2002) 038 [hep-ph/0208277] [INSPIRE].
- [87] XENON100 collaboration, E. Aprile et al., *Dark matter results from 100 live days of XENON100 data*, *Phys. Rev. Lett.* **107** (2011) 131302 [arXiv:1104.2549] [INSPIRE].
- [88] H. Baer, A. Belyaev, T. Krupovnickas and J. O’Farrill, *Indirect, direct and collider detection of neutralino dark matter*, *JCAP* **08** (2004) 005 [hep-ph/0405210] [INSPIRE].
- [89] H. Baer, E.-K. Park and X. Tata, *Collider, direct and indirect detection of supersymmetric dark matter*, *New J. Phys.* **11** (2009) 105024 [arXiv:0903.0555] [INSPIRE].
- [90] H. Baer and S. Profumo, *Low energy antideuterons: shedding light on dark matter*, *JCAP* **12** (2005) 008 [astro-ph/0510722] [INSPIRE].
- [91] FERMI LAT collaboration, I. Kuznetsova and J. Rafelski, *Electron-positron plasma drop formed by ultra-intense laser pulses*, arXiv:1109.3546 [INSPIRE].
- [92] K. Rajagopal, M.S. Turner and F. Wilczek, *Cosmological implications of axinos*, *Nucl. Phys. B* **358** (1991) 447 [INSPIRE].
- [93] L. Covi, J.E. Kim and L. Roszkowski, *Axinos as cold dark matter*, *Phys. Rev. Lett.* **82** (1999) 4180 [hep-ph/9905212] [INSPIRE].
- [94] L. Covi, H.-B. Kim, J.E. Kim and L. Roszkowski, *Axinos as dark matter*, *JHEP* **05** (2001) 033 [hep-ph/0101009] [INSPIRE].
- [95] H. Baer, A.D. Box and H. Summy, *Mainly axion cold dark matter in the minimal supergravity model*, *JHEP* **08** (2009) 080 [arXiv:0906.2595] [INSPIRE].
- [96] F.D. Steffen, *Dark matter candidates-axions, neutralinos, gravitinos and axinos*, *Eur. Phys. J. C* **59** (2009) 557 [arXiv:0811.3347] [INSPIRE].
- [97] H. Baer, C.-h. Chen, F. Paige and X. Tata, *Trileptons from chargino-neutralino production at the CERN Large Hadron Collider*, *Phys. Rev. D* **50** (1994) 4508 [hep-ph/9404212] [INSPIRE].
- [98] H. Baer, T. Krupovnickas, S. Profumo and P. Ullio, *Model independent approach to focus point supersymmetry: from dark matter to collider searches*, *JHEP* **10** (2005) 020 [hep-ph/0507282] [INSPIRE].
- [99] T. Sjöstrand, S. Mrenna and P.Z. Skands, *PYTHIA 6.4 physics and manual*, *JHEP* **05** (2006) 026 [hep-ph/0603175] [INSPIRE].
- [100] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer and T. Stelzer, *MadGraph 5: going beyond*, *JHEP* **06** (2011) 128 [arXiv:1106.0522] [INSPIRE].
- [101] J. Conway, *PGS — Pretty Good Simulation*, online at <http://physics.ucdavis.edu/~conway/research/software/pgs/pgs4-general.htm>.
- [102] H. Baer, K. Hagiwara and X. Tata, *Gauginos as a signal for supersymmetry at  $p\bar{p}$  colliders*, *Phys. Rev. D* **35** (1987) 1598 [INSPIRE].
- [103] H. Baer, D.D. Karatas and X. Tata, *Gluino and squark production in association with gauginos at hadron supercolliders*, *Phys. Rev. D* **42** (1990) 2259 [INSPIRE].

- [104] H. Baer, C. Kao and X. Tata, *Aspects of chargino-neutralino production at the Tevatron collider*, *Phys. Rev. D* **48** (1993) 5175 [[hep-ph/9307347](#)] [[INSPIRE](#)].
- [105] I. Hinchliffe, F. Paige, M. Shapiro, J. Soderqvist and W. Yao, *Precision SUSY measurements at CERN LHC*, *Phys. Rev. D* **55** (1997) 5520 [[hep-ph/9610544](#)] [[INSPIRE](#)].
- [106] H. Bachacou, I. Hinchliffe and F.E. Paige, *Measurements of masses in SUGRA models at CERN LHC*, *Phys. Rev. D* **62** (2000) 015009 [[hep-ph/9907518](#)] [[INSPIRE](#)].
- [107] ATLAS collaboration, *Expected performance of the ATLAS experiment: detector, trigger and physics*, [CERN-OPEN-2008-020](#) (2009).
- [108] CMS Collaboration, *Physics technical design report. Volume II: physics performance*, [CERN-LHCC-2006-021](#) (2006).
- [109] H. Baer, R.B. Munroe and X. Tata, *Supersymmetry studies at future linear  $e^+e^-$  colliders*, *Phys. Rev. D* **54** (1996) 6735 [*Erratum ibid.* **D 56** (1997) 4424] [[hep-ph/9606325](#)] [[INSPIRE](#)].
- [110] H. Baer, A. Belyaev, T. Krupovnickas and X. Tata, *Linear collider capabilities for supersymmetry in dark matter allowed regions of the mSUGRA model*, *JHEP* **02** (2004) 007 [[hep-ph/0311351](#)] [[INSPIRE](#)].