

2008

Interpretation and evaluation of combined measurement techniques for soil CO₂ efflux: Discrete surface chambers and continuous soil CO₂ concentration probes

Diego Andrés Riveros-Iregui
University of Nebraska - Lincoln, driveros2@unl.edu

Brian L. McGlynn
Montana State University - Bozeman

Howard E. Epstein
University of Virginia, Charlottesville

Daniel L. Welsch
Canaan Valley Institute, Davis, West Virginia

Follow this and additional works at: <http://digitalcommons.unl.edu/natrespapers>

 Part of the [Natural Resources and Conservation Commons](#)

Riveros-Iregui, Diego Andrés; McGlynn, Brian L.; Epstein, Howard E.; and Welsch, Daniel L., "Interpretation and evaluation of combined measurement techniques for soil CO₂ efflux: Discrete surface chambers and continuous soil CO₂ concentration probes" (2008). *Papers in Natural Resources*. 211.
<http://digitalcommons.unl.edu/natrespapers/211>

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Natural Resources by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Interpretation and evaluation of combined measurement techniques for soil CO₂ efflux: Discrete surface chambers and continuous soil CO₂ concentration probes

Diego A. Riveros-Iregui,¹ Brian L. McGlynn,¹ Howard E. Epstein,² and Daniel L. Welsch³

Received 24 June 2008; revised 17 September 2008; accepted 27 October 2008; published 9 December 2008.

[1] Soil CO₂ efflux is a large respiratory flux from terrestrial ecosystems and a critical component of the global carbon (C) cycle. Lack of process understanding of the spatiotemporal controls on soil CO₂ efflux limits our ability to extrapolate from fluxes measured at point scales to scales useful for corroboration with other ecosystem level measures of C exchange. Additional complications are introduced by the effects of soil water content seasonality and rainfall on the performance of measurement techniques. In this paper we present measurements of soil CO₂ efflux made at two contrasting sites within a characteristic subalpine forest of the northern Rocky Mountains. Comparison of measurements between the soil respiration chamber technique and the soil CO₂ profile technique over daily and seasonal time scales indicated that soil water content plays a major role in the magnitude and seasonality of soil CO₂ efflux, especially after snowmelt or following summer rainfall. Agreement between both techniques was limited during high soil water content conditions and after summer rainfall. Differences in diel hysteresis patterns of soil CO₂ efflux between sites were controlled by the effects of canopy cover and temporal differences in photosynthetic activity of vegetation. Our results indicate that an accurate parameterization of soil water content heterogeneity in space and time must be a critical component of realistic model representations of soil CO₂ efflux from heterogeneous landscapes.

Citation: Riveros-Iregui, D. A., B. L. McGlynn, H. E. Epstein, and D. L. Welsch (2008), Interpretation and evaluation of combined measurement techniques for soil CO₂ efflux: Discrete surface chambers and continuous soil CO₂ concentration probes, *J. Geophys. Res.*, 113, G04027, doi:10.1029/2008JG000811.

1. Introduction

[2] Soil CO₂ efflux is a natural process by which soil carbon is released into the atmosphere through autotrophic and heterotrophic respiration. Evaluating and predicting soil CO₂ efflux response to differences in hydrologic conditions (e.g., groundwater recharge and discharge areas, soil water content, precipitation, and land cover) are largely constrained by the methods used to measure, interpret, and model soil CO₂ efflux. Rates of soil CO₂ efflux are currently estimated from a wide range of ecosystems with manual soil respiration chambers [Fang and Moncrieff, 1996; Subke and Tenhunen, 2004; Welsch and Hornberger, 2004], automated soil respiration chambers [Goulden and Crill, 1997; Savage and Davidson, 2001; Burrows et al., 2005], and the soil CO₂ profile technique [Tang et al., 2003; Jassal et al., 2005; Tang and Baldocchi, 2005]. Particularly in the last 5 years, the soil CO₂ profile technique has gained popularity because

it can provide continuous and automated measurements at temporal scales useful for comparison with other techniques of ecosystem C exchange such as eddy covariance towers [Baldocchi et al., 2006]. While a wealth of ongoing studies use either technique, a direct comparison of their performance, measurements, strengths, and limitations in space and time is lacking.

[3] It has been suggested that the interactions among precipitation, infiltration, evaporation, transpiration, and soil drainage exert a major control on vegetation activity in water-limited ecosystems [Ridolfi et al., 2000; Porporato et al., 2002]. Large gaps exist in our understanding of the variability of soil CO₂ efflux in response to changing hydrologic conditions across space and time. Traditionally, studies addressing the variability of soil CO₂ efflux focus on its temporal component (e.g., diel, seasonal, and yearly variability) but tend to omit the spatial component inherent to this flux (i.e., landscape-induced variability). This omission limits the capability of temperature-based models [Lloyd and Taylor, 1994] to accurately estimate soil CO₂ efflux from areas having different characteristics within similar ecosystems. More importantly, this omission restricts our understanding of how CO₂-producing processes simultaneously develop in space and time to generate the

¹Department of Land Resources and Environmental Sciences, Montana State University, Bozeman, Montana, USA.

²Department of Environmental Sciences, University of Virginia, Charlottesville, Virginia, USA.

³Canaan Valley Institute, Davis, West Virginia, USA.




Figure 1. Location of the Tenderfoot Creek Experimental Forest in the Little Belt Mountains of central Montana. The two most common ecosystems of these mountains were selected for this study: a wet riparian meadow and a dry upland forest.

soil CO₂ rates that chambers or soil profile techniques measure.

[4] Recent studies [Riveros-Iregui *et al.*, 2007] demonstrated that soil water content controls the relationship between soil CO₂ efflux and soil temperature, as soil water content (1) enhances soil CO₂ production and (2) inhibits soil CO₂ diffusion. Furthermore, the seasonality of soil water content can control the switch from diffusion- to production-limited soil CO₂ efflux [Riveros-Iregui *et al.*, 2007; Riveros-Iregui, 2008; Pacific *et al.*, 2008]. This concept becomes especially important in ecosystems with considerable spatial variability in soil water content induced by landscape morphology (i.e., convex versus concave areas [Scott-Denton *et al.*, 2003; Pacific *et al.*, 2008]). Vegetation cover and soil characteristics further control ecosystem response to changes in environmental conditions [Huxman *et al.*, 2004]. As a result, marked differences in soil water content regimes play a major role in ecosystem response, particularly soil CO₂ efflux, of heterogeneous forests. In this paper, we provide a comparison, over diel and seasonal time scales, of discrete (soil respiration chamber) and continuous (soil CO₂ profile technique) measurements of soil CO₂ efflux made at two sites: a wet riparian meadow and a dry upland forest. Both sites are collocated with eddy covariance towers and are within a characteristic subalpine forest of the northern Rocky Mountains. Through this space-time comparison we seek to (1) determine the mechanisms driving variability in soil CO₂ efflux from riparian meadows and upland forests; (2) compare the performance of soil respiration chambers and solid-state CO₂ probes throughout an entire growing season; and (3) assess, both mechanistically and methodologically, the confidence that these methods offer as providers of soil CO₂ efflux rates from heterogeneous landscapes. This information is essential to improving process understanding of soil CO₂ efflux from large areas, establishing a conceptual framework for soil CO₂ efflux

modeling studies, and adding confidence to current and developing measurement techniques.

2. Methods

2.1. Study Site

[5] This study was located in the Tenderfoot Creek Experimental Forest, in the Little Belt Mountains of central Montana (Figure 1). These mountains are characteristic of the subalpine forests of the northern Rocky Mountains. Two contrasting ecosystems that represent the two dominant systems of these mountains were selected to address the objectives of this study: a wet riparian meadow (hereafter riparian site) and a dry upland forest (hereafter upland site). Vegetation cover at the riparian site is predominantly *Calamagrostis canadensis* (bluejoint reedgrass), whereas the upland site is covered mostly by *Pinus contorta* (lodgepole pine) and *Pseudotsuga menziesii* (Douglas fir) in the overstory and *Vaccinium* spp. in the understory. Elevations are 2169 and 2305 m at the riparian and upland sites, respectively. Mean annual precipitation is 880 mm with ~70% falling as snow [Farnes *et al.*, 1995], and peak snowpack accumulations occur between late March and mid-April [Woods *et al.*, 2006]. Mean annual temperature is 0°C, and the growing season lasts from 45 to 75 days.

2.2. Environmental Variables

[6] Between 9 June 2006 and 7 September 2006 we measured volumetric soil water content (θ) (CSI model 616, Campbell Scientific Inc., Logan, Utah) and soil temperature (T_s) (CSI model 107, Campbell Scientific Inc., Logan, Utah) at 20 cm below the soil surface, at 20-min intervals, and collected the data with a logger (model CR-10x, Campbell Scientific Inc., Logan, Utah). Manual measurements ($n = 3$) of θ were also taken with a portable meter (Hydrosense, Campbell Scientific Inc., Utah, USA) to obtain an integrated estimate of soil water content over the top 20 cm of the soil profile. Measurements from the CSI Hydrosense meter were experimentally corroborated with regular time domain reflectometry instruments ($r^2 = 0.986$ and $n = 121$) to confirm applicability of the CSI Hydrosense instrument under a full range of soil water contents. Precipitation was measured by a tipping bucket rain gauge (TR-525M, accurate to within 1% for up to 50 mm h⁻¹, Texas Electronics, Dallas, Texas, USA), at 20-min intervals and reported on a daily basis.

2.3. Measurements of Soil CO₂ Efflux

[7] Soil CO₂ efflux was measured independently at each site by the soil respiration chamber technique (discrete measurements) and by the soil CO₂ profile technique (continuous measurements). While the performance of soil respiration chambers has been amply evaluated [Norman *et al.*, 1997; Hutchinson and Livingston, 2001; Davidson *et al.*, 2002], the performance of the relatively newer soil CO₂ profile technique remains to be critically evaluated in space and time, as well as against soil respiration chambers. Discrete measurements were based on a soil respiration chamber model SRC-1 (footprint of 314.2 cm², accuracy within 1% of calibrated range (0–9.99 g CO₂ m⁻² h⁻¹), PP Systems, Massachusetts, USA) equipped with an infrared gas analyzer (EGM-4, accuracy within 1% of calibrated

range (0–2000 ppm), PP Systems, Massachusetts, USA). Chamber measurements were collected in triplicate every 2–7 days at each site. At each site, a 0.5-m² area was roped off to minimize disturbance and vegetation was clipped once a week after measurements were collected. Roots were left intact to minimize disturbance to belowground respiration. Before each measurement, the chamber was flushed with ambient air for 15 s and placed onto the soil, ensuring a good seal between the chamber and the soil surface. Soil CO₂ efflux was calculated by measuring the rate of increase in CO₂ concentration within the chamber and fitting a quadratic equation to the relationship between the increasing CO₂ concentration and elapsed time (as recommended by the manufacturer). In order to minimize introduction of biases during sampling, no chamber measurements were taken before 1000 LT or after 1600 LT.

[8] Continuous soil CO₂ concentration measurements were collected with solid-state CO₂ probes (GMP221 with transmitter, Vaisala, Helsinki, Finland) installed at 5 cm below the soil surface, logging continuously at 20-min intervals with a data logger (model CR10x, Campbell Scientific Inc., Logan, Utah). Soil CO₂ concentrations measured by the probes were corrected for temperature and pressure following compensatory procedures described by *Tang et al.* [2003] and according to the manufacturer's recommendation. When buried in the soil, these probes respond to changes in CO₂ concentrations in less than 5 min [*Tang et al.*, 2003]. Because it is difficult to measure soil CO₂ concentrations near the soil-atmosphere interface (e.g., $z = 0$), we tested the sensitivity of soil CO₂ efflux to variability in atmospheric concentrations by simultaneously using three different CO₂ concentration values (350, 450, and 550 ppm) at the soil-atmosphere interface. We chose these three values on the basis of the range of variability of initial surface CO₂ concentrations measured at each deployment of the soil respiration chamber (ranging between ~390 and ~530 ppm). Our results demonstrate that the assumed values do not compromise calculation of soil CO₂ efflux, as the diel variability of soil CO₂ concentration at 5 cm is much greater than the diel variability of soil CO₂ above the soil surface given the atmospheric buffer. Previous studies [*Tang et al.*, 2005b] have used similar assumptions (~370 ppm at 0.5 m above the soil surface). However, our approach of using all three values provides a confidence error of 200 ppm (at $z = 0$), demonstrating that variations in concentrations at the soil surface introduce little variation in estimated soil CO₂ efflux given the natural diel variability of soil CO₂ at depth (>5000 ppm). Nevertheless, to illustrate this effect, confidence bounds (for 350 and 550 ppm) were estimated and presented with the results. Additional corroboration and confidence is given when comparing continuous and discrete chamber soil CO₂ efflux estimates (see section 3).

[9] Continuous soil CO₂ efflux based on solid-state probes has been estimated in numerous recent studies across multiple ecosystems [*Tang et al.*, 2003; *Jassal et al.*, 2005; *Tang et al.*, 2005a, 2005b; *Baldocchi et al.*, 2006; *Vargas and Allen*, 2008]. To estimate surface efflux, these studies assumed that flux of soil CO₂ was linear with depth, with increasing concentrations with increasing depth. However, these assumptions become invalid when soil CO₂ is greater in shallow soils than in deeper soils, leading to bidirectional concentration gradients and fluxes. This scenario can occur

in response to disturbances such as precipitation and shallow soil wetting [e.g., *Tang et al.*, 2005b]. Because precipitation was sporadic throughout the growing season in our system and we sought to investigate CO₂ flux toward the surface at both diel and seasonal time scales, we calculated soil CO₂ efflux based on concentrations at 0 and 0.05 m using Fick's first law of diffusion,

$$F = -D_P \frac{\partial[\text{CO}_2]}{\partial z}, \quad (1)$$

where F is CO₂ flux between two depths and D_P is the diffusion coefficient for CO₂ in the air-filled pore space. The diffusion coefficient (D_P) was calculated as a function of total porosity (Φ) and air-filled porosity (ε) and using the model proposed by *Moldrup et al.* [1999]:

$$\frac{D_P}{D_O} = \Phi^2 \left(\frac{\varepsilon}{\Phi} \right)^{2+\frac{1}{b}}, \quad (2)$$

where D_O is the gas diffusion coefficient in free air and b is the *Campbell* [1974] pore size distribution parameter. This parameter has been found to be strongly related ($r^2 = 0.96$) to clay fraction content (CF) through the following relationship [*Clapp and Hornberger*, 1978; *Olesen et al.*, 1996; *Rolston and Moldrup*, 2002]:

$$b = 13.6\text{CF} + 3.5. \quad (3)$$

[10] Diffusivity of CO₂ in the gas phase is about 4 orders of magnitude higher than in the liquid phase [*Šimunek and Suarez*, 1993; *Welsch and Hornberger*, 2004]; therefore, we assumed that solubility of the gas phase CO₂ is negligible. The characterization of the distribution of new moisture inputs in the soil, particularly in arid and semiarid environments, remains challenging because of wetting front instability or heterogeneity [*Wang et al.*, 2007]. Soil macropores caused by decaying roots, worm holes, and similar disturbances can cause preferential flow and differences in infiltration patterns [*Geiger and Durnford*, 2000; *Devitt and Smith*, 2002] under ponding [*Hill and Parlange*, 1972; *Glass et al.*, 1989a, 1989b; *Baker and Hillel*, 1990] and nonponding conditions [*Selker et al.*, 1992; *Babel et al.*, 1995; *Hendrickx and Yao*, 1996; *Yao and Hendrickx*, 1996]. Thus, we used the integrated 0–20 cm soil water content based on three replicates as a measure of volumetric water content over the top 20 cm of soil. Previous studies have studied the heterogeneity of new moisture distribution by applying vertical integrations [*Noborio et al.*, 1996; *Timlin and Pachepsky*, 2002] and have found that this estimate is a good representation of soil water content, even in extremely nonuniform conditions [*Topp et al.*, 1982a, 1982b; *Robinson et al.*, 2003]. Similarly, we assumed constant D_P/D_O and temperature (used for density estimation in D_O calculations) over the top 20 cm of the soil. We acknowledge potential inaccuracies introduced by this approach and current probe design constraints for water content profile estimation; however, this is an effective approximation of volumetric soil water content and the D_P/D_O parameter.




Figure 2. Variability of (a) precipitation and soil water content (θ) for (b) riparian and (c) upland sites for the 2006 growing season. Measurements of θ were made continuously at 20 cm and discretely integrating over 0–20 cm as indicated by the symbols.

[11] Because solid-state soil CO_2 probes are known to release heat after operating for long periods of time [Hirano *et al.*, 2003; Jassal *et al.*, 2005], we installed a double-pole double-throw relay (6 VDC coil voltage, 115 mA, Tyco Electronics, Berwyn, Pennsylvania) in the power line between the battery bank and the solid-state soil CO_2 probes, controlled by the data logger. This setup allowed us to switch the probes on prior to each measurement, including warming time as recommended by the manufacturer, and to switch them off to prevent long-term heating while saving >75% of battery power.

2.4. Ecosystem Respiration

[12] Continuous measurements of land-atmosphere CO_2 and water vapor exchange were made above the canopy of both ecosystems with the eddy covariance method [Baldocchi, 2003]. Wind velocity was measured with a triaxial sonic anemometer (CSAT3, Campbell Scientific Inc., Logan, Utah). Carbon dioxide and water vapor fluctuations were measured with an open path, infrared absorption gas analyzer (7500, LI-COR, Lincoln, Nebraska). Measurements were made at 10-Hz frequencies for the duration of the study. Estimates of nighttime ecosystem respiration were selected on the basis of fluxes between 2300 LT and 0400 LT and reported on a daily basis. A U^* threshold of 0.2 m s^{-1} was used to ensure periods with enough turbulence and reliable eddy covariance estimates. Because the purpose of the eddy covariance measurements was exclusively to provide a relative comparison, values are presented as nighttime ecosystem respiration fluxes, and no daytime

correction was applied to fluxes to avoid postprocessing and modeling abstraction.

3. Results

3.1. Environmental Variables

[13] The variability of continuous and discrete measurements of volumetric soil water content (θ) is presented in Figure 2. Throughout the growing season, values of θ decreased from over 50 to $\sim 10\%$ at the riparian site and from ~ 18 to $\sim 12\%$ at the upland site (Figure 2). Given that 70% of precipitation comes as snow and that peak runoff usually occurs between mid-May and early June [Woods *et al.*, 2006], these ecosystems are subject to a rapid spring wet up followed by a prolonged seasonal dry down, typical of subalpine forests. Rainfall was higher in magnitude and occurred more frequently before mid-July, after which rainfall decreased and occurred on only 2 days late in the season (Figure 2). The effects of rainfall on θ differed in magnitude between the wet riparian site and the drier upland site (Figure 2) and depended on antecedent conditions (wet soil versus dry soil). These effects were also reflected, both mechanistically and methodologically, on measured soil CO_2 fluxes (section 3.2).

[14] Soil temperature (T_s) varied both daily and seasonally (Figure 3), with a seasonal maximum toward the end of July at both sites. This time corresponds with the minimum or near-minimum values of θ (Figure 2), maximum soil thermal diffusivity [Ochsner *et al.*, 2001], maximum soil gas diffusivity [Riveros-Iregui *et al.*, 2007], and the initial decrease in potential for biological activity, sporadically




Figure 3. Variability in (a) precipitation, (b, d) soil temperature, and (c, e) soil CO₂ efflux at the riparian and upland sites for the 2006 growing season. CO₂ efflux was measured by the soil CO₂ profile technique (solid-state CO₂ sensors installed at depth) and by the soil respiration chamber technique. Boxes represent the mean, and error bars represent the standard deviation of three chamber measurements.

reset by precipitation events (e.g., 18 August). While the timing of daily T_s maxima was well synchronized between sites, the amplitude of the diel variability of T_s was lower at the upland site (Figure 3) because of the energy buffer imposed by the canopy cover. Overnight freezing temperatures in August drove a decrease in daily T_s maxima at both sites, indicating the decline of the growing season.

3.2. Soil CO₂ Efflux

[15] Both the soil respiration chamber and the soil CO₂ technique measured increasing fluxes early in the growing season at the riparian and upland sites (Figure 3), with the soil CO₂ profile having the advantage of increased sampling frequency, allowing for detailed visualization of the diel dynamics of soil CO₂ efflux. Similar diel dynamics have

been previously studied in detail [Tang *et al.*, 2005a; Riveros-Iregui *et al.*, 2007]; however, high-frequency measurements of soil CO₂ efflux responding to seasonal changes in environmental conditions (e.g., snowmelt and early and late season rainfall) between two sites within the same small-watershed ecosystem is, to our knowledge, unprecedented. Confidence bounds for 350 and 550 ppm at the soil-atmosphere interface ($z = 0$) are shown along with soil CO₂ efflux from the profile technique at both sites. The dotted lines in Figures 3c and 3e, almost identical to the solid lines, demonstrate that the error introduced by our approach is minimal compared to the natural variability of soil CO₂ efflux on a diel and seasonal basis. For the remainder of this paper we will refer to calculations and analyses based on




Figure 4. Comparison of instantaneous soil CO₂ efflux measured by both techniques during the 2006 growing season. Data include measurements at the riparian and the upland sites during rainy and dry days.

soil CO₂ concentration of 450 ppm at the soil-atmosphere interface.

[16] Seasonal dynamics recorded by the chamber technique agreed well with estimates by the soil CO₂ profile technique at both sites (Figure 3). However, marked differences throughout the study were imposed by high soil water content and sporadic rainfall. Comparing chamber measurements and instantaneous efflux from the profile technique for the entire season shows moderate agreement for the entire growing season at both sites ($r^2 = 0.51$; Figure 4). This agreement improved considerably when measurements within 2 days of at least 1-mm rainfall were excluded and when chamber measurements made on wet soil ($\theta > 0.25 \text{ m}^3 \text{ m}^{-3}$) were further excluded from the comparison at each site (Figure 5). However, by excluding such environmental disturbances from technique comparison, little can be learned about ways to improve technique performance or overcome their limitations; therefore, disagreement due to disturbances must not be omitted from these types of studies.

[17] Chamber measurements fell within the range of diel values of soil CO₂ efflux, except early in the season at the riparian site or following precipitation events at both sites (e.g., 10 and 13 July, Figure 6a). Accumulated on a diel basis (Figure 6b), rates of soil CO₂ efflux indicated that both techniques provide comparable estimates with the exception of those days following precipitation events, when chamber measurements were up to 84% higher than estimates by the soil CO₂ profile technique.

[18] Because the soil CO₂ profile technique is currently applied at different sampling frequencies (e.g., 5 min [Vargas and Allen, 2008] and 20 min (this study)), we tested the susceptibility of this technique to sampling frequency and time-of-day biases. Our results demonstrate that high sampling frequencies do not necessarily improve seasonal estimates of soil CO₂ efflux rates (Figure 7). Using the time of day at which the soil respiration chamber was deployed at each site (between 1000 LT and 1600 LT) to

capture instantaneous efflux by the soil CO₂ profile technique, similar seasonal estimates were found using all profile measurements (5973 data points) or interpolating between 34 daytime measurements at the riparian site and 10 daytime measurements at the upland site (Figure 7).

4. Discussion

4.1. What Are the Mechanisms Driving the Main Differences in Soil CO₂ Efflux From Riparian Meadows and Upland Forests?

[19] A major challenge in process understanding of soil CO₂ generation and efflux lies in the spatiotemporal nature of its biophysical controls. The interaction among soil temperature, vegetation, soil substrate, soil physical properties, and the landscape-induced redistribution of soil water can exhibit confounding effects on soil CO₂ efflux processes [Davidson *et al.*, 1998]. However, particularly in subalpine ecosystems, an important element that can be used to our advantage is the redistribution and seasonality in soil water content. There exists a degree of predictability in that snowmelt controls the time of the most dramatic increase in soil water content. Furthermore, landscape morphology redistributes that moisture downslope to lower areas of the landscape. Only through sporadic convective summer storms does the ecosystem receive new moisture inputs that can enhance biological activity. In our study, a seasonal comparison based on landscape position (Figure 3) demonstrates that soil CO₂ efflux at the riparian site was higher for 72% of the growing season, particularly after snowmelt, and when rainfall drove θ higher at the riparian site. Only when similar θ values were found between sites (e.g., after 29 July)




Figure 5. Comparison of techniques between (a, c) riparian and (b, d) upland sites, with all measurements at each site (Figures 5a and 5b), and removing measurements on soil with $\theta > 0.25 \text{ (m}^3 \text{ m}^{-3}\text{)}$ and measurements within 2 days of precipitation over 1 mm (Figures 5c and 5d).




Figure 6. (a) Comparison of soil CO_2 efflux measurements and soil temperature (T_s) made on 3 different days at the (left) riparian and (right) upland sites. Symbols denote day of the year. Open symbols denote measurements by the soil profile technique, and filled symbols denote measurements by the soil respiration chamber. After rain events (i.e., 10–13 July), measurements by the soil respiration chamber are much greater than estimates by the soil profile technique. (b) Comparison of cumulative fluxes over 24-h intervals at different times of the year at the (top) riparian and (bottom) upland sites. Dates of measurements are indicated in each plot and correspond to dates of measurements in Figure 6a. A total of 10.7 mm of rain occurred over 4 days prior to 10–13 July measurements.

were soil CO_2 efflux rates similar or higher at the upland site (Figure 3).

[20] A comparison of soil CO_2 efflux and soil temperature during early, middle, and late season (Figure 6a) between the riparian (wet) and upland (dry) sites demonstrated spatial and temporal differences in diel hysteresis patterns introduced by differences in diel T_s and CO_2 efflux. Recent studies have highlighted the evolution of diel patterns in both soil CO_2 concentrations [Riveros-Iregui *et al.*, 2007] and soil CO_2 effluxes [Carbone *et al.*, 2008]. Diel hysteresis in soil CO_2 concentrations is controlled by water content-limited soil CO_2 diffusion [Riveros-Iregui *et al.*, 2007]. A decline in θ results in enhanced soil CO_2 diffusion, allowing belowground concentrations to remain at steady state ($\partial CO_2/\partial t = 0$ [Riveros-Iregui *et al.*, 2007]). However, while the belowground concentrations can remain at steady state, the observed aboveground efflux (F) is not at steady

state ($\partial F/\partial t \neq 0$), indicating that, particularly in dry soils, diel hysteresis in soil CO_2 efflux is production limited and represents a rapid response of combined heterotrophic and autotrophic activities. Greater hysteresis patterns in efflux at the riparian site (e.g., 23 June at the riparian site versus 22 June at the upland site) are likely due to the effects of faster short vegetation response to photosynthetic activity at this site [Carbone and Trumbore, 2007] and the effects of a taller and more complete canopy cover on T_s at the upland site. More circular hysteresis patterns late in the season at the upland site (Figure 6a) indicate enhanced photosynthetic activity of the forest canopy with respect to riparian grasses [Emanuel, 2007], and that riparian vegetation (senescing by this time of the year) underwent late season water stress before upland vegetation. This suggests that upland vegetation is better adapted to lower θ , whereas riparian vegetation is adapted to higher θ and is sensitive to θ reduction over the




Figure 7. Comparison of cumulative soil CO₂ efflux estimates by the soil CO₂ profile technique, modifying its sampling frequency. Solid lines indicate estimates from 20-min sampling over 83 days ($n = 5973$). Dashed ($n = 34$) and dotted ($n = 10$) lines indicate reduced sampling frequency (every 2–7 days), using the time of the soil respiration chamber deployment as an example. Agreement between high and moderate frequencies suggests little time-of-day and sampling frequency bias. This comparison demonstrates that on a cumulative basis the use of the soil CO₂ profile technique at 20-min intervals yields similar results as when it is used every 2–7 days.

growing season. As a result, the riparian vegetation influence on soil CO₂ efflux diminished over the course of seasonal dry down.

[21] These findings demonstrate how soil water content distribution across the landscape exerts a major control on both spatial and temporal (seasonal) differences of soil CO₂ efflux. We suggest that parameterization of water content heterogeneity in space and time must be a critical component for realistic model representations of soil CO₂ efflux (understood as the balance between production and transport [*Pacific et al.*, 2008]) from heterogeneous landscapes. To date, this fundamental concept remains to be robustly applied and integrated within studies of land-atmosphere exchange at the ecosystem level.

4.2. How Do Two of the Most Commonly Used Methods to Measure Soil CO₂ Efflux Compare Across Sites and Across the Growing Season?

[22] To assess the effects of high soil water content and rainfall on discrepancies between techniques, we compared each site separately, both including and excluding measurements following rainfall and during early season high θ (Figure 5). At both sites technique agreement significantly improved when measurements on wet soil days ($\theta > 0.25 \text{ m}^3 \text{ m}^{-3}$) and measurements taken within 2 days of ≥ 1 -mm rainfall were removed from the comparison. The disagreement between techniques following precipitation is to be expected as new water inputs can cause a CO₂ burst in soil air because of a rapid gas displace-

ment in the pore space followed by enhanced biological activity [*Cable and Huxman*, 2004; *Huxman et al.*, 2004]. However, the disagreement caused by the removal of measurements with high θ indicates that parameterization of the soil CO₂ profile method needs to be strongly improved and most likely differentiated between high θ and low θ , especially in ecosystems with large variability of θ . Previous studies attempting this technique corroboration [*Baldocchi et al.*, 2006; *Vargas and Allen*, 2008] do not provide context for when chamber measurements were taken with respect to the seasonality of θ , hence, little can be learned from the agreement (or disagreement) of their techniques. Our results suggest that while good technique comparison can be attained during periods of stable conditions (e.g., constant θ and no rainfall), environmental disturbances will affect method corroboration in space and time. Excluding chamber measurements taken following rainfall would improve technique correlation, but through this or similar exclusions, information on primary controls on soil CO₂ efflux is lost. The strengths and limitations of each method, as well as full system understanding, can only be achieved with the direct comparison of both approaches. A context for environmental conditions under which measurements were taken is necessary to understand technique performance (strengths and limitations) and variability of the fluxes (i.e., distinction between ecophysiological processes and environmental biases).

[23] Analyzing instantaneous fluxes from the soil CO₂ profile technique at different sampling frequencies indicates that on a cumulative basis the soil profile technique is not biased by the time of the day at which sampling occurs or by the sampling frequency itself (Figure 7). This means that on a cumulative basis the use of the soil CO₂ profile technique at 20-min frequency intervals yields similar results as when it is used every 2–7 days. Given that daily minima of soil CO₂ in this system occur before sunrise and daily maxima occur during early to late evening [*Riveros-Iregui et al.*, 2007], sampling soil CO₂ efflux between 1000 LT and 1600 LT (as in the example presented in Figure 7) may correspond to the 24-h mean of soil CO₂ efflux. Our findings suggest that on a seasonal basis, it is more critical to capture spatial variability and seasonal dynamics driven primarily by changes in soil water content than the diel dynamics caused by soil temperature and plant activity. These results demonstrate that while the soil CO₂ probes provide important resolution for short time scales, long-term (seasonal) measurements do not necessarily benefit from this high-frequency sampling. The tradeoff between spatial coverage of chambers and temporal resolution of the soil CO₂ profile technique greatly depends on study goals and whether one is interested in seasonal estimates of soil CO₂ efflux rates or rapid dynamics of this flux.

4.3. What Are the Implications of These Findings for Process Understanding of Soil CO₂ Efflux From Subalpine Ecosystems?

[24] A comparison of measurements by the soil respiration chamber and the soil CO₂ profile technique demonstrates that, accumulated over the growing season, both techniques are within 7% of measurements for the riparian site and within 32% of measurements for the upland site (Figure 8). Similar agreements between techniques have




Figure 8. (a) Relative precipitation. (b, c, d) Comparison of cumulative discrete (instantaneous) measurements made by the soil respiration chamber technique (including all data), 20-min measurements by the soil profile technique, and nighttime ecosystem respiration (R_E) by an eddy covariance system at a riparian and an upland site.

been reported [Tang *et al.*, 2003; Baldocchi *et al.*, 2006], but the difference in agreement across sites had not been previously observed. Better agreement at the riparian site is likely due to the higher θ in these areas, which leads to a more homogeneous water content profile in the top 20 cm of the soil, even after precipitation events (Figure 2). New moisture from precipitation can be distributed more homogeneously within the top section of the soil profile. Conversely, because θ is lower at the upland site, the distribution of new moisture from precipitation in the topsoil does not occur as homogeneously, and new moisture does not penetrate as deeply, causing larger differences in diffusivity throughout the soil profile, thus limiting the soil profile technique, especially late in the season.

[25] Compared to nighttime ecosystem respiration, both techniques provide comparable effluxes for each site (Figure 8). While this comparison is not intended for quantitative purposes, it provides the foundations for potential, detailed examinations. For example, the difference in soil CO₂ efflux response between the riparian and upland sites, particularly after snowmelt and precipitation, demonstrates that different landscape positions may not respond uniformly to environmental disturbances, particularly because of differences in θ . On a cumulative basis riparian areas exhibit higher soil CO₂ efflux and ecosystem respiration than upland sites throughout the growing season (Figure 8d). While riparian meadows can occupy a smaller fraction of an entire forest ($\sim 2\%$) [Riveros-Iregui, 2008], soil CO₂ efflux from these areas is larger than effluxes from upland forests. Soil CO₂ efflux from chamber measurements was within 16% of nighttime ecosystem respiration at the riparian site and within 15% of nighttime ecosystem respiration at the upland site. However, these relationships, and

the magnitude of differences between sites and throughout the growing season, are nonlinear, which warrants future investigations on how the parameterization of a nonstationary behavior of the landscape can be important to improve current estimates of soil CO₂ efflux from large areas and to improve comparisons with other estimates of C exchange at the ecosystem scale. Direct comparison of multiple techniques (soil respiration chambers, soil CO₂ profile technique, and eddy covariance towers) is necessary to understand the spatiotemporal nature of C fluxes. The findings presented here are essential for enhancing process understanding of soil CO₂ efflux from heterogeneous landscapes, providing a conceptual framework of soil CO₂ efflux useful for modeling studies, and gaining confidence in current and developing soil CO₂ efflux measuring techniques.

5. Conclusions and Implications

[26] 1. Soil water content was a major control on both spatial and temporal (particularly seasonal) differences of soil CO₂ efflux between a riparian meadow and an upland forest, especially after snowmelt and rainfall. Parameterization of water content heterogeneity in space and time must be a critical component of realistic model representations of soil CO₂ efflux rates from heterogeneous landscapes.

[27] 2. Good agreement between the soil respiration chamber technique and the soil CO₂ profile technique can be attained during periods of stable conditions (e.g., constant θ and no rainfall). However, seasonality of soil water content and sporadic rainfall introduce physical effects that limit this agreement and play a major role in method corroboration. Providing a context for environmental conditions under which measurements were taken is necessary

to understanding performance of techniques and the source of the variability in measured efflux.

[28] 3. On a 24-h basis both techniques yield comparable results, except during periods of sporadic precipitation, when the chamber technique yields soil CO₂ efflux rates much larger than those by the soil profile technique. This means that rapid changes in soil physical properties, respiration enhancement, and water-caused displacement of CO₂ within the soil pore space might not be adequately captured by solid-state CO₂ sensors. Nonetheless, these sensors remain a useful tool for capturing changes in soil CO₂ caused by less transient, nonhydrological, ecophysiological processes (i.e., responses of plant and microbial activity to changing environmental conditions).

[29] 4. Differences in diel hysteresis patterns of soil CO₂ efflux between sites were controlled by the effects of canopy cover on soil temperature and temporal differences in photosynthetic activity of vegetation. Particularly in dry soils, diel hysteresis in soil CO₂ efflux is production limited and represents a rapid response of combined heterotrophic and autotrophic activities. More circular hysteresis patterns late in the season at the upland site suggest that upland vegetation is better adapted to low soil water content and that riparian vegetation influence on soil CO₂ efflux diminished over the course of seasonal dry down.

[30] 5. On a seasonal basis soil CO₂ efflux measurements were not biased by the time of the day at which sampling occurred, meaning that long-term (seasonal) measurements do not necessarily benefit from high-frequency sampling by soil CO₂ probes. Furthermore, if the study focuses on seasonality, capturing the spatial variability and seasonal dynamics of efflux driven primarily by changes in soil water content is more important than capturing diel dynamics caused by soil temperature and plant activity.

[31] 6. Both techniques can yield comparable efflux rates with the exception of transient flux events caused by precipitation or diffusion-limited flux caused by wet soils. Direct comparison of multiple techniques (soil respiration chambers, soil CO₂ profile technique, and eddy covariance towers) is necessary to gain insight into the primary controls on soil CO₂ production and transport and the spatiotemporal nature of efflux. Acknowledging technique limitations is important for reporting realistic rates of soil CO₂ efflux.

[32] 7. These results have implications for interpreting and evaluating rates of soil CO₂ efflux measured by soil respiration chambers and the soil CO₂ profile technique, from both mechanistic and methodological perspectives. While the findings presented here were attained in a subalpine forest, they can be applied to current and future studies in a wide range of ecosystems. These implications should be considered when measuring and modeling the dynamics of C cycling at the ecosystem level.

[33] **Acknowledgments.** Financial support was provided by the U.S. National Science Foundation Integrated Carbon Cycle Research Program (grants EAR-0404130, EAR-0403924, and EAR-0403906). D.A.R.-I. acknowledges support from the 2007 American Geophysical Union Horton Research Grant, a NSF Doctoral Dissertation Improvement grant (DEB-0807272), and the USGS 104b Grant Program, administered by the Montana Water Center. The authors thank the Tenderfoot Creek Experimental Forest and the USDA, Forest Service, Rocky Mountain Research Station, particularly Ward McCaughey, V. J. Pacific, R. E. Emanuel, and D. J. Muth provided helpful discussions during the devel-

opment of this work. Two anonymous reviewers provided helpful comments on a previous version of this paper.

References

- Babel, M. S., R. Loof, and A. Dasgupta (1995), Fingering preferential flow in unsaturated homogeneous coarse sands, *Hydrol. Sci. J.*, 40(1), 1–17.
- Baker, R. S., and D. Hillel (1990), Laboratory tests of a theory of fingering during infiltration into layered soils, *Soil Sci. Soc. Am. J.*, 54(1), 20–30.
- Baldocchi, D. D. (2003), Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of ecosystems: Past, present and future, *Global Change Biol.*, 9(4), 479–492, doi:10.1046/j.1365-2486.2003.00629.x.
- Baldocchi, D., J. Tang, and L. Xu (2006), How switches and lags in biophysical regulators affect spatial-temporal variation of soil respiration in an oak-grass savanna, *J. Geophys. Res.*, 111, G02008, doi:10.1029/2005JG000063.
- Burrows, E. H., J. L. Bubier, A. Mosedale, G. W. Cobb, and P. M. Crill (2005), Net ecosystem exchange of carbon dioxide in a temperate poor fen: A comparison of automated and manual chamber techniques, *Biogeochemistry*, 76(1), 21–45, doi:10.1007/s10533-004-6334-6.
- Cable, J. M., and T. E. Huxman (2004), Precipitation pulse size effects on Sonoran Desert soil microbial crusts, *Oecologia*, 141(2), 317–324, doi:10.1007/s00442-003-1461-7.
- Campbell, G. S. (1974), A simple method for determining unsaturated conductivity from moisture retention data, *Soil Sci.*, 117(6), 311–314, doi:10.1097/00010694-197406000-00001.
- Carbone, M. S., and S. E. Trumbore (2007), Contribution of new photosynthetic assimilates to respiration by perennial grasses and shrubs: Residence times and allocation patterns, *New Phytol.*, 176(1), 124–135, doi:10.1111/j.1469-8137.2007.02153.x.
- Carbone, M. S., G. C. Winston, and S. E. Trumbore (2008), Soil respiration in perennial grass and shrub ecosystems: Linking environmental controls with plant and microbial sources on seasonal and diel timescales, *J. Geophys. Res.*, 113, G02022, doi:10.1029/2007JG000611.
- Clapp, R. B., and G. M. Hornberger (1978), Empirical equations for some soil hydraulic properties, *Water Resour. Res.*, 14(4), 601–604, doi:10.1029/WR014i004p06001.
- Davidson, E. A., E. Belk, and R. D. Boone (1998), Soil water content and temperature as independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest, *Global Change Biol.*, 4(2), 217–227, doi:10.1046/j.1365-2486.1998.00128.x.
- Davidson, E. A., K. Savage, L. V. Verchot, and R. Navarro (2002), Minimizing artifacts and biases in chamber-based measurements of soil respiration, *Agric. For. Meteorol.*, 113(1–4), 21–37, doi:10.1016/S0168-1923(02)00100-4.
- Devitt, D. A., and S. D. Smith (2002), Root channel macropores enhance downward movement of water in a Mojave Desert ecosystem, *J. Arid Environ.*, 50(1), 99–108, doi:10.1006/jare.2001.0853.
- Emanuel, R. E. (2007), Vegetation water stress and its impact on land-atmosphere interaction, Ph.D. dissertation, 162 pp., Univ. of Va., Charlottesville.
- Fang, C., and J. B. Moncrieff (1996), An improved dynamic chamber technique for measuring CO₂ efflux from the surface of soil, *Funct. Ecol.*, 10(2), 297–305, doi:10.2307/2389856.
- Fames, P. E., R. C. Shearer, W. W. McCaughey, and K. J. Hansen (1995), Comparisons of hydrology, geology, and physical characteristics between Tenderfoot Creek Experimental Forest (east side) Montana, and Coram Experimental Forest (west side) Montana, *Final Rep. RJVA-INT-92734*, For. Sci. Lab., Intermountain Res. Stn., U.S. Dep. Agric. For. Serv., Bozeman, Mont.
- Geiger, S. L., and D. S. Durnford (2000), Infiltration in homogeneous sands and a mechanistic model of unstable flow, *Soil Sci. Soc. Am. J.*, 64(2), 460–469.
- Glass, R. J., J.-Y. Parlange, and T. S. Steenhuis (1989a), Wetting front instability: 1. Theoretical discussion and dimensional analysis, *Water Resour. Res.*, 25(6), 1187–1194, doi:10.1029/WR025i006p01187.
- Glass, R. J., T. S. Steenhuis, and J.-Y. Parlange (1989b), Wetting front instability: 2. Experimental determination of relationships between system parameters and two-dimensional unstable flow field behavior in initially dry porous media, *Water Resour. Res.*, 25(6), 1195–1207, doi:10.1029/WR025i006p01195.
- Goulden, M. L., and P. M. Crill (1997), Automated measurements of CO₂ exchange at the moss surface of a black spruce forest, *Tree Physiol.*, 17(8–9), 537–542.
- Hendrickx, J. M. H., and T. M. Yao (1996), Prediction of wetting front stability in dry field soils using soil and precipitation data, *Geoderma*, 70(2–4), 265–280, doi:10.1016/0016-7061(95)00082-8.
- Hill, D. E., and J.-Y. Parlange (1972), Wetting front instability in layered soils, *Soil Sci. Soc. Am. Proc.*, 36(5), 697–702.

- Hirano, T., H. Kim, and Y. Tanaka (2003), Long-term half-hourly measurement of soil CO₂ concentration and soil respiration in a temperate deciduous forest, *J. Geophys. Res.*, 108(D20), 4631, doi:10.1029/2003JD003766.
- Hutchinson, G. L., and G. P. Livingston (2001), Vents and seals in non-steady-state chambers used for measuring gas exchange between soil and the atmosphere, *Eur. J. Soil Sci.*, 52(4), 675–682, doi:10.1046/j.1365-2389.2001.00415.x.
- Huxman, T. E., J. M. Cable, D. D. Ignace, J. A. Eilts, N. B. English, J. Weltzin, and D. G. Williams (2004), Response of net ecosystem gas exchange to a simulated precipitation pulse in a semi-arid grassland: The role of native versus non-native grasses and soil texture, *Oecologia*, 141(2), 295–305.
- Jassal, R., A. Black, M. Novak, K. Morgenstern, Z. Nestic, and D. Gaumont-Guay (2005), Relationship between soil CO₂ concentrations and forest-floor CO₂ effluxes, *Agric. For. Meteorol.*, 130(3–4), 176–192, doi:10.1016/j.agrformet.2005.03.005.
- Lloyd, J., and J. A. Taylor (1994), On the temperature-dependence of soil respiration, *Funct. Ecol.*, 8(3), 315–323, doi:10.2307/2389824.
- Moldrup, P., T. Olesen, T. Yamaguchi, P. Schjonning, and D. E. Rolston (1999), Modeling diffusion and reaction in soils: IX. The Buckingham-Burdine-Campbell equation for gas diffusivity in undisturbed soil, *Soil Sci.*, 164(8), 542–551, doi:10.1097/00010694-199908000-00002.
- Noborio, K., K. J. McInnes, and J. L. Heilman (1996), Measurements of soil water content, heat capacity, and thermal conductivity with a single TDR probe, *Soil Sci.*, 161(1), 22–28, doi:10.1097/00010694-199601000-00004.
- Norman, J. M., C. J. Kucharik, S. T. Gower, D. D. Baldocchi, P. M. Crill, M. Rayment, K. Savage, and R. G. Striegl (1997), A comparison of six methods for measuring soil-surface carbon dioxide fluxes, *J. Geophys. Res.*, 102(D24), 28,771–28,777, doi:10.1029/97JD01440.
- Ochsner, T. E., R. Horton, and T. H. Ren (2001), A new perspective on soil thermal properties, *Soil Sci. Soc. Am. J.*, 65(6), 1641–1647.
- Olesen, T., P. Moldrup, K. Henriksen, and L. W. Petersen (1996), Modeling diffusion and reaction in soils. 4. New models for predicting ion diffusivity, *Soil Sci.*, 161(10), 633–645, doi:10.1097/00010694-199610000-00001.
- Pacific, V. J., B. L. McGlynn, D. A. Riveros-Iregui, D. L. Welsch, and H. E. Epstein (2008), Variability in soil respiration across riparian-hillslope transitions, *Biogeochemistry*, doi:10.1007/s10533-008-9258-8.
- Porporato, A., P. D'Odorico, F. Laio, L. Ridolfi, and I. Rodriguez-Iturbe (2002), Ecohydrology of water-controlled ecosystems, *Adv. Water Resour.*, 25(8–12), 1335–1348.
- Ridolfi, L., P. D'Odorico, A. Porporato, and I. Rodriguez-Iturbe (2000), Impact of climate variability on the vegetation water stress, *J. Geophys. Res.*, 105(D14), 18,013–18,025, doi:10.1029/2000JD900206.
- Riveros-Iregui, D. A. (2008), Hydrologic-carbon cycle linkages in a subalpine catchment, Ph.D. dissertation, 220 pp., Mont. State Univ., Bozeman.
- Riveros-Iregui, D. A., R. E. Emanuel, D. J. Muth, B. L. McGlynn, H. E. Epstein, D. L. Welsch, V. J. Pacific, and J. M. Wraith (2007), Diurnal hysteresis between soil CO₂ and soil temperature is controlled by soil water content, *Geophys. Res. Lett.*, 34, L17404, doi:10.1029/2007GL030938.
- Robinson, D. A., S. B. Jones, J. M. Wraith, D. Or, and S. P. Friedman (2003), A review of advances in dielectric and electrical conductivity measurement in soils using time domain reflectometry, *Vadose Zone J.*, 2, 444–475.
- Rolston, D. E., and P. Moldrup (2002), Gas diffusivity, in *Methods of Soil Analysis: Part 4—Physical Methods*, edited by J. H. Dane and G. C. Topp, pp. 1113–1139, Soil Sci. Soc. of Am., Madison, Wis.
- Savage, K. E., and E. A. Davidson (2001), Interannual variation of soil respiration in two New England forests, *Global Biogeochem. Cycles*, 15(2), 337–350, doi:10.1029/1999GB001248.
- Scott-Denton, L. E., K. L. Sparks, and R. K. Monson (2003), Spatial and temporal controls of soil respiration rate in a high-elevation, subalpine forest, *Soil Biol. Biochem.*, 35(4), 525–534, doi:10.1016/S0038-0717(03)00007-5.
- Selker, J., P. Leclercq, J.-Y. Parlange, and T. Steenhuis (1992), Fingering flow in two dimensions: I. Measurement of matric potential, *Water Resour. Res.*, 28(9), 2513–2521, doi:10.1029/92WR00963.
- Šimunek, J., and D. L. Suarez (1993), Modeling of carbon dioxide transport and production in soil: I. Model development, *Water Resour. Res.*, 29(2), 487–497, doi:10.1029/92WR02225.
- Subke, J. A., and J. D. Tenhunen (2004), Direct measurements of CO₂ flux below a spruce forest canopy, *Agric. For. Meteorol.*, 126(1–2), 157–168, doi:10.1016/j.agrformet.2004.06.007.
- Tang, J. W., and D. D. Baldocchi (2005), Spatial-temporal variation in soil respiration in an oak-grass savanna ecosystem in California and its partitioning into autotrophic and heterotrophic components, *Biogeochemistry*, 73(1), 183–207, doi:10.1007/s10533-004-5889-6.
- Tang, J. W., D. D. Baldocchi, Y. Qi, and L. K. Xu (2003), Assessing soil CO₂ efflux using continuous measurements of CO₂ profiles in soils with small solid-state sensors, *Agric. For. Meteorol.*, 118(3–4), 207–220, doi:10.1016/S0168-1923(03)00112-6.
- Tang, J. W., D. D. Baldocchi, and L. Xu (2005a), Tree photosynthesis modulates soil respiration on a diurnal time scale, *Global Change Biol.*, 11(8), 1298–1304, doi:10.1111/j.1365-2486.2005.00978.x.
- Tang, J. W., L. Misson, A. Gershenson, W. X. Cheng, and A. H. Goldstein (2005b), Continuous measurements of soil respiration with and without roots in a ponderosa pine plantation in the Sierra Nevada Mountains, *Agric. For. Meteorol.*, 132(3–4), 212–227, doi:10.1016/j.agrformet.2005.07.011.
- Timlin, D., and Y. Pachepsky (2002), Infiltration measurement using a vertical time-domain reflectometry probe and a reflection simulation model, *Soil Sci.*, 167(1), 1–8, doi:10.1097/00010694-200201000-00001.
- Topp, G. C., J. L. Davis, and A. P. Annan (1982a), Electromagnetic determination of soil water content using TDR: I. Applications to wetting fronts and steep gradients, *Soil Sci. Soc. Am. J.*, 46(4), 672–678.
- Topp, G. C., J. L. Davis, and A. P. Annan (1982b), Electromagnetic determination of soil water content using TDR: II. Evaluation of installation and configuration of parallel transmission lines, *Soil Sci. Soc. Am. J.*, 46(4), 678–684.
- Vargas, R., and M. F. Allen (2008), Diel patterns of soil respiration in a tropical forest after Hurricane Wilma, *J. Geophys. Res.*, 113, G03021, doi:10.1029/2007JG000620.
- Wang, X.-P., X.-R. Li, H.-L. Xiao, R. Berndtsson, and Y.-X. Pan (2007), Effects of surface characteristics on infiltration patterns in an arid shrub desert, *Hydrol. Process.*, 21(1), 72–79, doi:10.1002/hyp.6185.
- Welsch, D. L., and G. M. Hornberger (2004), Spatial and temporal simulation of soil CO₂ concentrations in a small forested catchment in Virginia, *Biogeochemistry*, 71(3), 413–434, doi:10.1023/B:BI0G.0000049350.24911.e9.
- Woods, S. W., R. Ahl, J. Sappington, and W. McCaughey (2006), Snow accumulation in thinned lodgepole pine stands, Montana, USA, *For. Ecol. Manage.*, 235(1–3), 202–211, doi:10.1016/j.foreco.2006.08.013.
- Yao, T. M., and J. M. H. Hendrickx (1996), Stability of wetting fronts in dry homogeneous soils under low infiltration rates, *Soil Sci. Soc. Am. J.*, 60(1), 20–28.

H. E. Epstein, Department of Environmental Sciences, University of Virginia, Charlottesville, VA 22904, USA.

B. L. McGlynn and D. A. Riveros-Iregui, Department of Land Resources and Environmental Sciences, Montana State University, 334 Leon Johnson Hall, Bozeman, MT 59717, USA. (diego.riverosiregui@myportal.montana.edu)

D. L. Welsch, Canaan Valley Institute, P.O. Box 673, Davis, WV 26260, USA.