

2006

The Case Against Evening-Up Litters Until Weaning

Duane Reese

University of Nebraska-Lincoln, dreese1@unl.edu

Barbara Straw

University of Nebraska-Lincoln

Follow this and additional works at: http://digitalcommons.unl.edu/coopext_swine



Part of the [Animal Sciences Commons](#)

Reese, Duane and Straw, Barbara, "The Case Against Evening-Up Litters Until Weaning" (2006). *Nebraska Swine Reports*. 212.
http://digitalcommons.unl.edu/coopext_swine/212

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Swine Reports by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



The Case Against Evening-Up Litters Until Weaning

Duane E. Reese
Barbara E. Straw¹

Summary and Implications

A literature review on the effect of fostering or moving individual piglets from one litter to another after they are 24 to 48 hours of age was conducted. Late fostering disrupts nursing, increases fighting, and impairs the growth rate of adopted piglets and their littermates. Pig body weight at weaning was reduced 13 to 24% in extensively fostered litters vs. those where no piglets were fostered after 48 hours of age. No evidence was found that late fostering improves preweaning survival. For the greater good of all piglets, producers are encouraged to resist the urge to even-up litters or foster individual piglets after they are 24 hours old. Piglets that fall behind or grow slower than littermates after the initial fostering is done should be transferred to nurse sows where an entirely new litter(s) of older pigs is made. Milk replacers can also play a role in providing slower-growing or starving piglets more milk.

Introduction

Fostering or moving piglets from one litter to another is commonly practiced in swine operations to adjust litter size so that all piglets have good access to sow milk during lactation. Proper fostering reduces preweaning mortality and probably the number of substandard pigs at weaning. Many farrowing managers and employees know all fostering should be completed before the piglets are 24 to 48 hours old for best results. However, in some operations, moving individual piglets between litters or “evening-up” continues until weaning. Some

farrowing house personnel hate to see one litter with 10 pigs and the one next to it with seven. Also, some believe that a piglet in one litter that is falling behind littermates would be better off living in another litter of more similar-sized piglets, especially if there are fewer piglets in the recipient litter. Basically the goal is to have all litters in the farrowing area uniform or looking like “peas in a pod.” This paper will review the literature on fostering to clarify the issue for people who continue to even-up litters until weaning. Also, options for accommodating fall-outs or piglets that grow slower than littermates before weaning will be presented.

Research Summary

Michigan State University researchers conducted a study with 80 litters on a farm where extensive transfer of pigs between litters was being done on a daily basis until weaning. In 40 litters the usual practice of continuous fostering until weaning was continued. In another 40 litters fostering was limited to the first two days of life.

The effect of extensive fostering on pig body weight and standard deviation of body weight at weaning and preweaning mortality is presented in Table 1. As expected, extensive fostering

resulted in a lower average within-litter standard deviation of pig body weight at weaning (i.e., pigs were more uniform in size within litter); however, it also reduced pig weaning weight by 2.2 lb or 20%. Mortality was not significantly different between treatment groups, although it was numerically higher in the continuous fostering treatment. This research demonstrated that continuous fostering results in more uniform litters at weaning, but at the expense of growth rate and possibility survival.

In half of the 32 litters Canadian researchers studied, two piglets were exchanged between pairs of litters at 6 ± 1 day of age. Thus, there were three types of piglets in the study; adopted (piglets that were exchanged), resident (piglets that were not exchanged but were littermates to the adopted piglets), and control (no fostering). Piglets were weighed at birth, fostering, weaning (day 18 ± 1) and weekly during the next month.

The effect of fostering on pig body weight during the course of the trial is shown in Figure 1. There was no significant difference in body weight between piglet types at birth or just before fostering. However, at every time period after fostering, the average body weight of piglets in the fostered litters (those containing adopted and resident piglets) was significantly

(Continued on next page)

Table 1. Effect of limited vs. continuous fostering on pig performance during lactation.^a

Item	Limited fostering ^b	Continuous fostering
Average within-litter standard deviation of body weight at weaning ^c	2.0	0.7
Weaning weight, lb ^c	11.6	9.4
Mortality, %	8.0	8.8

^aStraw et al., 1998.

^bFirst two days of life only.

^cP < 0.008.




Figure 1. Effect of fostering on day 6 \pm 1 of lactation on piglet body weight before and after weaning (day 18 \pm 1). ^aBody weight between fostered (adopted + resident) and control litters differed ($P < 0.05$); ^bBody weight between adopted and resident piglets differed ($P < 0.05$). Adapted from Giroux et al., 2000.

reduced. Within the fostered litters the body weight of adopted piglets was significantly reduced compared to that of the resident piglets at each period. In conclusion, fostering had a marked effect on the growth rate of adopted piglets such that they gained only 76% of the weight of those in stable litters. While supporting the results of the Michigan State study, this research further demonstrated that adopted piglets may continue to be smaller after weaning.

In another Canadian study the behavior and growth of 13 control and 14 fostered litters was compared. Once every three days (from day one to 16 of lactation), all piglets were weighed and three piglets were switched between two fostered litters. Thus, there were three types of piglets in the study: adopted (piglets that were exchanged), resident (piglets that were not exchanged but were littermates to the adopted piglets), and control (no fostering). Behavior was observed for two hours after weighing and (or) fostering and during one nursing period 24 hours later.

Fights were significantly more

frequent in the fostered vs. control litters during and between nursings at all fostering periods except on day one. Most of the fights occurred between resident and fostered piglets except on day 1. While nursing, piglets in fostered litters fought significantly more than those in control litters at 24 hours after fostering except on day one and 16. Failed nursings and snaps by the sow toward piglets were significantly more frequent in fostered vs. control litters. Moreover, sows rearing fostered litters spent 15 to 30% less time lying on their sides at day four, seven, 13 and 16 ($P < 0.05$). Adopted piglets weighed 13% less than controls at weaning ($P < 0.01$); resident piglets were significantly heavier than adopted piglets, but smaller ($P = 0.1$) than controls.

This study provided insight into why continuous or late fostering reduces piglet weight gain. The presence of alien piglets in the litter disrupts nursing and therefore milk intake, not only as a result of fighting between piglets, but the sow is less accommodating to the nutritional and comfort needs of her litter. The study also con-

firmed that fostering is appropriate through the first day of life.

Better Management Options

It's common for one or more piglets in a few litters to fall behind or grow slower than littermates during lactation. These piglets are commonly called fall-outs, fall-backs, or runts. Many fall-outs will flourish once they have the opportunity to receive more milk. Producers can use other sows (i.e., nurse sows) and/or milk replacers to provide fall-outs more milk.

Nurse sows

Nurse sows can be created and utilized two ways. The preferred method is to identify a well-milking sow(s) to raise fall-outs that are collected from other sows. The procedure involves finding a newly farrowed gilt, i.e., one that finished farrowing six to 12 hours earlier. It is better to use a gilt, because she has smaller teats that are easier for piglets to grasp. Being newly farrowed is an advantage, because she doesn't know that the pigs she is about to receive




Figure 2. Example of bump weaning.

aren't her pigs. Identify eight to 10 fall-out pigs that are five to seven days of age and move them to the freed-up gilt. This moves them to a younger age group, but they are likely just starving and not sick. To be sure fall-outs are just starving and not sick, check their littermates to see if they are healthy. Also, observe if the fall-outs are being crowded out at the udder or are nursing a lower-producing teat (usually located at the rear of the udder). You do not want to move sick piglets, because that spreads disease around. The difficulty with this method is having enough spare sow capacity to take care of the gilt's original litter. It is important to wait six to 12 hours after the gilt has finished farrowing before her piglets are fostered to other sows to ensure all her piglets

receive a good dose of colostrum. Remember to foster the gilt's piglets to other sows that are nursing similar-aged piglets.

The other procedure, commonly called "bump weaning," involves moving fall-outs to a later lactation, good milking sow until they reach the normal weaning age for the farm (Figure 2). For example, assume there are three good-milking sows: sow A has lactated for three days, sow B has lactated for eight days and sow C has lactated for 14 days (five to seven days before she will be weaned). Sow C's piglets are weaned and Sow B's piglets are moved to Sow C. Sow A's piglets are moved to Sow B. Sow A is given 2 to 3 day-old fall-outs collected from several litters. The main disadvantage with this procedure is that too

often one or two adopted piglets per litter get injured by vicious sows. Ultimately, however, the detriment to the bumped pigs is probably outweighed by the benefit to the fall-outs (which were likely to die without some milk). For these reasons, bump weaning should be used as a last resort instead of a routine procedure. Bump weaning ensures that no piglets will be weaned later than the age limit set for the farm and that entire rooms of sows and litters can be weaned at the same time. Note that pigs are always moved forward and not backward in the system. Also, the key to making bump weaning work is to identify candidate pigs early in lactation (2 to 3 days of age) rather than later.

Milk replacers

Milk replacers offer another way for fall-outs to obtain more milk. Milk can be provided free choice in plastic milk feeders or baby bottles. Or you can place the fall-out in a plastic bin containing a feeder while it drinks. This method ensures the fall-out is not competing for milk and you can be sure it drinks. Initially the fall-out must be trained to drink from a bottle, but after a few feedings it catches on and takes advantage of the additional milk without competition. Fall-outs can also consume milk from a pan or bowl; some will need to be trained, however. To train fall-outs to drink from a bowl, place their snout in the milk for a few seconds every hour until they appear to have learned to drink on their own. Use bowls or pans that attach to or are held down to the floor so they cannot be knocked over.

Some producers place a deck or pen containing a milk feeder in each farrowing room (one deck per 12 crates, for example) to manage fall-outs. The best milking sow that has lactated for about 10 days is

(Continued on next page)



identified. Her pigs are removed and placed in the deck and fed milk replacer. Eight to 10 fall-outs are collected from various litters in the room and placed on the newly weaned sow.

Conclusion

Fostering piglets after they are 24 hours old disrupts nursing, increases fighting, and significantly

impairs the growth rate of adopted piglets and their littermates. Also, no evidence was found that late fostering improves preweaning survival. Therefore, for the greater good of all piglets, resist the urge to even-up litters or foster individual piglets after they are 24 hours old. Piglets that fall behind or grow slower than littermates after the initial fostering is done should be transferred to nurse sows where

an entirely new litter(s) of older pigs is made. Milk replacers can also play a role in providing fall-outs more milk.

¹Duane E. Reese is an extension swine specialist in the Animal Science Department at the University of Nebraska–Lincoln. Barbara E. Straw is an extension veterinarian in the Department of Large Animal Clinical Sciences at Michigan State University. References are available from dreese1@unl.edu by request.

Patterns of Drinking Water Use in Pork Production Facilities

Michael C. Brumm¹

Summary and Implications

The amount of drinking water needed daily by the pig depends on numerous influences, including temperature, diet, stage of production and health. Within a 24-hour period under thermal-neutral conditions, grow-finish and gestating swine demonstrate a peak in water usage in late afternoon while lactating females consume water more consistently throughout the day. In times of heat stress, grow-finish pigs alter their water usage pattern with a peak between 8 to 9 a.m. and second peak around 5 to 8 p.m. Daily drinking water needs for pigs range from less than 0.5 gal/pig/day for newly weaned pigs to greater than 1.5 gal/pig/day for grow-finish pigs using nipple drinkers. Water requirements for breeding swine range from 3 to 4 gal/day for gestating females and 6 gal/day for lactating swine. Knowledge of the daily water needs of pigs, and the patterns of water usage within the day allow for the appropriate sizing of delivery devices and prediction of the impact of pork production on available water supplies. Daily charting of drinking water usage can serve as a predictor of the on-set of swine health challenges such as swine influenza. As more sophisticated methods

become available to record water usage, other predictors of performance may be developed depending on the patterns detected.

Introduction

With the on-going drought in central and western Nebraska and the controversy surrounding the environmental impact of pork production facilities, a basic understanding of the water usage patterns in pork production facilities is important. In addition, deviations from normal patterns may be a predictor of health and future performance.

How much water does a pig drink?

Daily drinking water needs for pigs range from less than 0.5 gal/pig/day for newly weaned pigs to greater than 1.5 gal/pig/day for grow-finish pigs using nipple drinkers in warm conditions. Grow-finish pigs using bowl/cup drinkers or wet/dry feeders use less water, generally averaging just over 1.0 gal/pig/day. Water requirements for the breeding herd range from 3 to 4 gal/day for the gestating female to 5 to 6 gal/day for the lactating female.

Using the above numbers, it is possible to predict the yearly water

usage by various pork production facilities. For example, a 1,000 head grow-finish facility typically has a pen space utilization rate of 85-90%. That is, there are pigs occupying pen spaces 310 to 330 days per year. If the facility has nipple drinkers and a 90% facility utilization rate, total drinking water use for the facility will be:

$$1,000 \text{ spaces} \times 330 \text{ days/year} \\ \times 1.5 \text{ gal/space/day} = 495,000 \\ \text{gal}$$

While 495,000 gallons of water seems like a big number, when compared to the water used for irrigated crop production, it is minor. An acre-inch of water (an inch of water covering an acre of ground) is equivalent to 27,154 gallons of water. This means the example finisher will use just over 18 acre-inches of water.

If drinkers that have been proven to waste less water are used such as bowl drinkers or wet/dry feeders, total drinking water use for the facility is estimated to be:

$$1,000 \text{ spaces} \times 330 \text{ days/year} \\ \times 1.05 \text{ gal/space/day} = 346,500 \\ \text{gal}$$

This equates to 12.8 acre-inches of water.