

June 2008

Measurement of the ratio of the $pp \rightarrow W+c$ -jet cross section to the inclusive $p^-p \rightarrow W^+$ jets cross section

V. M. Abazov

Joint Institute for Nuclear Research, Dubna, Russia

Kenneth A. Bloom

University of Nebraska - Lincoln, kbloom2@unl.edu

Gregory Snow

University of Nebraska - Lincoln, gsnow1@unl.edu

D0 Collaboration

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsbloom>

 Part of the [Physics Commons](#)

Abazov, V. M.; Bloom, Kenneth A.; Snow, Gregory; and Collaboration, D0, "Measurement of the ratio of the $pp \rightarrow W+c$ -jet cross section to the inclusive $p^-p \rightarrow W^+$ jets cross section" (2008). *Kenneth Bloom Publications*. 245.
<http://digitalcommons.unl.edu/physicsbloom/245>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Kenneth Bloom Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Measurement of the ratio of the $p\bar{p} \rightarrow W + c$ -jet cross section to the inclusive $p\bar{p} \rightarrow W + \text{jets}$ cross section

DØ Collaboration

V.M. Abazov^{aj}, B. Abbott^{bw}, M. Abolins^{bm}, B.S. Acharya^{ac}, M. Adams^{ay}, T. Adams^{aw}, E. Aguilo^f, S.H. Ahn^{ae}, M. Ahsan^{bg,*}, G.D. Alexeev^{aj}, G. Alkhazov^{an}, A. Alton^{bl,1}, G. Alverson^{bk}, G.A. Alves^b, M. Anastasoiaie^{ai}, L.S. Ancu^{ai}, T. Andeen^{ba}, S. Anderson^{as}, B. Andrieu^q, M.S. Anzels^{ba}, M. Aoki^{ax}, Y. Arnoudⁿ, M. Arov^{bh}, M. Arthaud^r, A. Askew^{aw}, B. Åsman^{ao}, A.C.S. Assis Jesus^c, O. Atramentov^{aw}, C. Avila^h, C. Ay^x, F. Badaud^m, A. Baden^{bi}, L. Bagby^{ax}, B. Baldin^{ax}, D.V. Bandurin^{bg}, P. Banerjee^{ac}, S. Banerjee^{ac}, E. Barberis^{bk}, A.-F. Barfuss^o, P. Bargassa^{cb}, P. Baringer^{bf}, J. Barreto^b, J.F. Bartlett^{ax}, U. Bassler^r, D. Bauer^{aq}, S. Beale^f, A. Bean^{bf}, M. Begalli^c, M. Begel^{bu}, C. Belanger-Champagne^{ao}, L. Bellantoni^{ax}, A. Bellavance^{ax}, J.A. Benitez^{bm}, S.B. Beri^{aa}, G. Bernardi^q, R. Bernhard^w, I. Bertram^{ap}, M. Besançon^r, R. Beuselinck^{aq}, V.A. Bezzubov^{am}, P.C. Bhat^{ax}, V. Bhatnagar^{aa}, C. Biscarat^t, G. Blazey^{az}, F. Blekman^{aq}, S. Blessing^{aw}, D. Bloch^s, K. Bloom^{bo}, A. Boehnlein^{ax}, D. Boline^{bj}, T.A. Bolton^{bg}, G. Borissov^{ap}, T. Bose^{by}, A. Brandt^{bz}, R. Brock^{bm}, G. Brooijmans^{br}, A. Bross^{ax}, D. Brown^{cc}, N.J. Buchanan^{aw}, D. Buchholz^{ba}, M. Buehler^{cc}, V. Buescher^v, V. Bunichev^{al}, S. Burdin^{ap,2}, S. Burke^{as}, T.H. Burnett^{cd}, C.P. Buszello^{aq}, J.M. Butler^{bj}, P. Calfayan^y, S. Calvet^p, J. Cammin^{bs}, W. Carvalho^c, B.C.K. Casey^{ax}, H. Castilla-Valdez^{ag}, S. Chakrabarti^r, D. Chakraborty^{az}, K. Chan^f, K.M. Chan^{bc}, A. Chandra^{av}, F. Charles^{s,*}, E. Cheu^{as}, F. Chevallierⁿ, D.K. Cho^{bj}, S. Choi^{af}, B. Choudhary^{ab}, L. Christofek^{by}, T. Christoudias^{aq}, S. Cihangir^{ax}, D. Claes^{bo}, Y. Coadou^f, M. Cooke^{cb}, W.E. Cooper^{ax}, M. Corcoran^{cb}, F. Couderc^r, M.-C. Cousinou^o, S. Crépe-Renaudinⁿ, D. Cutts^{by}, M. Ćwiok^{ad}, H. da Motta^b, A. Das^{as}, G. Davies^{aq}, K. De^{bz}, S.J. de Jong^{ai}, E. De La Cruz-Burelo^{bl}, C. De Oliveira Martins^c, J.D. Degenhardt^{bl}, F. Déliot^r, M. Demarteau^{ax}, R. Demina^{bs}, D. Denisov^{ax}, S.P. Denisov^{am}, S. Desai^{ax}, H.T. Diehl^{ax}, M. Diesburg^{ax}, A. Dominguez^{bo}, H. Dong^{bt}, L.V. Dudko^{al}, L. Duflost^p, S.R. Dugad^{ac}, D. Duggan^{aw}, A. Duperrin^o, J. Dyer^{bm}, A. Dyshkant^{az}, M. Eads^{bo}, D. Edmunds^{bm}, J. Ellison^{av}, V.D. Elvira^{ax}, Y. Enari^{by}, S. Eno^{bi}, P. Ermolov^{al}, H. Evans^{bb}, A. Evdokimov^{bu}, V.N. Evdokimov^{am}, A.V. Ferapontov^{bg}, T. Ferbel^{bs}, F. Fiedler^x, F. Filthaut^{ai}, W. Fisher^{ax}, H.E. Fisk^{ax}, M. Fortner^{az}, H. Fox^{ap}, S. Fu^{ax}, S. Fuess^{ax}, T. Gadfort^{br}, C.F. Galea^{ai}, E. Gallas^{ax}, C. Garcia^{bs}, A. Garcia-Bellido^{cd}, V. Gavrillov^{ak}, P. Gay^m, W. Geist^s, D. Gelé^s, C.E. Gerber^{ay}, Y. Gershtein^{aw}, D. Gillberg^f, G. Ginter^{bs}, N. Gollub^{ao}, B. Gómez^h, A. Goussiou^{cd}, P.D. Grannis^{bt}, H. Greenlee^{ax}, Z.D. Greenwood^{bh}, E.M. Gregores^d, G. Grenier^t, Ph. Gris^m, J.-F. Grivaz^p, A. Grohsjean^y, S. Grünendahl^{ax}, M.W. Grünewald^{ad}, F. Guo^{bt}, J. Guo^{bt}, G. Gutierrez^{ax}, P. Gutierrez^{bw}, A. Haas^{br}, N.J. Hadley^{bi}, P. Haefner^y, S. Hagopian^{aw}, J. Haley^{bp}, I. Hall^{bm}, R.E. Hall^{au}, L. Han^g, K. Harder^{ar}, A. Harel^{bs}, R. Harrington^{bk}, J.M. Hauptman^{be}, R. Hauser^{bm}, J. Hays^{aq}, T. Hebbeker^u, D. Hedin^{az}, J.G. Hegeman^{ah}, J.M. Heinmiller^{ay}, A.P. Heinson^{av}, U. Heintz^{bj}, C. Hensel^{bf}, K. Herner^{bt}, G. Hesketh^{bk}, M.D. Hildreth^{bc}, R. Hirosky^{cc}, J.D. Hobbs^{bt}, B. Hoeneisen^l, H. Hoeth^z, M. Hohlfield^v, S.J. Hong^{ae}, S. Hossain^{bw}, P. Houben^{ah}, Y. Hu^{bt}, Z. Hubacek^j, V. Hynekⁱ, I. Iashvili^{bq}, R. Illingworth^{ax}, A.S. Ito^{ax}, S. Jabeen^{bj}, M. Jaffré^p, S. Jain^{bw}, K. Jakobs^w, C. Jarvis^{bi}, R. Jesik^{aq}, K. Johns^{as}, C. Johnson^{br}, M. Johnson^{ax}, A. Jonckheere^{ax}, P. Jonsson^{aq}, A. Juste^{ax}, E. Kajfasz^o, A.M. Kalinin^{aj}, J.M. Kalk^{bh}, S. Kappler^u, D. Karmanov^{al}, P.A. Kasper^{ax}, I. Katsanos^{br}, D. Kau^{aw}, V. Kaushik^{bz}, R. Kehoe^{ca}, S. Kermiche^o, N. Khalatyan^{ax}, A. Khanov^{bx}, A. Kharchilava^{bq}, Y.M. Kharzheev^{aj}, D. Khatidze^{br}, T.J. Kim^{ae}, M.H. Kirby^{ba}, M. Kirsch^u, B. Klima^{ax}, J.M. Kohli^{aa}, J.-P. Konrath^w, V.M. Korablev^{am}, A.V. Kozelov^{am}, J. Kraus^{bm}, D. Krop^{bb}, T. Kuhl^x, A. Kumar^{bq}, A. Kupco^k, T. Kurča^t, J. Kvitaⁱ, F. Lacroix^m, D. Lam^{bc}, S. Lammers^{br}, G. Landsberg^{by}, P. Lebrun^t, W.M. Lee^{ax}, A. Leflat^{al}, J. Lellouch^q, J. Leveque^{as}, J. Li^{bz}, L. Li^{av}, Q.Z. Li^{ax}, S.M. Lietti^e, J.G.R. Lima^{az}, D. Lincoln^{ax}, J. Linnemann^{bm}, V.V. Lipaev^{am}, R. Lipton^{ax}, Y. Liu^g, Z. Liu^f, A. Lobodenko^{an}, M. Lokajicek^k, P. Love^{ap}, H.J. Lubatti^{cd}, R. Luna^c, A.L. Lyon^{ax}, A.K.A. Maciel^b, D. Mackin^{cb}, R.J. Madaras^{at}, P. Mättig^z, C. Magass^u, A. Magerkurth^{bl}, P.K. Mal^{cd}, H.B. Malbouisson^c, S. Malik^{bo}, V.L. Malyshev^{aj}, H.S. Mao^{ax}, Y. Maravin^{bg}, B. Martinⁿ, R. McCarthy^{bt}, A. Melnitchouk^{bn}, L. Mendoza^h, P.G. Mercadante^e

M. Merkin^{al}, K.W. Merritt^{ax}, A. Meyer^u, J. Meyer^{v,3}, T. Millet^t, J. Mitrevski^{br}, J. Molina^c, R.K. Mommsen^{ar}, N.K. Mondal^{ac}, R.W. Moore^f, T. Moulik^{bf}, G.S. Muanza^t, M. Mulders^{ax}, M. Mulhearn^{br}, O. Mundal^v, L. Mundim^c, E. Nagy^o, M. Naimuddin^{ax}, M. Narain^{by}, N.A. Naumann^{ai}, H.A. Neal^{bl}, J.P. Negret^h, P. Neustroev^{an}, H. Nilsen^w, H. Nogima^c, S.F. Novaes^e, T. Nunnemann^y, V. O'Dell^{ax}, D.C. O'Neil^f, G. Obrant^{an}, C. Ochando^p, D. Onoprienko^{bg}, N. Oshima^{ax}, N. Osman^{aq}, J. Osta^{bc}, R. Otec^j, G.J. Otero y Garzón^{ax}, M. Owen^{ar}, P. Padley^{cb}, M. Pangilinan^{by}, N. Parashar^{bd}, S.-J. Park^{bs}, S.K. Park^{ae}, J. Parsons^{br}, R. Partridge^{by}, N. Parua^{bb}, A. Patwa^{bu}, G. Pawloski^{cb}, B. Penning^w, M. Perfilov^{al}, K. Peters^{ar}, Y. Peters^z, P. Pétroff^p, M. Petteni^{aq}, R. Piegaia^a, J. Piper^{bm}, M.-A. Pleier^v, P.L.M. Podesta-Lerma^{ag,4}, V.M. Podstavkov^{ax}, Y. Pogorelov^{bc}, M.-E. Pol^b, P. Polozov^{ak}, B.G. Pope^{bm}, A.V. Popov^{am}, C. Potter^f, W.L. Prado da Silva^c, H.B. Prosper^{aw}, S. Protopopescu^{bu}, J. Qian^{bl}, A. Quadt^{v,3}, B. Quinn^{bn}, A. Rakitine^{ap}, M.S. Rangel^b, K. Ranjan^{ab}, P.N. Ratoff^{ap}, P. Renkel^{ca}, S. Reucroft^{bk}, P. Rich^{ar}, J. Rieger^{bb}, M. Rijssenbeek^{bt}, I. Ripp-Baudot^s, F. Rizatdinova^{bx}, S. Robinson^{aq}, R.F. Rodrigues^c, M. Rominsky^{bw}, C. Royon^r, P. Rubinov^{ax}, R. Ruchti^{bc}, G. Safronov^{ak}, G. Sajotⁿ, A. Sánchez-Hernández^{ag}, M.P. Sanders^q, A. Santoro^c, G. Savage^{ax}, L. Sawyer^{bh}, T. Scanlon^{aq}, D. Schaile^y, R.D. Schamberger^{bt}, Y. Scheglov^{an}, H. Schellman^{ba}, T. Schliephake^z, C. Schwanenberger^{ar}, A. Schwartzman^{bp}, R. Schwienhorst^{bm}, J. Sekaric^{aw}, H. Severini^{bw}, E. Shabalina^{ay}, M. Shamim^{bg}, V. Shary^r, A.A. Shchukin^{am}, R.K. Shivpuri^{ab}, V. Siccaldi^s, V. Simak^j, V. Sirotenko^{ax}, P. Skubic^{bw}, P. Slattery^{bs}, D. Smirnov^{bc}, G.R. Snow^{bo}, J. Snow^{bv}, S. Snyder^{bu}, S. Söldner-Rembold^{ar}, L. Sonnenschein^q, A. Sopczak^{ap}, M. Sosebee^{bz}, K. Soustruznikⁱ, B. Spurlock^{bz}, J. Starkⁿ, J. Steele^{bh}, V. Stolin^{ak}, D.A. Stoyanova^{am}, J. Strandberg^{bl}, S. Strandberg^{ao}, M.A. Strang^{bq}, E. Strauss^{bt}, M. Strauss^{bw}, R. Ströhmer^y, D. Strom^{ba}, L. Stutte^{ax}, S. Sumowidagdo^{aw}, P. Svoisky^{bc}, A. Sznajder^c, P. Tamburello^{as}, A. Tanasijczuk^a, W. Taylor^f, J. Temple^{as}, B. Tiller^y, F. Tissandier^m, M. Titov^r, V.V. Tokmenin^{aj}, T. Toole^{bi}, I. Torchiani^w, T. Trefzger^x, D. Tsybychev^{bt}, B. Tuchming^r, C. Tully^{bp}, P.M. Tuts^{br}, R. Unalan^{bm}, L. Uvarov^{an}, S. Uvarov^{an}, S. Uzunyan^{az}, B. Vachon^f, P.J. van den Berg^{ah}, R. Van Kooten^{bb}, W.M. van Leeuwen^{ah}, N. Varelas^{ay}, E.W. Varnes^{as}, I.A. Vasilyev^{am}, M. Vaupel^z, P. Verdier^t, L.S. Vertogradov^{aj}, M. Verzocchi^{ax}, F. Villeneuve-Seguié^{aq}, P. Vint^{aq}, P. Vokac^j, E. Von Toerne^{bg}, M. Voutilainen^{bp,5}, R. Wagner^{bp}, H.D. Wahl^{aw}, L. Wang^{bi}, M.H.L.S. Wang^{ax}, J. Warchol^{bc}, G. Watts^{cd}, M. Wayne^{bc}, G. Weber^x, M. Weber^{ax}, L. Welty-Rieger^{bb}, A. Wenger^{w,6}, N. Wermes^v, M. Wetstein^{bi}, A. White^{bz}, D. Wicke^z, G.W. Wilson^{bf}, S.J. Wimpenny^{av}, M. Wobisch^{bh}, D.R. Wood^{bk}, T.R. Wyatt^{ar}, Y. Xie^{by}, S. Yacoub^{ba}, R. Yamada^{ax}, M. Yan^{bi}, T. Yasuda^{ax}, Y.A. Yatsunenko^{aj}, K. Yip^{bu}, H.D. Yoo^{by}, S.W. Youn^{ba}, J. Yu^{bz}, A. Zatserklyaniy^{az}, C. Zeitnitz^z, T. Zhao^{cd}, B. Zhou^{bl}, J. Zhu^{bt}, M. Zielinski^{bs}, D. Zieminska^{bb}, A. Zieminski^{bb,x}, L. Zivkovic^{br}, V. Zutshi^{az}, E.G. Zverev^{al}

^a Universidad de Buenos Aires, Buenos Aires, Argentina

^b LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

^c Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

^d Universidade Federal do ABC, Santo André, Brazil

^e Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil

^f University of Alberta, Edmonton, Alberta, Simon Fraser University, Burnaby, British Columbia, York University, Toronto, Ontario, and McGill University, Montreal, Quebec, Canada

^g University of Science and Technology of China, Hefei, People's Republic of China

^h Universidad de los Andes, Bogotá, Colombia

ⁱ Center for Particle Physics, Charles University, Prague, Czech Republic

^j Czech Technical University, Prague, Czech Republic

^k Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic

^l Universidad San Francisco de Quito, Quito, Ecuador

^m LPC, Univ Blaise Pascal, CNRS/IN2P3, Clermont, France

ⁿ LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, France

^o CPPM, IN2P3/CNRS, Université de la Méditerranée, Marseille, France

^p LAL, Université Paris-Sud, IN2P3/CNRS, Orsay, France

^q LPNHE, IN2P3/CNRS, Universités Paris VI and VII, Paris, France

^r DAPNIA/Service de Physique des Particules, CEA, Saclay, France

^s IPHC, Université Louis Pasteur et Université de Haute Alsace, CNRS/IN2P3, Strasbourg, France

^t IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, and Université de Lyon, Lyon, France

^u III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany

^v Physikalisches Institut, Universität Bonn, Bonn, Germany

^w Physikalisches Institut, Universität Freiburg, Freiburg, Germany

^x Institut für Physik, Universität Mainz, Mainz, Germany

^y Ludwig-Maximilians-Universität München, München, Germany

^z Fachbereich Physik, University of Wuppertal, Wuppertal, Germany

^{aa} Panjab University, Chandigarh, India

^{ab} Delhi University, Delhi, India

^{ac} Tata Institute of Fundamental Research, Mumbai, India

^{ad} University College Dublin, Dublin, Ireland

^{ae} Korea Detector Laboratory, Korea University, Seoul, Republic of Korea

^{af} SungKyunKwan University, Suwon, Republic of Korea

^{ag} CINVESTAV, Mexico City, Mexico

^{ah} FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands

^{ai} Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands

^{aj} Joint Institute for Nuclear Research, Dubna, Russia

^{ak} Institute for Theoretical and Experimental Physics, Moscow, Russia

^{al} Moscow State University, Moscow, Russia

- ^{am} Institute for High Energy Physics, Protvino, Russia
^{an} Petersburg Nuclear Physics Institute, St. Petersburg, Russia
^{ao} Lund University, Lund, Royal Institute of Technology and Stockholm University, Stockholm, and Uppsala University, Uppsala, Sweden
^{ap} Lancaster University, Lancaster, United Kingdom
^{aq} Imperial College, London, United Kingdom
^{ar} University of Manchester, Manchester, United Kingdom
^{as} University of Arizona, Tucson, AZ 85721, USA
^{at} Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA
^{au} California State University, Fresno, CA 93740, USA
^{av} University of California, Riverside, CA 92521, USA
^{aw} Florida State University, Tallahassee, FL 32306, USA
^{ax} Fermi National Accelerator Laboratory, Batavia, IL 60510, USA
^{ay} University of Illinois at Chicago, Chicago, IL 60607, USA
^{az} Northern Illinois University, DeKalb, IL 60115, USA
^{ba} Northwestern University, Evanston, IL 60208, USA
^{bb} Indiana University, Bloomington, IN 47405, USA
^{bc} University of Notre Dame, Notre Dame, IN 46556, USA
^{bd} Purdue University Calumet, Hammond, IN 46323, USA
^{be} Iowa State University, Ames, IA 50011, USA
^{bf} University of Kansas, Lawrence, KS 66045, USA
^{bg} Kansas State University, Manhattan, KS 66506, USA
^{bh} Louisiana Tech University, Ruston, LA 71272, USA
^{bi} University of Maryland, College Park, MD 20742, USA
^{bj} Boston University, Boston, MA 02215, USA
^{bk} Northeastern University, Boston, MA 02115, USA
^{bl} University of Michigan, Ann Arbor, MI 48109, USA
^{bm} Michigan State University, East Lansing, MI 48824, USA
^{bn} University of Mississippi, University, MS 38677, USA
^{bo} University of Nebraska, Lincoln, NE 68588, USA
^{bp} Princeton University, Princeton, NJ 08544, USA
^{bq} State University of New York, Buffalo, NY 14260, USA
^{br} Columbia University, New York, NY 10027, USA
^{bs} University of Rochester, Rochester, NY 14627, USA
^{bt} State University of New York, Stony Brook, NY 11794, USA
^{bu} Brookhaven National Laboratory, Upton, NY 11973, USA
^{bv} Langston University, Langston, OK 73050, USA
^{bw} University of Oklahoma, Norman, OK 73019, USA
^{bx} Oklahoma State University, Stillwater, OK 74078, USA
^{by} Brown University, Providence, RI 02912, USA
^{bz} University of Texas, Arlington, TX 76019, USA
^{ca} Southern Methodist University, Dallas, TX 75275, USA
^{cb} Rice University, Houston, TX 77005, USA
^{cc} University of Virginia, Charlottesville, VA 22901, USA
^{cd} University of Washington, Seattle, WA 98195, USA

ARTICLE INFO

Article history:

Received 15 March 2008
 Received in revised form 19 May 2008
 Accepted 26 June 2008
 Available online 4 July 2008
 Editor: H. Weerts

PACS:

12.15.Ji
 12.38.Qk
 13.85.Ni
 13.85.Qk
 14.70.Fm
 14.65.Dw

ABSTRACT

We present a measurement of the fraction of inclusive W + jets events produced with net charm quantum number ± 1 , denoted $W + c$ -jet, in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using approximately 1 fb^{-1} of data collected by the DØ detector at the Fermilab Tevatron Collider. We identify the W + jets events via the leptonic W boson decays. Candidate $W + c$ -jet events are selected by requiring a jet containing a muon in association with a reconstructed W boson and exploiting the charge correlation between this muon and W boson decay lepton to perform a nearly model-independent background subtraction. We measure the fraction of $W + c$ -jet events in the inclusive W + jets sample for jet $p_T > 20$ GeV and pseudorapidity $|\eta| < 2.5$ to be $0.074 \pm 0.019(\text{stat.}) \pm_{0.014}^{0.012}(\text{syst.})$, in agreement with theoretical predictions. The probability that background fluctuations could produce the observed fraction of $W + c$ -jet events is estimated to be 2.5×10^{-4} , which corresponds to a 3.5σ statistical significance.

Published by Elsevier B.V.

* Corresponding author.

E-mail address: mahsana@phys.ksu.edu (M. Ahsan).

¹ Visitor from Augustana College, Sioux Falls, SD, USA.

² Visitor from The University of Liverpool, Liverpool, UK.

³ Visitor from II. Physikalisches Institut, Georg-August-University, Göttingen, Germany.

⁴ Visitor from ICN-UNAM, Mexico City, Mexico.

⁵ Visitor from Helsinki Institute of Physics, Helsinki, Finland.

⁶ Visitor from Universität Zürich, Zürich, Switzerland.

✘ Deceased.

In hadron–hadron collisions, the $W/Z + b$ - or c -jet final state can signal the presence of new physics; however, only a few measurements [1–3] of cross sections for these Standard Model processes exist. Charm quark production in association with a W boson can be a significant background, for example, to measurements of single top, $t\bar{t}$ pair, and Higgs boson production, and to supersymmetric top quark (stop) pair production when only the $\tilde{t} \rightarrow c\tilde{\chi}_1^0$ decay channel is allowed by the mass difference between the stop quark and the neutralino. Moreover, as the squared Cabibbo–Kobayashi–Maskawa matrix element, $|V_{cd}|^2$, suppresses the expected leading order d quark–gluon fusion production mech-

anism, $W + c$ -quark production provides direct sensitivity to the proton's s quark parton distribution function (PDF), $s(x, Q^2)$, where x is the momentum fraction of the proton carried by the s -quark and $\sqrt{Q^2}$ is the hard scatter scale [4]. This PDF has been measured directly only in fixed target neutrino-nucleon deep inelastic scattering experiments using relatively low momentum transfer squared, Q^2 , of the order 1–100 GeV² [5–11]. A probe of the s quark PDF at the Tevatron tests the universality of $s(x, Q^2)$ and its QCD evolution up to $Q^2 = 10^4$ GeV². The strange quark PDF initiates both Standard Model (e.g., $sg \rightarrow W^- + c$) and possible new physics processes (e.g., $s\bar{c} \rightarrow H^-$) [12] at both the Fermilab Tevatron and CERN LHC colliders.

In this Letter, we present a measurement of the cross section ratio $\sigma(p\bar{p} \rightarrow W + c\text{-jet})/\sigma(p\bar{p} \rightarrow W + \text{jets})$ as a function of jet transverse momentum p_T , where $W + c\text{-jet}$ denotes a W boson plus jets final state in which the jets have a net charm quantum number $C = \pm 1$, and $W + \text{jets}$ denotes any W boson final state with at least one jet. Several experimental uncertainties (e.g., luminosity, jet energy scale, and reconstruction efficiency) and theoretical uncertainties (e.g., renormalization and factorization scales) largely cancel in this ratio.

This measurement utilizes approximately 1 fb⁻¹ of $p\bar{p}$ collision data at a center-of-mass energy $\sqrt{s} = 1.96$ TeV collected with the DØ detector at the Fermilab Tevatron collider. We identify W bosons through their leptonic decays, $W \rightarrow \ell\nu$, where $\ell = e, \mu$. W bosons decaying to tau leptons are included for leptonic tau decays $\tau \rightarrow e\bar{\nu}_e\nu_\tau$ or $\tau \rightarrow \mu\bar{\nu}_\mu\nu_\tau$. The electron or muon from W boson decays is required to be isolated, and its transverse momentum p_T must satisfy $p_T > 20$ GeV. The presence of a neutrino is inferred from the requirement that the missing transverse energy \cancel{E}_T satisfies $\cancel{E}_T > 20$ GeV. Jets are defined using the iterative seed-based midpoint cone algorithm [13] with cone radius of 0.5. We restrict the transverse momentum of the jet to $p_T > 20$ GeV after it is calibrated for the calorimeter jet energy scale (JES), and its pseudorapidity to $|\eta| < 2.5$, where $\eta = -\ln[\tan(\theta/2)]$ and θ is the polar angle with respect to the proton beam direction. We correct the jet measurement to the particle level [14] for comparison with theory.

A muon reconstructed within a jet (a “jet-muon”) identifies that jet (a “ μ -tagged jet”) as a charm quark candidate. Events containing a jet-muon enrich a sample in b/c semileptonic decays. Events with the jet-muon's charge opposite to or equal to that of the W boson are denoted as “OS” or “SS” events, respectively. In the $gs \rightarrow Wc$ process charge conservation only allows the W^-c or $W^+\bar{c}$ pairings. Therefore the muon from the charm quark carries an opposite-sign charge compared to the W boson's charge and the numbers of OS and SS events, N_{OS} and N_{SS} , satisfy $N_{OS} \gg N_{SS}$. N_{SS} can be non-zero because a jet initiated by a c quark has a small probability of containing a muon from the decay of particles other than the leading charm quark. Other vector boson + jets physics processes ($W + g$, $W + c\bar{c}$, $W + b\bar{b}$, $Z + \text{jets}$) can produce μ -tagged jets, but the charge of the jet-muon is uncorrelated with that of the boson, hence $N_{OS} \approx N_{SS}$ for these sources. Processes with light-quark (u , d or s) initiated jets recoiling against the W boson can produce a small fraction of charge-correlated μ -tagged jets owing to leading particle effects [15]. While charge correlation effects are large in the WW production, their contributions to the background remain minor owing to the small WW cross section. The WZ and ZZ processes only rarely produce charge-correlated jets. Other final states that can produce charge-correlated jets ($t\bar{t}$, $t\bar{b}$, $W + b\bar{c}$ and $W + b$) are suppressed by small production cross sections or tiny CKM matrix elements. These considerations allow a measurement of the $W + c\text{-jet}$ production rate from OS events with the backgrounds determined *in situ* from SS events, up to small weakly model-dependent theory corrections.

The DØ detector [16] is a multi-purpose device built to investigate $p\bar{p}$ collisions. The innermost silicon microstrip detectors followed by the scintillating fiber tracking detector, covering pseudorapidity $|\eta| \lesssim 3.0$ and located inside the 2 T superconducting solenoid, are used for tracking and vertexing. The liquid-argon and uranium calorimeter, a finely segmented detector in the transverse and the longitudinal directions, is used as the primary tool to reconstruct electrons, jets, and the missing transverse energy. It is housed in three cryostats, one central calorimeter in the region $|\eta| < 1.1$ and two end caps extending the coverage to $|\eta| \approx 4.0$. The outermost subsystem of the DØ detector is the muon spectrometer, consisting of three layers of muon tracking subdetectors and scintillation trigger counters, which is used to construct muons up to $|\eta| \approx 2.0$. The first layer is situated before the 1.8 T iron toroid and the other two layers are outside, enclosing the detector.

Candidate events in the electron (muon) decay channel of the W boson must pass at least one of the single electron (muon) three-level (L1, L2 and L3) triggers used in each data taking period. Each level of trigger imposes tighter restrictions on the events compared to those of the preceding level. The single muon triggers at L1 impose hit requirements in the muon scintillators. Some of the triggers also require spatially matched hits in the muon tracking detectors. The conditions at L2 require a reconstructed muon with p_T above a threshold in the range 0–5 GeV for various triggers. At L3, certain triggers require a track reconstructed in the inner tracking system with $p_T > 10$ GeV. The ratio measurement benefits from full cancellation of the trigger efficiency in the electron channel. This cancellation is partial in the muon channel due to the presence of two muons in the $W + c\text{-jet}$ sample.

Selection of $W \rightarrow e\nu$ candidates begins with the requirement that a cluster of energy be found that is consistent with the presence of an electron in the calorimeter. The cluster must: have at least 90% of its energy contained in the electromagnetic part of the calorimeter; have a reconstructed track from the inner tracking system pointing to it; be isolated from other clusters in that the fraction of the energy deposited in an annulus ($0.2 < \Delta\mathcal{R} = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} < 0.4$, where ϕ is the azimuthal angle) around the EM cluster is less than 15% of the electromagnetic energy within the cone of radius $\Delta\mathcal{R} = 0.2$; have longitudinal and transverse energy deposition profiles consistent with those expected for an electron; and satisfy a likelihood discriminant selection that combines tracker and calorimeter information using the expected distributions for electrons and jet background. The electron track's point of closest approach to the z -axis must be within 3 cm of the $p\bar{p}$ interaction point, which must lie within 60 cm of the nominal detector center.

Selection of $W \rightarrow \mu\nu$ candidates begins by requiring that a muon candidate be found in the muon spectrometer with a track matched to one found in the central tracker. Rejection of cosmic ray background events demands that the central tracker track pass within 0.02 or 0.2 cm of the beam crossing point in the transverse plane, depending on whether the track is reconstructed with or without hits, respectively, in the silicon detector, and that the point of closest approach of the track should be within 3 cm of the interaction point along the z -axis. Further cosmic ray rejection comes from scintillator timing information in the muon spectrometer. Requiring the W boson candidate muon track to be separated from the axis formed by any jet found in the event by $\Delta\mathcal{R}(\mu, \text{jet}) > 0.5$ suppresses backgrounds from semileptonic decays of heavy flavor quarks in multi-jet events.

For the final selection in both channels, each event must satisfy the transverse mass requirement $40 \leq M_T \leq 120$ GeV, where $M_T = \sqrt{2\cancel{E}_T p_T^\ell [1 - \cos \Delta\phi(\cancel{E}_T, p_T^\ell)]}$ is computed from the isolated lepton p_T^ℓ and the \cancel{E}_T . The azimuthal angular separation between

the isolated lepton and \cancel{E}_T directions must be greater than 0.4. Events must contain at least one jet with $p_T > 20$ GeV after the calorimeter JES correction is applied, and $|\eta| < 2.5$. Upon application of all selection criteria, $N_{Wj}^e = 82747$ and $N_{Wj}^\mu = 57944$ W + jets candidates remain in the electron and muon channels, respectively.

Backgrounds originate from photons and jets that are misidentified as electrons and from $c\bar{c}$ and $b\bar{b}$ multi-jet events that produce an isolated muon. These multi-jet backgrounds are determined directly from the data using a “matrix method”. According to this procedure “loose” $W(\rightarrow \ell\nu) + \text{jets}$ datasets are selected by applying all previously described selection criteria in each channel with the exception that the lepton isolation requirements are loosened. This produces a set of loose candidate events, N_L^ℓ , in each lepton channel consisting of a mixture of real W + jets events, N_W^ℓ , and multi-jet background events, N_{MJ}^ℓ , with $N_L^\ell = N_W^\ell + N_{MJ}^\ell$. Application of the stricter lepton isolation criteria used to extract the signal changes this mixture to $N_T^\ell = \epsilon_W^\ell N_W^\ell + \epsilon_{MJ}^\ell N_{MJ}^\ell$, where N_T^ℓ signifies the number of events in each channel satisfying the tighter isolation criteria and ϵ_W^ℓ and ϵ_{MJ}^ℓ denote the relative probabilities for loosely selected W boson and multi-jet events, respectively, to satisfy the stricter isolation criteria. A large sample of two-jet events is used to measure ϵ_{MJ}^ℓ , and ϵ_{MJ}^ℓ is estimated from a similar two-jet dataset using a sample of back-to-back muon-plus-jet events with low \cancel{E}_T . The factors ϵ_W^ℓ and ϵ_{MJ}^ℓ are obtained from large data samples of $Z \rightarrow e^+e^-$ and $Z \rightarrow \mu^+\mu^-$ events, respectively. Solving the equations $N_L^\ell = N_W^\ell + N_{MJ}^\ell$ and $N_T^\ell = \epsilon_W^\ell N_W^\ell + \epsilon_{MJ}^\ell N_{MJ}^\ell$ yields the estimates for N_W^ℓ and N_{MJ}^ℓ . The fractional contributions of multi-jet background to the inclusive W + jets signal is $f_{MJ}^e = (3.2 \pm 0.8)\%$ in the electron channel and $f_{MJ}^\mu = (4.1 \pm 3.0)\%$ in the muon channel.

The channel $Z(\rightarrow \ell^+\ell^-) + \text{jets}$ contributes as background when one of the leptons from the Z boson decay fails to be reconstructed. An estimate of this background follows from MC simulations of Z + jets production produced with ALPGEN v2.05 [17] using the CTEQ6L [18] PDF set, the PYTHIA v6.323 [19] generator for the parton fragmentation and hadronization, the MLM prescription [20] to avoid an over-counting of final state jets, and EVTGEN [21] to decay the heavy hadrons. A GEANT [22] based program simulates the effects of detector response, and the same reconstruction software is employed as used for data. This procedure results in estimates for the fractional contaminations of $f_Z^e = (0.9 \pm 0.1)\%$ for $Z(\rightarrow e^+e^-) + \text{jets}$ and $f_Z^\mu = (5.0 \pm 0.7)\%$ for $Z(\rightarrow \mu^+\mu^-) + \text{jets}$. The small background from $Z \rightarrow \tau^+\tau^-$ process contributes to the estimated fractions f_Z^e and f_Z^μ . Quoted uncertainties derive mainly from systematic effects in the Z + jets ALPGEN cross section model that are estimated by varying the relative cross section of W + jets with respect to Z + jets by its uncertainties.

Extraction of samples of W + c -jet event candidates from the W + jets samples follows from selecting events with a μ -tagged jet. This jet must contain a reconstructed muon with $p_T > 4$ GeV and $|\eta| < 2.0$ that lies within a cone of $\Delta\mathcal{R}(\mu, \text{jet}) < 0.5$ with respect to the jet axis, and have JES corrected $p_T > 20$ GeV before including the muon and neutrino energies. The muon must be detected in both of the outer two layers of the muon spectrometer, and its muon spectrometer track must be matched to a reconstructed track in the central tracker. Background suppression of $Z(\rightarrow \mu^+\mu^-) + \text{jets}$ events entails rejecting events in which the dimuon invariant mass exceeds 70 GeV in the muon channel without restricting the charges of the muons. Application of all selection criteria yields N_{OS}^e and N_{SS}^e events in the electron channel, and N_{OS}^μ and N_{SS}^μ events in the muon channel as reported in Table 1. Estimated multi-jet backgrounds in the μ -tagged jet data samples, determined using the matrix method, are $N_{OS}^{e,MJ}$ and $N_{SS}^{e,MJ}$ events

in the electron channel, and $N_{OS}^{\mu,MJ}$ and $N_{SS}^{\mu,MJ}$ events in the muon channel as listed in Table 1. Estimates of the WW , $t\bar{t}$ and single top backgrounds from MC are denoted $N_{OS}^{e,WW}$, $N_{SS}^{e,WW}$, $N_{OS}^{e,t\bar{t}}$, $N_{SS}^{e,t\bar{t}}$, $N_{OS}^{e,tb}$ and $N_{SS}^{e,tb}$ events, respectively, in the electron channel, and $N_{OS}^{\mu,WW}$, $N_{SS}^{\mu,WW}$, $N_{OS}^{\mu,t\bar{t}}$, $N_{SS}^{\mu,t\bar{t}}$, $N_{OS}^{\mu,tb}$ and $N_{SS}^{\mu,tb}$ events, respectively, in the muon channel as given in Table 1. The estimate of the single top background follows from the $t\bar{b}$ and $t\bar{b}q$ events produced with the COMPHEP [23] generator followed by full detector simulation. The quoted uncertainties on the WW , $t\bar{t}$ and single top background predictions given in Table 1 are dominated by the uncertainties in their cross section measurements [24–26]. Lepton charges are well measured at DØ, and uncertainties from charge mis-identification are very small.

The acceptance times efficiencies, ϵ_c^ℓ ($\ell = e, \mu$), of the W + c -jet selections relative to inclusive W + jets in each W boson decay channel is estimated from the MC simulation, and includes MC to data correction factors estimated using independent data calibration samples. The absolute efficiency of reconstructing a W boson with at least one jet cancels in the ratio. The relative acceptance includes effects of charm quark to hadron fragmentation, charmed hadron semi-muonic decay and the residual missing calorimeter energy from the muon and neutrino in the μ -tagged jet. The relative efficiency accounts for muon identification and track reconstruction effects. Charm quark fragmentation and charm hadron decay uncertainties are constrained by previous experiments [27, 28] and contribute 4.5% and 9.5%, respectively, to the acceptance uncertainties in both channels. The correction included in the acceptance for the missing contribution to the jet p_T from the muon and neutrino energies adjusts the jet p_T spectrum of W + c -jet candidate events appropriately to the particle level, as verified by a MC closure test. A large sample of $J/\psi \rightarrow \mu^+\mu^-$ events collected at DØ is employed to correct the jet-muon reconstruction efficiency, $(58.7 \pm 2.8)\%$, computed from the MC simulation, by a factor of 0.89 ± 0.06 . This correction is found to be independent of the jet p_T . The final acceptance times efficiencies are found to be $\epsilon_c^e = (1.24 \pm 0.22)\%$ and $\epsilon_c^\mu = (1.22 \pm 0.23)\%$.

The presence of two muons in the muon channel increases the trigger selection efficiency of the W + c -jet candidates compared to the inclusive W + jets data sample. The divisor factor $K_T^\mu = 1.18 \pm 0.12$, extracted from the probability of events being selected when only the jet-muon fires the trigger, corrects for the bias in W + c -jet selection in the muon channel. In the electron channel the factor K_T^e is unity as the trigger efficiency cancels in the ratio.

The W + c -jet cross section ratio is extracted using

$$\frac{\sigma[W(\rightarrow \ell\nu) + c\text{-jet}]}{\sigma[W(\rightarrow \ell\nu) + \text{jets}]} = \frac{\frac{1}{\epsilon_c^\ell K_T^\ell} [N_{OS}^\ell - f_c^\ell (N_{SS}^\ell - N_{SS}^{\ell,B}) - N_{OS}^{\ell,B}]}{(1 - f_Z^\ell - f_{MJ}^\ell) N_{Wj}^\ell},$$

where $N_{SS}^{\ell,B} = N_{SS}^{\ell,MJ} + N_{SS}^{\ell,WW} + N_{SS}^{\ell,t\bar{t}} + N_{SS}^{\ell,tb}$ and $N_{OS}^{\ell,B} = N_{OS}^{\ell,MJ} + N_{OS}^{\ell,WW} + N_{OS}^{\ell,t\bar{t}} + N_{OS}^{\ell,tb}$. This equation requires one further correction in each channel, f_c^ℓ , for the small correlation between the jet-muon and W boson charges that arises in W + light-quark jet events. The factor f_c^ℓ is determined from fully simulated W + jet events as the ratio of the predicted number of OS μ -tagged jets to SS μ -tagged jets in all background samples that pass the same selection criteria as defined for the data sample. Processes considered include $W + u, d, s, W + g, W + c\bar{c}, W + b\bar{b}$, and $W + c$ -jet, where the c quark does not decay semi-muonically in the last case. The f_c^ℓ are parameterized in terms of jet p_T as $f_c^\ell = a_\ell + b_\ell \times p_T$, with $a_e = 1.223 \pm 0.016$, $a_\mu = 1.241 \pm 0.023$, $b_e = -0.0017 \pm 0.0003$, and $b_\mu = -0.0019 \pm 0.0004$, where all quoted uncertainties of the parameters are statistical; f_c^ℓ decreases

Table 1Summary of quantities to estimate the $W + c$ -jet cross section ratio. The first uncertainties quoted are statistical and the second systematic

Jet p_T [GeV]	(20–30)	(30–45)	(45–200)	(20–200)
$W \rightarrow e\nu$ decay channel				
N_{Wj}^e	35695	24412	22640	82747
N_{OS}^e	83	77	85	245
N_{SS}^e	45	41	68	154
$N_{OS}^{e,MJ}$	$4.5 \pm 1.0 \pm 1.2$	$4.2 \pm 0.9 \pm 1.1$	$4.6 \pm 1.0 \pm 1.2$	$13.3 \pm 2.6 \pm 3.4$
$N_{SS}^{e,MJ}$	$5.6 \pm 1.1 \pm 1.4$	$5.1 \pm 1.0 \pm 1.3$	$8.5 \pm 1.5 \pm 2.2$	$19.3 \pm 2.9 \pm 4.9$
$N_{OS}^{e,WW}$	1.8 ± 0.6	2.1 ± 0.7	2.3 ± 0.8	6.2 ± 2.1
$N_{SS}^{e,WW}$	0.4 ± 0.1	0.6 ± 0.2	0.9 ± 0.3	1.9 ± 0.5
$N_{OS}^{e,t\bar{t}}$	2.4 ± 0.6	4.6 ± 1.1	11.8 ± 2.8	18.8 ± 4.5
$N_{SS}^{e,t\bar{t}}$	2.1 ± 0.5	4.1 ± 1.0	10.0 ± 2.4	16.1 ± 3.9
$N_{OS}^{e,tb}$	1.1 ± 0.3	2.1 ± 0.6	3.1 ± 0.9	6.3 ± 1.8
$N_{SS}^{e,tb}$	0.8 ± 0.2	1.4 ± 0.4	2.5 ± 0.7	4.6 ± 1.3
f_c^e	$1.183 \pm 0.017 \pm 0.018$	$1.164 \pm 0.019 \pm 0.017$	$1.118 \pm 0.024 \pm 0.017$	$1.149 \pm 0.007 \pm 0.017$
ϵ_c^e	$0.0113 \pm 0.0015^{+0.0017}_{-0.0017}$	$0.0125 \pm 0.0011^{+0.0019}_{-0.0019}$	$0.0125 \pm 0.0020^{+0.0019}_{-0.0019}$	$0.0124 \pm 0.0012^{+0.0019}_{-0.0019}$
$\frac{\sigma[W(\rightarrow e\nu)+c\text{-jet}]}{\sigma[W(\rightarrow e\nu)+\text{jets}]}$	$0.079 \pm 0.031^{+0.013}_{-0.022}$	$0.100 \pm 0.038^{+0.017}_{-0.016}$	$0.043 \pm 0.049^{+0.007}_{-0.007}$	$0.073 \pm 0.023^{+0.012}_{-0.014}$
$W \rightarrow \mu\nu$ decay channel				
N_{Wj}^μ	27378	17325	13241	57944
N_{OS}^μ	76	64	63	203
N_{SS}^μ	28	38	56	122
$N_{OS}^{\mu,MJ}$	$4.6 \pm 1.8 \pm 3.3$	$3.8 \pm 1.5 \pm 2.7$	$3.8 \pm 1.5 \pm 2.7$	$12.2 \pm 4.6 \pm 8.7$
$N_{SS}^{\mu,MJ}$	$2.0 \pm 1.3 \pm 1.4$	$2.8 \pm 1.7 \pm 2.0$	$4.1 \pm 2.5 \pm 2.9$	$8.8 \pm 5.4 \pm 6.3$
$N_{OS}^{\mu,WW}$	0.8 ± 0.3	1.6 ± 0.5	1.8 ± 0.6	4.2 ± 1.6
$N_{SS}^{\mu,WW}$	0.3 ± 0.1	0.3 ± 0.1	0.6 ± 0.2	1.2 ± 0.4
$N_{OS}^{\mu,t\bar{t}}$	1.2 ± 0.3	2.3 ± 0.6	5.8 ± 1.4	9.3 ± 2.2
$N_{SS}^{\mu,t\bar{t}}$	1.0 ± 0.2	2.0 ± 0.5	5.1 ± 1.2	8.1 ± 1.9
$N_{OS}^{\mu,tb}$	0.7 ± 0.2	1.4 ± 0.4	2.1 ± 0.6	4.2 ± 1.2
$N_{SS}^{\mu,tb}$	0.5 ± 0.1	0.8 ± 0.1	1.8 ± 0.5	3.1 ± 0.9
f_c^μ	$1.195 \pm 0.025 \pm 0.014$	$1.174 \pm 0.027 \pm 0.013$	$1.121 \pm 0.035 \pm 0.013$	$1.148 \pm 0.007 \pm 0.013$
ϵ_c^μ	$0.0110 \pm 0.0011^{+0.0016}_{-0.0017}$	$0.0122 \pm 0.0013^{+0.0018}_{-0.0019}$	$0.0148 \pm 0.0018^{+0.0022}_{-0.0023}$	$0.0122 \pm 0.0012^{+0.0018}_{-0.0019}$
K_T^μ	$1.18 \pm 0.02 \pm 0.12$	$1.18 \pm 0.02 \pm 0.12$	$1.18 \pm 0.02 \pm 0.12$	$1.18 \pm 0.02 \pm 0.12$
$\frac{\sigma[W(\rightarrow \mu\nu)+c\text{-jet}]}{\sigma[W(\rightarrow \mu\nu)+\text{jets}]}$	$0.123 \pm 0.037^{+0.024}_{-0.033}$	$0.076 \pm 0.050^{+0.016}_{-0.013}$	$0.000 \pm 0.058^{+0.014}_{-0.008}$	$0.075 \pm 0.031^{+0.015}_{-0.017}$
Combined $W \rightarrow e\nu$ and $W \rightarrow \mu\nu$ decay channels				
$\frac{\sigma[W(\rightarrow \ell\nu)+c\text{-jet}]}{\sigma[W(\rightarrow \ell\nu)+\text{jets}]}$	$0.097 \pm 0.024^{+0.016}_{-0.026}$	$0.091 \pm 0.031^{+0.016}_{-0.015}$	$0.025 \pm 0.038^{+0.005}_{-0.004}$	$0.074 \pm 0.019^{+0.012}_{-0.014}$

with increasing jet p_T because the sub-process $q\bar{q} \rightarrow Wg$ dominates $qg \rightarrow Wq'$ at high jet p_T . Systematic uncertainties in f_c^ℓ arise mainly from the cross section and jet fragmentation models. The f_c^ℓ are nearly independent of the absolute charged multiplicity per jet and the $W +$ light-jets cross section. This has been verified by comparing the ratio of all OS tracks to all SS tracks found in jets in the inclusive $W +$ jets data sample. The f_c^ℓ depend instead on the K^\pm/π^\pm ratio per jet and the relative cross section for W boson plus heavy quark jet final states compared to $W +$ light-jets. A 6% uncertainty is assigned to the weighted π^\pm multiplicity based on a comparison of the difference between tracking efficiency in data and simulation, and a 20% uncertainty on the K^\pm/π^\pm ratio is estimated based on comparing K_S^0 production in data to MC. Uncertainties in ALPGEN cross sections are estimated to be 50% for $W + b\bar{b}$, $W + c\bar{c}$, and $W + c$ -jet, relative to $W +$ light-jets [29]. A change of the $W + c$ -jet cross section by $\pm 100\%$ does not lead to a significant effect in f_c^ℓ . The uncertainty due to PDFs on f_c^ℓ is estimated to be $^{+0.97\%}_{-0.64\%}$. Overall systematic uncertainties are found to be 1.5% for f_c^e and 1.1% for f_c^μ , in which the relative cross section contributions dominate. Adding a 0.6% uncertainty in each channel due to MC statistics yields $f_c^e = 1.149 \pm 0.018$ and $f_c^\mu = 1.148 \pm 0.015$ averaged over all $p_T > 20$ GeV.

Table 1 summarizes the cross section ratio measurements and their uncertainties for the electron and the muon channels for events with jet $p_T > 20$ GeV and jet $|\eta| < 2.5$ as well as for

Table 2Fractional systematic uncertainties on the measurement in the $W \rightarrow e\nu$ and the $W \rightarrow \mu\nu$ channels

p_T GeV	e channel				μ channel				
	JES %	JPR %	f_c^e %	ϵ_c^e %	JES %	JPR %	f_c^μ %	ϵ_c^μ %	K_T^μ %
20–30	+0 –21.6	+2.4 –4.8	+3.8 –4.1	+15.7 –15.6	+0 –17.6	+5.0 –7.5	+2.5 –3.3	+15.3 –16.2	± 10
30–45	+6.4 –4.3	+2.1 –4.3	+4.3 –4.7	+14.5 –14.4	+9.8 –0.7	+4.5 –6.7	+3.1 –4.3	+14.1 –14.9	± 10
45–200	+2.2 –2.2	+2.2 –4.5	+6.9 –7.6	+14.7 –14.6	+27.7 –0	+4.7 –7.0	+4.0 –5.2	+15.0 –15.8	± 10
20–200	+0 –9.0	+2.3 –4.5	+4.5 –5.2	+15.1 –15.0	+5.9 –2.4	+4.7 –7.1	+3.1 –4.3	+14.9 –15.7	± 10

three different jet p_T bins. The ratio measurements benefit from cancellation of several uncertainties, notably the integrated luminosity [30], lepton detection efficiency, and jet energy scale (JES). Table 2 lists the remaining absolute systematic uncertainties on the measurement estimated from the MC simulation. These arise mainly from second order JES effects, jet p_T resolution (JPR), c -jet tagging efficiency, and the $W + c$ -jet background correction factors f_c^ℓ .

The measured $W + c$ -jet fractions integrated over $p_T > 20$ GeV and $|\eta| < 2.5$ are

$$\frac{\sigma[W(\rightarrow e\nu) + c\text{-jet}]}{\sigma[W(\rightarrow e\nu) + \text{jets}]} = 0.073 \pm 0.023(\text{stat.})^{+0.012}_{-0.014}(\text{syst.}),$$




Fig. 1. Measured ratio $[\sigma(W+c\text{-jet})/\sigma(W+\text{jets})]$ for jet $p_T > 20$ GeV and $|\eta| < 2.5$. The inner error bars around the data points show the statistical only uncertainties and the full bars represent the quadratic sum of statistical and systematic uncertainties. The systematic uncertainty on $W+c$ -jet fraction includes the uncertainties due to JES, the jet p_T resolution, the background correction factor f_c^ℓ , and the product of the relative acceptance and efficiencies ϵ_c^ℓ . It also includes the uncertainty due to K_T^μ in the muon channel. The dashed line represents the prediction from ALPGEN and PYTHIA.

$$\frac{\sigma[W(\rightarrow \mu\nu) + c\text{-jet}]}{\sigma[W(\rightarrow \mu\nu) + \text{jets}]} = 0.075 \pm 0.031(\text{stat.})_{-0.017}^{+0.015}(\text{syst.}).$$

Since the $W \rightarrow e\nu$ and $W \rightarrow \mu\nu$ measurements are consistent with one another, and statistical uncertainties dominate, the two lepton channels are combined to yield

$$\frac{\sigma[W + c\text{-jet}]}{\sigma[W + \text{jets}]} = 0.074 \pm 0.019(\text{stat.})_{-0.014}^{+0.012}(\text{syst.}).$$

Systematic uncertainties are taken to be fully correlated in the two channels. This measurement can be compared to the $W+c$ -jet fraction predicted by ALPGEN and PYTHIA of 0.044 ± 0.003 , where the quoted theoretical uncertainty derives from the uncertainty on the CTEQ6.5 PDFs. Due to the relatively small contributions of $W+bb$ and $W+c\bar{c}$ to inclusive $W+\text{jets}$, the model prediction of the $W+c$ -jet rate has $\lesssim 5\%$ sensitivity to their cross sections. Fig. 1 shows the differential $W+c$ -jet fraction, and compares the data to a model prediction using leading order QCD augmented by ALPGEN and PYTHIA.

As a test of the $W+c$ -jet signal hypothesis, Fig. 2(a) compares data to ALPGEN and PYTHIA expectations in the background-subtracted distribution of the signed impact parameter significance, a/σ_a , for the jet-muon, where a is the projected distance of closest approach of the jet-muon to the event interaction point in the transverse plane, and σ_a is the estimated uncertainty on a . Data show satisfactory agreement with expectations for $W+c$ -jet production and the underlying OS–SS ansatz after the subtraction of light and b quark jet contributions. Similarly, the distribution of the relative transverse momentum of the jet-muon with respect to the jet axis, p_T^{rel} , shows the consistency between data and the c -jet expectation as illustrated in Fig. 2(b).

To quantify the probability that the observed excess of OS events over SS events is due exclusively to background fluctuations, ensembles for OS, SS backgrounds and inclusive $W+\text{jets}$ are generated that incorporate all the systematic uncertainties together with the correlations among the OS, SS backgrounds and $W+\text{jets}$ expectations using Gaussian samplings of the uncertainties. The probability that background fluctuations could produce the observed fraction of the signal events in the inclusive $W+\text{jets}$





Fig. 2. Comparison of the background-subtracted ($N_{OS}^\ell - f_c^\ell N_{SS}^\ell$) data in the combined electron and muon channels with the simulation. (a) Signed significance in impact parameter of the jet-muon track with respect to the interaction point, (b) jet-muon transverse momentum relative to the jet axis (p_T^{rel}).

sample is 2.5×10^{-4} , corresponding to a 3.5σ significance for the $W+c$ -jet hypothesis.

In conclusion, we have performed a measurement of the $W+c$ -jet/ $W+\text{jets}$ cross section ratio at a hadron collider using both electron and muon decay channels of the W boson and utilizing the correlation between the charge of the jet-muon with that of the W boson. The probability that background fluctuations could produce an estimated $W+c$ -jet fraction in $W+\text{jets}$ equal to or larger than the one measured in data is 2.5×10^{-4} , which corresponds to a 3.5σ significance of the observation. We find our measurement to be consistent with LO perturbative QCD predictions of the $W+c$ -jet production rate and with an s quark PDF evolved from Q^2 scales two orders of magnitude below those of this measurement. The measurement further provides direct experimental evidence of the underlying partonic process $qg \rightarrow Wq'$ that should dominate W boson production at the CERN Large Hadron Collider (LHC).

Acknowledgements

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Republic of Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC (United Kingdom); MSMT and GACR (Czech Re-

public); CRC Program, CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); and the Alexander von Humboldt Foundation.

References

- [1] A. Abulencia, et al., CDF Collaboration, Phys. Rev. D 74 (2006) 032008.
- [2] V.M. Abazov, et al., DØ Collaboration, Phys. Rev. Lett. 94 (2005) 161801.
- [3] T. Aaltonen, et al., CDF Collaboration, Phys. Rev. Lett. 100 (2008) 091803.
- [4] U. Baur, et al., Phys. Lett. B 318 (1993) 544.
- [5] D.A. Mason, et al., NuTeV Collaboration, Phys. Rev. Lett. 99 (2007) 192001.
- [6] M. Goncharov, et al., NuTeV Collaboration, Phys. Rev. D 64 (2001) 112006.
- [7] S.A. Rabinowitz, et al., CCFR Collaboration, Phys. Rev. Lett. 70 (1993) 134.
- [8] A.O. Bazarko, et al., CCFR Collaboration, Z. Phys. C 65 (1995) 189.
- [9] P. Vilain, et al., Charm II Collaboration, Eur. Phys. J. C 11 (1999) 19.
- [10] H. Abramowicz, et al., CDHS Collaboration, Z. Phys. C 15 (1982) 19.
- [11] M. Holder, et al., CDHS Collaboration, Phys. Lett. B 69 (1977) 377.
- [12] H.L. Lai, et al., hep-ph/0702268.
- [13] G.C. Blazey, et al., in: U. Baur, R.K. Ellis, D. Zeppenfeld (Eds.), Proceedings of the Workshop: QCD and Weak Boson Physics in Run II, Fermilab-Pub-00/297 (2000).
- [14] C. Butter, et al., hep-ph/0803.0678.
- [15] K. Abe, et al., SLD Collaboration, Phys. Rev. Lett. 78 (1997) 3442.
- [16] V.M. Abazov, et al., DØ Collaboration, Nucl. Instrum. Methods Phys. Res. A 565 (2006) 463.
- [17] M.L. Mangano, et al., JHEP 0307 (2003) 001. ALPGEN version 2.05 is used throughout.
- [18] J. Pumplin, et al., JHEP 0207 (2002) 012; D. Stump, et al., JHEP 0310 (2003) 046.
- [19] T. Sjöstrand, et al., Comput. Phys. Commun. 135 (2001) 238. PYTHIA version 6.323 is used throughout.
- [20] S. Hoeche, et al., hep-ph/0602031.
- [21] D.J. Lange, Nucl. Instrum. Methods Phys. Res. A 462 (2001) 152.
- [22] R. Brun, F. Carminati, CERN Program Library Long Writeup W5013 (1993).
- [23] E. Boos, et al., CompHEP Collaboration, Nucl. Instrum. Methods Phys. Res. A 534 (2004) 250.
- [24] V.M. Abazov, et al., DØ Collaboration, Phys. Rev. Lett. 94 (2005) 151801.
- [25] V.M. Abazov, et al., DØ Collaboration, Phys. Rev. D 76 (2007) 092007.
- [26] V.M. Abazov, et al., DØ Collaboration, Phys. Rev. Lett. 98 (2007) 181802.
- [27] M. Artuso, et al., CLEO Collaboration, Phys. Rev. D 70 (2004) 112001.
- [28] W.-M. Yao, et al., Particle Data Group, J. Phys. G 33 (2006) 1.
- [29] V.M. Abazov, et al., DØ Collaboration, Phys. Rev. Lett. 98 (2007) 181902.
- [30] T. Andeen, et al., FERMLAB-TM-2365 (2007).