

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Faculty Publications from the Harold W. Manter  
Laboratory of Parasitology

Parasitology, Harold W. Manter Laboratory of

---

1-2001

# Phylogenetic Analysis of the Rhabdocoela (Platyhelminthes) with emphasis on the Neodermata and Relatives

David Zamparo  
*University of Toronto*

Daniel R. Brooks  
*University of Toronto, dnlbrooks@gmail.com*

Deborah A. McLennan  
*University of Toronto*

Eric P. Hoberg  
*United States Department of Agriculture*

Follow this and additional works at: <https://digitalcommons.unl.edu/parasitologyfacpubs>



Part of the [Parasitology Commons](#)

---

Zamparo, David; Brooks, Daniel R.; McLennan, Deborah A.; and Hoberg, Eric P., "Phylogenetic Analysis of the Rhabdocoela (Platyhelminthes) with emphasis on the Neodermata and Relatives" (2001). *Faculty Publications from the Harold W. Manter Laboratory of Parasitology*. 251.

<https://digitalcommons.unl.edu/parasitologyfacpubs/251>

This Article is brought to you for free and open access by the Parasitology, Harold W. Manter Laboratory of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications from the Harold W. Manter Laboratory of Parasitology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

# Phylogenetic analysis of the Rhabdozoa (Platyhelminthes) with emphasis on the Neodermata and relatives

DAVID ZAMPARO, DANIEL R. BROOKS, ERIC P. HOBERG & DEBORAH A. MCLENNAN

---

Accepted: 1 September 2000

Zamparo, D., Brooks, D. R., Hoberg, E. P. & McLennan, D. A. (2001) Phylogenetic analysis of the Rhabdozoa (Platyhelminthes) with emphasis on the Neodermata and relatives. — *Zoologica Scripta*, 30, 59–77.

Phylogenetic systematic analysis of 24 taxa representing the rhabdozoan platyhelminths, based on a suite of 89 morphological characters, produced two equally parsimonious trees, 181 steps long, with a consistency index (CI) of 0.69 and a rescaled consistency index (RCI) of 0.56, differing only with respect to that portion of the tree containing Umagillidae, Acholadidae, Graffillinae, Pseudograffillinae, Pterastericolidae and Hypoblepharinidae. Our results accommodate all previously proposed sister taxa to the Neodermata in a single clade in which ((Dalyelliidae + Temnocephalida) Typhloplanidae) is the sister group of ((Fecampiidae + *Urastoma*) (*Udonella* ((Aspidogastrea + Digenea) (Monogenea (Gyrocotylidea (Amphilinidea + Eucestoda)))))). Bootstrap and jackknife analyses indicate that the groupings of ((Dalyelliidae + Temnocephalida) Typhloplanidae) and of ((Fecampiidae + *Urastoma*) (*Udonella* ((Aspidogastrea + Digenea) (Monogenea (Gyrocotylidea (Amphilinidea + Eucestoda)))))) are highly robust, with the latter clade having a CI of 90% and RCI of 82%. Disagreements among previous analyses of these taxa have been due to the influence of missing data for critical characters in key taxa and differences in the taxa analysed, rather than any inherent weakness in the morphological data. Non-phylogenetic systematic approaches to homology assessment and misconceptions regarding phylogenetic systematic methodology are discussed. Recent analyses combining sequence data with a subset of approximately 60% of the morphological characters should be re-assessed using the entire morphological database. Even if *Udonella* is a monogenean, it is most parsimonious to suggest that the common ancestor of the Neodermata had a vertebrate–arthropod two-host life cycle.

David Zamparo, Daniel R. Brooks, Deborah A. McLennan, Centre for Comparative Biology and Biodiversity, Department of Zoology, University of Toronto, Toronto, Ontario M5S 3G5, Canada. E-mail: zamparo@zoo.utoronto.ca

Eric P. Hoberg, Animal Biosystematics Laboratory, ARS USDA, BARC East 1180, Beltsville, Maryland 20705, U.S.A.

## Introduction

The phylogenetic relationships among members of the phylum Platyhelminthes have received extensive scrutiny for nearly 20 years. Ehlers (1984) published the first phylogenetic systematic treatment for the phylum at about the same time as parasitologists were turning their attention towards intensive phylogenetic analysis of the parasitic groups within that phylum (the Neodermata and relatives: Brooks 1982, 1989a,b; Brooks *et al.* 1985a,b). Although most of the studies since those initial attempts have produced remarkably congruent results, there have been some disagreements, especially about the identity of the sister group to the Neodermata (Brooks 1982, 1989a,b; Ehlers 1984, 1985a,b, 1986; Brooks *et al.* 1985a; Rohde 1990, 1991;

Rohde *et al.* 1990; Brooks & McLennan 1993c; Jondelius & Tholleson 1993; Williams 1993; Watson 1997).

Congruence notwithstanding, some parasite taxonomists (e.g. Rohde 1990, 1994a, 1996) have objected to the hypothesized relationships among the parasitic groups, the choice of characters and the evolutionary implications of the phylogenetic systematic analyses, which call into question a number of long-standing myths about parasite evolution (Brooks & McLennan 1993a,b,c). More recently, the debate has shifted to assertions that molecular data are inherently superior to morphological data as markers of phylogeny (e.g. Justine 1998b; Littlewood *et al.* 1999a,b; Litvaitis & Rohde 1999). Recent molecular studies, for example, have either ignored (e.g. Baverstock *et al.* 1991; Blair 1993; Litvaitis & Rohde

1999) or minimized (Rohde *et al.* 1995; Littlewood *et al.* 1999a,b) the extensive morphological database that has been collected for the parasitic platyhelminths over the last 200 years. The assertion that morphological data are not as reliable as molecular data is a curious one, given that: (1) morphological studies routinely produce fewer equally parsimonious trees with better goodness of fit values than their wholly molecular counterparts; and (2) molecular studies have often produced results virtually identical to those already published by morphologists (e.g. Hoberg *et al.* 1997, in press; Mariaux 1998). This same debate has been carried out by systematists working on many different taxa. The result of the debate has been widespread agreement that the goal of systematics should be the production of phylogenetic hypotheses based on the most parsimonious (i.e. most scientifically robust) arrangement of all available evidence (see Kluge 1989, 1997, 1998a,b, 1999).

Jondelius & Tholleson (1993) provided the first direct phylogenetic systematic link between intense analysis of the parasitic groups and extensive analysis of the Platyhelminthes as a whole with their pioneering analysis of the Rhabdozoa. In this study, we present an updated analysis of the Rhabdozoa incorporating new character information that has been collected since the study by Jondelius & Tholleson (1993). The emphasis of this study is on the Neodermata and their closest relatives. We are particularly interested in answering two questions. What is the sister group of the Neodermata? Do the new data support or refute previous hypotheses of phylogenetic relationships within the Neodermata? In doing so, we will discuss the rationale for *a priori* exclusion of many morphological characters from recent phylogenetic analyses of the taxa herein considered. In this regard, we hope to show that the database of suitable morphological characters is far larger than that used in recent 'total evidence' studies.

## Materials and methods

### Taxa

The following taxa were included in this study (see also Jondelius & Tholleson 1993): Umagillidae, Pseudograffillinae, Graffillinae, Acholadidae, Pterastericolidae, Fecampiidae, Hypoblepharidae, Dalyelliidae, Provorticidae, Temnocephalida, Kytorrhynchidae, Promesostomidae, Solenopharyngidae, Trigonostomidae, Typhloplanidae, Kalyptorhynchia, *Urastoma*, *Udonella*, Aspidogastrea, Digenea, Monogenea, Gyrocotylidae, Amphilinidea and the Eucestoda.

### Character list

Characters were recorded based upon extensive descriptions in the literature: Aken'ova & Lester (1996); Bandoni & Brooks (1987a,b); Boeger & Kritsky (1993, 1997); Brooks (1982,

1989a,b); Brooks & McLennan (1993a,b,c); Brooks *et al.* (1985a,b, 1989, 1991); Bullock (1965); Cannon (1982, 1987); Ching & Leighton (1993); Christensen (1976); Christensen & Kannevorff (1965); DeClerk & Schockaert (1995); Ehlers (1984, 1985a,b, 1986, 1995); Ehlers & Sopott-Ehlers (1993); Fleming (1986); Fleming *et al.* (1981); Hoberg *et al.* (1997, in press); Hyman (1951); Ivanov (1952); Joffe & Kornakova (1998); Jondelius (1991, 1992); Jondelius & Tholleson (1993); Justine (1990, 1991, 1993, 1995, 1998a); Kannevorff & Christensen (1966); Kornakova & Joffe (1999); Koie & Bresciani (1973); Lee (1972); Littlewood *et al.* (1998, 1999a); Noury-Srairi *et al.* (1989a,b); Rohde (1986a,b, 1987, 1989, 1990, 1991, 1994b, 1998); Rohde & Watson (1993); Rohde *et al.* (1987a,b, 1989a,b, 1992, 1995, 1999); Shinn & Christensen (1985); Sopott-Ehlers (1991, 1996, 1998, 2000); Sopott-Ehlers & Ehlers (1995, 1997, 1998); Watson (1997, 1998a,b); Watson & Jondelius (1995); Watson & L'Hardy (1995); Watson & Rohde (1994a,b, 1995a,b,c); Watson & Schockaert (1996, 1997); Watson *et al.* (1992, 1995); Williams (1993); Wirth (1984); Xylander (1986, 1987a,b,c,d, 1988a,b, 1989, 1990). Characters were polarized using information on platyhelminth groups other than the Rhabdozoa summarized primarily in Ehlers (1984, 1985a,b, 1986, 1995), Jondelius & Tholleson (1993) and Littlewood *et al.* (1998, 1999a). '?' indicates that the state of the character is unknown in a particular taxon. Higher taxa that are polymorphic for a character were coded with the plesiomorphic state, as per Jondelius & Tholleson (1993) and standard phylogenetic systematic practice (Wiley 1981; Brooks & McLennan 1991; Wiley *et al.* 1991, in press; McLennan & Brooks, in press). Table 1 is the data matrix.

### Spermatozoal ultrastructure

- 1 Number of sperm axonemes. Two (0); none (1).
- 2 Axonemes. Free (0); incorporated into sperm cell body by proximo-distal fusion (1); incorporated into sperm cell body by distal proximal fusion (2).
- 3 Dense bodies. Present (0); absent (1).
- 4 Reverting migration which leads to the nucleus occupying a more distal position relative to the basal bodies. Absent (0); present (1).
- 5 Reverting migration includes a backward movement of the basal bodies and their axonemes to a proximal position. Absent (0); present (1).
- 6 Basal bodies retain their proximal position. Absent (0); present (1).
- 7 Electron-dense granules. Absent (0); present (1).
- 8 Spermatogenesis. Mature spermatozoa lacking dense heel, rotation of flagella and spur (0); mature spermatozoa possessing dense heel, rotation of flagella and spur (1).
- 9 Intercentriolar body during spermatogenesis. Present, weakly developed (0); absent (1).

**Table 1** Data matrix for phylogenetic analysis of the rhabdoceols. In this study, 91 morphological transformation series were considered. The most robust and inclusive results are based on 89 transformation series (17, 28 excluded) and with characters 16, 22, 24, 41, 60, 61, 78, 79 and 82 ordered. For identities of characters and states, refer to text. 0, plesiomorphic state; 1, 2, 3, 4, 5, apomorphic states; ?, unknown; OG, outgroup function (composite outgroup based on character argumentations for each transformation series).

Taxa	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
Outgroup	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Umagillidae	1	0	0	0	0	0	0	0	0	0	0	?	0	1	1	?	?	0	0	0	0	?	0	0	0	1	1	1	2	0	
Pseudograffillinae	?	?	?	0	0	0	?	0	?	0	0	?	0	?	?	?	?	0	0	0	0	?	0	?	0	1	1	0	0	0	
Graffillinae	1	0	0	0	0	0	1	0	0	0	0	?	0	?	?	?	?	0	0	0	0	0	0	0	0	1	1	?	?	0	
Acholadidae	?	?	?	0	0	0	?	?	?	0	0	?	0	?	?	?	?	0	0	0	0	0	0	0	0	1	1	1	2	0	
Pterastericolidae	1	0	0	0	0	0	0	0	0	0	0	2	0	1	1	?	?	0	0	0	0	0	0	0	0	1	1	1	2	0	
Hypoblepharinidae	?	?	?	0	0	0	?	?	?	0	0	?	0	0	?	?	?	0	0	0	0	0	0	0	0	1	0	1	1	2	0
Provorticidae	1	?	0	0	0	0	0	0	0	0	0	?	0	1	?	?	?	0	0	0	0	0	0	0	1	0	1	1	?	1	0
Kytorhynchidae	?	?	?	0	0	0	?	?	?	0	0	?	0	?	?	?	?	0	0	0	0	0	0	0	1	0	0	?	?	?	0
Promesostomidae	1	0	?	0	0	0	0	0	?	0	0	?	0	0	?	?	?	0	0	0	0	0	0	0	0	0	?	?	?	0	
Solenopharyngidae	1	0	0	0	0	0	1	0	?	0	0	?	0	0	?	?	?	0	0	0	0	0	0	0	0	1	?	?	?	0	
Trigonostomidae	1	2	1	0	0	0	?	0	?	0	0	2	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
Kalyptorhynchia	1	2	0	0	0	0	0	0	0	0	0	2	0	1	1	?	?	0	0	0	0	0	0	0	1	0	0	0	?	?	1
Dalyelliidae	1	2	0	0	0	0	1	1	0	1	0	2	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	?	?	1	0
Temnocephalida	1	0	0	0	0	0	1	1	0	1	0	2	0	1	1	0	0	0	1	0	0	0	0	1	0	0	0	?	?	0	0
Typhloplanidae	1	0	0	0	0	0	0	1	0	1	0	2	0	1	1	0	?	0	0	0	0	0	0	0	0	0	?	0	?	1	0
<i>Urastoma</i>	1	1	1	1	0	1	0	0	1	0	0	1	1	0	0	?	1	0	0	0	0	0	0	?	0	?	?	0	1	1	
Fecampiidae	1	1	1	1	1	0	0	0	1	0	0	1	1	0	0	0	1	0	0	0	0	0	0	0	1	0	0	1	0	0	1
<i>Udonella</i>	1	1	1	1	0	1	0	0	1	0	0	1	0	0	0	0	0	1	1	0	0	0	0	0	1	2	1	1	1	2	0
Aspidogastrea	1	1	1	1	0	1	0	0	1	0	0	1	0	0	0	1	1	0	0	0	0	0	0	0	1	2	1	1	0	1	0
Digenea	1	1	1	1	0	1	0	0	1	0	0	1	0	0	0	1	1	0	0	0	0	1	0	1	2	1	1	1	0	1	0
Monogenea	1	1	1	1	0	1	0	0	1	0	0	1	0	0	0	1	1	0	0	0	0	1	0	1	2	1	1	2	1	0	0
Gyrocotylidea	1	1	1	1	0	1	0	0	1	0	0	1	0	0	0	2	0	0	0	0	1	0	1	1	2	2	1	1	1	2	0
Amphiliinidea	1	1	1	1	0	1	0	0	1	0	0	1	0	0	0	2	0	0	0	1	0	2	1	2	2	1	1	1	2	0	0
Eucestoda	1	1	1	1	0	1	0	0	1	0	1	1	0	0	0	2	0	0	0	1	1	2	1	2	2	1	1	1	2	0	0

Taxa	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
Outgroup	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Umagillidae	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	1
Pseudograffillinae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0
Graffillinae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1	0	0
Acholadidae	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0
Pterastericolidae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0
Hypoblepharinidae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	1	0	0	0
Provorticidae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Kytorhynchidae	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Promesostomidae	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1
Solenopharyngidae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0
Trigonostomidae	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1
Kalyptorhynchia	0	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	?	1	0	0	0	1
Dalyelliidae	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	1
Temnocephalida	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	1
Typhloplanidae	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1
<i>Urastoma</i>	1	?	?	0	0	?	0	0	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	
Fecampiidae	1	3	?	0	0	1	0	0	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	2	1	0	1	1	
<i>Udonella</i>	0	2	1	1	1	1	1	0	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	4	0	2	1	1	1	
Aspidogastrea	0	2	1	1	1	1	1	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	2	0	0	2	
Digenea	0	2	1	1	1	1	1	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	2	0	0	2	
Monogenea	0	2	1	1	1	1	1	0	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	1	1	2	0	0	3	
Gyrocotylidea	0	2	1	1	1	1	1	0	0	2	1	0	1	1	1	1	1	1	1	1	1	1	0	0	2	1	2	0	2	4
Amphiliinidea	0	2	1	1	1	1	1	0	0	2	1	0	1	1	1	1	1	0	0	1	1	1	0	2	1	3	0	2	4	
Eucestoda	0	2	1	1	1	2	1	1	0	2	1	0	1	1	1	1	0	0	0	1	1	1	1	1	1	0	0	0	2	4

Table 1 Continued

Taxa	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91
Outgroup	0	0	0	0	0	0	0	0	0	0	0	0	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Umagillidae	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Pseudograffillinae	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Graffillinae	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Acholadidae	0	0	0	0	0	0	0	0	0	0	1	1	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Pterastericolidae	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hypoblepharinidae	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Provorticidae	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Kytorhynchidae	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Promesostomidae	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Solenopharyngidae	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Trigonostomidae	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Kalyptorhynchia	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Dalyelliidae	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Temnocephalida	1	0	0	0	1	1	0	0	0	0	0	0	1	0	0	0	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0
Typhloplanidae	1	0	0	0	1	0	0	0	0	0	0	0	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Urastoma</i>	1	0	1	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Fecampiidae	1	1	2	1	2	0	0	0	0	0	1	1	3	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Udonella</i>	1	1	1	1	1	1	0	0	0	0	0	0	1	0	1	0	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0
Aspidogastrea	1	1	1	1	1	0	0	1	0	0	0	0	1	0	1	1	2	1	1	2	1	0	0	0	0	0	0	0	0	0	0
Digenea	1	1	1	1	1	0	0	1	0	0	0	0	1	1	1	1	2	1	1	2	0	0	0	0	1	1	1	0	0	0	0
Monogenea	1	1	1	2	1	0	0	2	0	0	0	0	1	0	1	1	1	2	1	1	0	1	0	0	0	0	0	0	0	0	0
Gyrocotylidea	2	1	1	1	0	0	1	2	0	0	0	0	1	0	2	0	1	2	2	1	0	2	1	1	0	0	0	0	0	0	0
Amphiliinidea	2	1	1	1	0	0	0	2	1	1	0	0	1	0	2	0	1	2	2	1	0	3	1	0	0	0	0	0	0	0	0
Eucestoda	2	1	1	1	0	0	0	2	0	0	0	0	1	0	2	0	1	2	2	1	0	4	1	0	0	0	0	1	1	1	1

**10** Peripheral layer of microtubules in spermatozoa. Not spirally arranged (0); spirally arranged (1).

**11** Mitochondria in sperm. Present (0); absent (1).

#### *Protonephridia ultrastructure*

**12** Longitudinal ribs (rods). Absent (0); present, in two rows, inner formed by terminal cell, outer formed by canal cell (1); present, in single row of longitudinal ribs formed by canal cell (2).

**13** Interdigitating processes of weir. Absent (0); present (1).

**14** Terminal perikaryon. Present (0); absent (not close to flame) (1).

**15** Support structure of ribs (rods). Microtubules absent (0); microtubules present (1).

**16** Pair of cytoplasmic cords from canal cell connected by a desmosome. Absent (0); present (1).

**17** Surface of capillary. 'Saccate'/simple (0); lamellae of connected spaces (1); microvilli (2).

#### *Osmoregulatory system microstructure*

**18** Secondary protonephridial system of canals and pores. Absent (0); present (1).

**19** Giant paranephrocytes. Absent (0); present (1).

**20** Osmoregulatory system. Never reticulate (0); becomes reticulate in late ontogeny (1).

**21** Osmoregulatory system in early ontogeny. Not reticulate (0); reticulate (1).

**22** Protonephridia in larvae. In anterior end of body (0); in anterior and posterior end of body (1); in posterior end of body (2).

**23** Desmosomes in the passage of the first excretory canal cell. Present (0); absent (1).

#### *Tegument*

**24** Tegument. Cellular (0); syncytial, protruding to surface between epidermal cells (1); syncytial, not protruding to surface between epidermal cells (2).

**25** Adult body ciliation. Completely ciliated (0); at least some body ciliation lost (1); all ciliation lost (2). Some umagillids have lost body ciliation (Jondelius 1991); we consider this to be a derived trait within the group and consider the family to be plesiomorphically ciliated.

**26** Rhabdites. Present (0); absent (1).

**27** Duo gland organ. Present (0); absent (1).

**28** Rhabdomeric eyes. Two (0); none (1); four (2).

**29** Lensing. Non-mitochondrial (0); mitochondrial (1); no lenses (2).

**30** Rhabdoids (large granular and vesicular bodies in epidermis). Absent (0); present (1).

**31** Spur projecting from the basal body opposite the

horizontal rootlet of epidermal cilia. Absent (0); present (1).

32 Pharyngeal musculature. Circular muscle innermost (0); longitudinal muscle innermost; (1) circular muscle layer only (2); pharynx absent (3).

33 Dictyosomes and endoplasmic reticulum in larval/juvenile epidermis. Present (0); absent (1).

34 Larval epidermis. Not shed at end of larval stage (0); shed at end of larval stage (1).

35 Cilia of larval epidermis. With more than one rostrally directed rootlet (0); with one rostrally directed rootlet (1).

36 Specialized microvilli and microtubules in epithelium. Absent (0); present (1); modified into microtriches (2).

37 Epithelial sensory cells. Electron-dense collars absent (0); electron-dense collars present (1).

38 Post-larval epidermis. Not syncytial (0); syncytial [neodermis] (1).

39 Excretory vesicles. Lateral, paired (0); single, medial opening postero-dorsally (1).

40 Cephalic tentacles. Absent (0); present (1).

41 Vitelloglands. Absent (0); present, lining not syncytial (1); present, lining syncytial (2).

42 Anterior and posterior nervous system commissures. Single bilobed units (0); two bilobed units (1).

43 Ciliary bands on embryo. Absent (0); present, in three rows (1).

44 Larval epidermis. Not syncytial (0); syncytial (1).

45 Endoderm. Present in embryos (0); absent in embryos (1).

46 Vitellogenic cells. With more than one kind of electron-dense vesiculated inclusions (0); with one kind of electron-dense vesiculated inclusion (1).

47 Inner longitudinal muscle layer. Poorly developed (0); well developed (1).

48 Antero-lateral notch. Absent (0); present (1).

49 Nuclei in larval epidermis. Present (0); absent (1).

50 Multiciliary nervous receptors. Present (0); absent (1).

51 Epithelial lining of genital ducts. Not syncytial (0); syncytial (1).

52 Protonephridial ductules. Ciliated (0); not ciliated (1).

53 Medullary and cortical distinction. Not apparent (0); apparent (1).

54 Protein embeddings in larval epidermis. Absent (0); present (1).

### **Reproductive system**

55 Male intromittent organ. Simple stylet (0); cirrus [sometimes mistakenly called a penis] (1); copulatory papilla (2); complex stylet (3); absent (4). Monogeneans do not have a copulatory stylet (the accessory piece in some monogeneans is an independently evolved structure, and a cirrus is plesiomorphic for the group: Boeger & Kritsky 1993,

1997). The copulatory papillae of Gyrocotylidea and Amphilinidea may be vestigial/reduced cirri.

56 Openings of male and female gonopores. Common genital atrium (0); separate (1); separate sexes (2).

57 Position of genital atrium or genital pores. Posterior (0); caudal (1); anterior (2); lateral (3).

58 Muscular copulatory bulb. Present (0); absent (1).

59 Testes. Paired (0); single (1); multiple, in two lateral bands (2). A single testis occurs convergently within Aspidogastrea, Digenea and Monogenea, but phylogenetic analyses (Brooks *et al.* 1985b, 1989; Boeger & Kritsky 1993, 1997) have shown that paired testes are plesiomorphic in each case.


60 Female reproductive system. Simple oviduct (0); oviduct expanded to form antrum (functional uterus) without separate opening (1); oviduct coiled, with small secondary tube (Laurer's canal) opening to the surface (not opening to surface or absent in derived taxa), used to vent excess material from oviduct (2); oviduct relatively straight, with secondary tube forming separate tubular uterus with uterine pore opening to surface (3); oviduct relatively straight, uterus highly coiled (4). Previous phylogenetic analyses of the Cercomeria (Brooks *et al.* 1985a) and Rhabdozoa (Jondelius & Tholleson 1993) have treated various portions of the female reproductive system as a series of separate characters. These include the presence or absence of a vagina, presence or absence of a uterus, and their position(s) relative to the male gonopore and to the body in general. This has been complicated in part by the fact that most neodermatans possess two (or even three) openings of the female reproductive tract.

The majority of cercomerideans (Trematoda + Cercomeromorphae) exhibit a bifurcated oviduct, with each bifurcation forming a tube that opens to the exterior. These tubes have been functionally defined in the parasitic taxa, i.e. any egg-containing tube is called the uterus, and the alternative tube is called the vagina. Thus, in the trematodes, the male gonopore and uterine pore are said to be proximate, with the vagina separate. The vagina (called the Laurer's canal) is almost always short, narrow and relatively straight (in many cases it does not open to the exterior or is even lost) and the uterus is generally coiled. In the Monogenea, all three pores are separate plesiomorphically, with the apomorphic state 'uterine and male pores proximate' being displayed by some taxa. The uterus and vagina are relatively well developed, short and straight. Doubling the vagina (considered by Brooks *et al.* 1985a to be an autapomorphy for the Monogenea) appears to be an apomorphic trait within the Monogenea (Boeger & Kritsky 1993, 1997). In the Gyrocotylidea, all three pores are proximal and separate (the plesiomorphic condition for the Monogenea). In the Cestodea (Amphilinidea + Eucestoda), the male pore and the vaginal pore

are proximal, with the uterine pore distantly situated. Finally, within the Cestodaria (Gyrocotylidae + Amphilinidea + Eucestoda), the uterus is plesiomorphically highly coiled (it is apomorphically saccate in the Eucestoda: Brooks *et al.* 1991; Hoberg *et al.* 1997, in press). Such coiling also occurs convergently within the Monogenea (Boeger & Kritsky 1993, 1997). Establishing homologies for these structures across taxa has been difficult, demanding complex evolutionary scenarios to explain the diversity of ducts, tubes, pores and their positions relative to each other. We suggest that these scenarios have been unnecessarily complex and propose the following alternative.

The basic unit of the platyhelminth female reproductive system is an ovary (paired plesiomorphically) connected to a tubular oviduct, a canal which originates from the ovary and terminates in a genital pore that communicates with the external environment. Plesiomorphically, this canal functions as both vagina (receiving sperm) and uterus (delivering eggs to the external environment) and is situated near the male genital pore, either sharing a common atrium with the male pore or not (Fig. 1). Within the rhabdozoans, including fecampiids, *Urastoma* and *Udonella*, the oviduct is expanded, producing a functional uterus, or antrum. The antrum may be symmetrical or asymmetrical, small, containing a single egg, or large, containing several eggs, and saccate or somewhat tubular.

We propose that, regardless of perceived function, the oviduct is the portion of the female reproductive system plesiomorphically proximal to the male genital pore, with which it may or may not share a common gonopore (genital atrium). The secondary duct may be proximal to (Monogenea, Gyrocotylidae) or distant from the openings of the oviduct and male genital pore (dorsal in the trematodes, ventral in the Amphilinidea and Eucestoda). The Laurer's canal is thus actually homologous with the uterus, not the vagina, of the Cercomeromorphae. The current function of the Laurer's canal, expulsion or digestion of sperm and other debris from the fertilization and egg-making process (e.g. Juel's organ in some hemiuriform digeneans), may well have been the original function of the duct. The widespread belief that the Laurer's canal is a vestigial vagina stems from discussions of the presumed degenerate evolutionary nature



**Fig. 1** Schematic representation of diversity in the female reproductive system of neodermatans. 0, 1, 2, 3, 4 refer to the character states used in this analysis. State 0 is the condition found among various rhabdozoans. State 1 occurs in *Urastoma*, Fecampiidae, *Udonella* and various rhabdozoans. State 2 is the condition found in trematodes. State 3 is the condition among the monogeneans. State 4 is the condition of the Cestodaria. A, antrum; L, Laurer's canal; M, metraterm; OD, oviduct; OV, ovary; S, sphincter; U, uterus.

of parasites beginning in the late 19th century. Actual evidence is rare. For example, without sectioning his material, Cohn (1902) stated that he had found one specimen of *Liolope copulans* extruding its cirrus into the Laurer's canal of another. Brooks & Overstreet (1978), however, noted that they never found any evidence of this behaviour in a close relative of *L. copulans*, *Dracovermis occidentalis* Brooks & Overstreet, 1978. They stated that '... based on the narrow Laurer's canal, wide cirrus, thick and large genital atrium, and uterus occasionally entirely packed with sperm in *Dracovermis occidentalis*, we doubt that Laurer's canal in that species serves for more than the elimination of excess products.' Increased egg-holding capacity in the trematodes is made possible by extensive coiling of the oviduct, while in the cercomeromorphs it is due to the elongation (Monogenea) and coiling (gyrocoylids, amphilinids and eucestodes) of the Laurer's canal, co-opted (an *exaptation*: Gould & Vrba 1982) as a functional uterus distinct from the oviduct.

The above proposal provides a succinct conception of the evolution of the number, nature and position of the ducts and pores of the female reproductive system in the Cercomeridea. Interestingly, it is also the scheme proposed by Looss (1893) but apparently forgotten until now. Our coding of characters associated with these structures reflects this new hypothesis. Finally, many trematodes have been described as exhibiting a glandular muscle surrounding the terminal end of the uterus called the 'metatherm' (Smyth 1994) or 'metraterm' (Noble *et al.* 1989). Many eucestodes have been described as having a muscular structure at the terminal end of the vagina called a 'vaginal sphincter'. If our hypothesis above is true, it is likely that these structures are homologous. At present, we lack sufficient information to use this as a character.

- 61 Ovary. Paired (0); single and spherical (1); single and bilobed (2).  
 62 Mehlis' gland. Absent (0); present (1).  
 63 Vitellaria. Paired, compact, medial (0); lateral and follicular (1); compact and medial vitellarium (2). Compact vitellaria occur convergently in a number of digenean and eucestode groups, but are apomorphic within these taxa (Brooks *et al.* 1985b, 1989, 1991; Hoberg *et al.* 1997, in press).  
 64 Cirrus. Absent (0); present, muscular and aspinose (1); present, muscular and spinose (2).  
 65 Testes. Preovarian (0); postovarian (1); dioecious (2). Dioecy appears convergently in some digenean (e.g. Schistosomatidae) and some eucestode (e.g. *Dioecotaenia*, *Dioecocestus*, *Shipleya*, *Gyrocoelia*) groups (Brooks *et al.* 1985b, 1989, 1991; Hoberg *et al.* 1997, in press). Because the Fecampiidae are dioecious, the character is inappropriate. We have the option of either coding the fecampiids as '9' — inappropriate, or as '2', as the condition is autapomorphic.

The choice of coding in this instance does not affect the analysis.

- 66 Eggs. Round adhesive disc at the end of filament where the substance of the disc is secreted later when the worm attaches the egg to the body of the host. Absent (0); present (1).  
 67 Vitellaria. Not encircling entire body (0); encircling entire body, extending along entire body length (1). The apomorphic state appears convergently in some eucestode groups (Hoberg *et al.* 1997, in press).  
 68 Permanent uterine pore. Absent (0); present, dorsal (1); present, ventral (2).  
 69 Uterine pore. Not proximal to pharynx (0); proximal to pharynx (1).  
 70 Uterus. Coiled, not 'N'-shaped (0); 'N'-shaped (1).

#### Digestive system

- 71 Mouth and pharynx. Present (0); absent (1). The apharyngeate condition exhibited by some monogeneans and digeneans is convergently evolved within these groups (Brooks *et al.* 1985b, 1989; Boeger & Kritsky 1993, 1997).  
 72 Doliiform pharynx (pharynx bulbosus of Jondelius & Tholleson 1993). Present (0); absent (1).  
 73 Pharynx placement. In anterior half of worm (1); medial to posterior half of worm (2); absent (3). This is a difficult character to polarize because most outgroups are polymorphic. Jondelius & Tholleson (1993) proposed that anterior was plesiomorphic for the rhabdocoels, but their own argument can also be used to support the contention that a pharynx in the mid to posterior half of the body is plesiomorphic; therefore, we have coded the outgroup state as '?' and given each ingroup state a non-zero number.  
 74 Oral sucker. Lacking a capsule (0); with a capsule (1).  
 75 Gut shape. Saccate (0); bifurcate (1); lacking in adults (2). Convergent reversal to a saccate gut from a plesiomorphically bifurcate gut occurs within the aspidogastreaans, digeneans and monogeneans (Brooks *et al.* 1985b, 1989; Boeger & Kritsky 1993, 1997).  
 76 Oral sucker. Absent (0); present (1).

#### Posterior adhesive organs

- 77 Posterior adhesive organ. Absent (0); present, not delimited by capsule (1); present, delimited by capsule (2).  
 78 Posterior adhesive organ. Absent (0); present, no hooks (1); present, with hooks (2).  
 79 Posterior adhesive organ. Absent (0); present throughout life (1); present only during early development, partially invaginated (2).  
 80 Posterior adhesive organ. Absent (0); present, terminal (1); present, ventral (2).  
 81 Posterior sucker. Without transverse septa (0); hypertrophy and linear subdivision of posterior sucker by transverse septa (1).



82 Hooks on posterior end of larva. Absent (0); sixteen equal-sized hooks (1); ten equal-sized hooks (2); six large and four small hooks (3); six hooks (4).

83 Posterior body invagination. Absent (0); present (1).

84 Rosette at posterior end of body. Absent (0); present (1).

**Ontogeny**

85 Miracidium. Absent (0); present (1).

86 Sporocyst. Absent (0); present (1).

87 Cercaria. Absent (0); present (1).

88 Proceroid. Absent (0); present (1).


89 Plerocercoid. Absent (0); present (1).

90 Cerebral development in larvae. Present (0); absent (1).


91 Extra-embryonic membrane. Not formed by embryo (0); formed by embryo (1).

**Analyses performed**

Data were analysed using standard Hennigian argumentation (see Hennig 1966; Wiley 1981; Brooks & McLennan 1991; Wiley *et al.* 1991, in press), and results were confirmed using the 'branch and bound option' on the computer program PAUP 4\*, implemented on Macintosh G3/400, G4/450 and G4/500 computers. Acctran and Deltran character optimization produced the same results. Bootstrap and jackknife analyses



**Fig. 2** Majority rule consensus tree for 24 rhabdozoel taxa based on 98 most parsimonious trees (MPTs) (tree length (TL) = 190), consistency index (CI = 67%) and rescaled consistency index (RCI = 55%) produced by phylogenetic systematic analysis of 91 morphological characters.



**Fig. 3** Two most parsimonious trees (MPTs) (tree length (TL) = 181, consistency index (CI) = 69%) and rescaled consistency index (RCI = 56%) for 24 rhabdoceol taxa produced by phylogenetic systematic analysis of 89 morphological characters. Ordering multistate characters 16, 22, 24, 41, 60, 61, 78, 79 and 82 produced the same results.


were performed using 10 000 replicates, with the exception of the complete data set, for which only 100 replicates were performed due to computational constraints.

### Results

The analysis of all 91 characters, unordered, produces 98 most parsimonious trees (MPTs), each 190 steps long, with a consistency index (CI) of 67% and rescaled consistency index (RCI) of 55%. Forty-one of these MPTs place the Kytorrhynchidae, Promesostomidae, Trigonostomidae, Typhloplanidae, Dalyelliidae and Temnocephalida at the base of the tree, similar to results reported by Jondelius & Tholleson (1993) and Littlewood *et al.* (1999a,b). The remaining 57 MPTs suggest that these taxa are part of an inclusive clade also containing the Neodermata, a result more similar to the hypothesis proposed by Ehlers (1984, 1985a,b, 1986, 1995) and Brooks and coworkers (Brooks *et al.* 1985a; Brooks 1989a,b; Brooks & McLennan 1993a). Figure 2 is the 50% majority rule consensus tree for the 98 MPTs. We have discovered that this 'dichotomous' result

in the placement of the Kytorrhynchidae, Promesostomidae, Trigonostomidae, Typhloplanidae, Dalyelliidae and Temnocephalida is the product of missing data for key taxa in characters 17 and 28. In computer-assisted phylogenetic studies, some configurations of missing data can produce effects similar to long branch attraction effects in the analysis of nucleotide sequence data (see Nixon & Davis 1991; Platnick *et al.* 1991; Maddison 1993; Wilkinson 1995). Other characters show low character consistencies on the tree as well, but their inclusion does not affect the stability of the results. We believe that characters 17 and 28 are too poorly documented at present to be useful.

Removing characters 17 and 28 produces two MPTs (Fig. 3), 181 steps long, with a CI of 0.69 and an RCI of 0.56, differing only in the degree of resolution of that portion of the tree containing the Umagillidae, Acholadidae, Graffillinae, Pseudograffillinae, Pterastericolidae and Hypoblepharinidae. Characters 16, 22, 24, 41, 60, 61, 78, 79 and 82 are multi-state transformation series produced by combining what were previously considered to be a series of binary characters



**Fig. 4** Bootstrap and jackknife consensus tree for 24 rhabdoceol taxa based on 89 morphological characters, with multistate characters 16, 22, 24, 41, 60, 61, 78, 79 and 82 ordered. Bootstrap/jackknife values appear on appropriate branches.

(Brooks & McLennan 1993c). The relationships shown in Fig. 3 supported the ordering of these transformation series. Phylogenetic analysis with these nine characters ordered produced the same results as Fig. 3. Successive approximations with reweighting of the data produced the single tree shown in Fig. 3(A), which we consider to be the most robust hypothesis incorporating the maximum amount of information possible for this data set.


Six taxa in the present study, the Acholadidae, Pseudograffillinae, Hypoblepharinidae, Solenopharyngidae, Promesostomidae and Kytorrhynchidae, have substantial missing data entries, and the portion of the tree containing the Umagillidae, Pseudograffillinae, Graffillinae, Acholadidae, Pterastericolidae and Hypoblepharinidae produces the two MPTs shown in Fig. 3. Not surprisingly, then, bootstrap and jackknife analyses indicate that only the groupings of ((Dalyelliidae + Temnocephalida) Typhloplanidae) and ((Fecampiidae + *Urastoma*) (*Udonella* ((Aspidogastrea + Digenea) (Monogenea (Gyrocotylidea (Amphilinidea + Eucestoda)))))) are robust (Fig. 4). Given recent successes at finding many

morphological traits for other platyhelminth groups (e.g. Lundin 2000), we feel confident that sufficient characters are there to be discovered, and a fully robust assessment of the Rhabdoceola is feasible. In the rest of this study, we will concentrate on the Neodermata and their closest relatives.

The placement of the Temnocephalida in our analysis precludes the interpretation that all posterior holdfast organs in this clade are homologous. The taxon *Cercomeria* Brooks, 1982 therefore cannot be maintained, as suggested by Ehlers & Sopott-Ehlers (1993) and Rohde & Watson (1995). The clade of Fecampiidae + *Urastoma* as the sister group of the Neodermata supports the monophyly of the Revertospermata Kornakova & Joffe, 1999, but not the Mediofusata Kornakova & Joffe, 1999.

## Discussion

Discussions of the phylogeny of the Neodermata revolve around two questions: (1) What is the sister group of the Neodermata? (2) How does the choice of sister group affect



**Fig. 5** Bootstrap and jackknife consensus trees for the Revertospermata Kornakova & Joffe (Neodermata (Fecampiidae + *Urastoma*)) based on 89 morphological characters, with multistate characters 16, 22, 24, 41, 60, 61, 78, 79 and 82 ordered. Bootstrap/jackknife values appear on appropriate branches.

the hypotheses of the relationships among taxa within the Neodermata? With regard to the first question, four taxa have been previously suggested as sister groups of the Neodermata: (1) the Dalyelliidae and Typhloplanidae (Ehlers 1984, 1985a,b, 1986, 1995; Ehlers & Sopott-Ehlers 1993); (2) the Temnocephalida (Brooks 1982, 1989a,b; Brooks *et al.* 1985a; Brooks & McLennan 1993c); (3) *Urastoma* (Rohde *et al.* 1990; Williams 1993; Watson 1997; Kornakova & Joffe 1999); and (4) the Fecampiidae (Rohde 1990, 1991; Litvaitis & Rohde 1999). This study included all four candidates in the same analysis, and the results indicate that they comprise the four closest relatives of the Neodermata (Fig. 3). With respect to the second question, the present analysis supports the monophyly of the Monogenea and the placement of *Udonella* as the basal member of the Neodermata as originally proposed by Brooks *et al.* (1985a). Re-analysing the present data set using any number and combination of the four putative sister groups as outgroup taxa produces the same result. This occurs because the data for relation-

ships within the Neodermata are highly robust (CI = 94%, RCI = 87%), making any combination of the four candidates suitable outgroups. The portion of the tree comprising the (Fecampiidae + *Urastoma*) + Neodermata is slightly less robust (CI = 90%, RCI = 82%) because the Fecampiidae + *Urastoma* clade is not as well supported (see bootstrap and jackknife values in Fig. 5).

Brooks *et al.* (1985a) used a data set of 39 transformation series in their initial analysis of the Neodermata; this produced a single MPT, 41 steps long (CI = 95%), depicting the same relationships as shown in Figs 2–5. In that analysis, the authors used only attributes deemed informative by authors of numerous earlier studies in order to demonstrate that differences in results were due to differences in methods of analysis, not in choice of characters. Adding more morphological traits produced a data set of 127 binary characters (Brooks 1989a,b), corroborating the original phylogenetic hypothesis, producing a single MPT, 131 steps long (CI = 97%). Brooks & McLennan (1993c) produced

the same MPT, 161 steps long, for 153 apomorphic traits (CI = 95%). In the present study, some characters were modified according to new findings, some redundant characters listed by Brooks & McLennan (1993c) were combined, and 47 fewer autapomorphies were used, resulting in the same MPT, 107 steps long, for 100 apomorphic traits (CI = 93%); including the 47 autapomorphies produces a single MPT for the Neodermata, 154 steps long, for 147 apomorphic traits (CI = 95%).

Despite consistent robust support for this hypothesis during the past 15 years, some researchers have felt uncomfortable with the results (Rohde 1990, 1994a, 1996; Justine 1998b; Littlewood *et al.* 1998, 1999a,b). We believe that misunderstandings about phylogenetic systematics have been responsible for the differences of opinion. The most fundamental misunderstanding stems from the way in which phylogeneticists determine homologous character states. All systematists begin the search for homology by using a set of criteria, such as those proposed by Remane (1952), to determine whether two or more characters are 'similar' (see discussion in de Pinna 1991). These similarities apply to both identity (a finger is a finger) and also transformation (a bird's wing is a tetrapod forearm). Assessing similarity based upon such biological criteria, without recourse to knowledge of underlying genealogical relationships, eliminates any hint of circularity in the process (see Eldredge & Cracraft 1980; Wiley 1981; Brooks & McLennan 1991; Wiley *et al.* 1991, in press; McLennan & Brooks, in press). The difference among systematists begins with how these similarities are treated next. Phylogenetic systematists use assessments of similarity to construct *hypotheses* of homology 'If a and b look the same (e.g. are in the same position, develop from the same tissue), then they are homologous'. This is called Hennig's auxiliary principle (see Hennig 1966; Wiley 1981; Brooks & McLennan 1991; Wiley *et al.* 1991, in press; McLennan & Brooks, in press). These hypotheses are tested by using phylogenetic systematics and are ultimately corroborated or rejected. In the latter case, we conclude that the similarity is due to homoplasy.

Some taxonomists, on the other hand, believe that they can make *a priori* judgements about which similarities are due to homology, and which are due to homoplasy, and thus eliminate some characters (the putative homoplasies) from the data set before the analysis begins. Such *a priori* judgements are valid only if they are supported by evidence. For example, experimental research has demonstrated that characters such as the number of vertebrae or fin rays in stickleback fish are strongly influenced by the temperature under which the larvae develop (Lindsey 1962; Hagen 1967). Reporting the number of vertebrae or fin rays without adjusting for developmental temperature, an almost impossible feat in wild caught fish, thus introduces a known source

of homoplasy into the data set. In this case, systematists are justified in eliminating these traits from their analysis *a priori*. Because such data are rare, however, it becomes important to ask: 'what supports the elimination of a particular character, or type of character, from an analysis?'

With regard to the Neodermata, it has been asserted that complex characters are more likely to be homologous than simple characters (Rohde 1990, 1994a, 1996; Littlewood *et al.* 1999a). What evidence is there to support this assertion? There is a large body of evidence documenting simple genetic bases for many homologous behavioural and morphological characters in *Drosophila* species. That alone would seem to falsify the hypothesis that simple characters are not likely to be homologous. This assertion stems, in part, from a misunderstanding of levels of homology. The presence of bristles may be homologous across *Drosophila*, but the exact number of bristles may display some homoplasy. In other words, there is no evidence indicating that sweeping generalities can be made about the nature of homology vs. homoplasy based upon a vague notion of simple vs. complex character structure.

The hypothesis about the relative merits of simple vs. complex characters as markers of genealogical relationships could be examined by assigning a 'simple' vs. 'complex' status to characters *a priori*, running those characters through a phylogenetic systematic analysis, and then asking whether there is a significant difference in homoplasy among the two character classes. Once this process has been repeated for a substantial number of data sets from different groups of organisms, we could then begin to determine the validity of the hypothesis. In lieu of this evidence, we opted to use all available characters, presuming maximum homology and character independence *a priori*, and relying on phylogenetic congruence among all characters *a posteriori* as the final arbiter of homology (Wiley 1981; Kluge 1989, 1997, 1998a,b, 1999; de Pinna 1991).

While the primary function of phylogenetic analysis is to produce a robust hypothesis of phylogenetic relationships, it also provides a means to help us determine when our *a priori* presumptions are incorrect. Once we have a phylogenetic hypothesis based on as many characters as possible, we can move from homology presumptions to homology determinations. Hennig (1966) considered such 'reciprocal illumination', using the overall analysis to assess individual *a priori* presumptions of homology, to be a primary benefit of phylogenetic systematics. The homologies are the traits that are congruent with the phylogenetic tree, whether they are complex or simple in nature; homoplasies are those that are incongruent with the tree. For example, this study supports the proposal by Ehlers & Sopott-Ehlers (1993) and Rohde & Watson (1995) that the holdfast organ of the temnocephalids is not homologous with the holdfasts of neodermatans

(characters 77–84). Brooks *et al.* (1985a) hypothesized that the various holdfasts, while demonstrably different, were all part of a homologous transformation series. Within phylogenetic systematic methodology, this hypothesis could not be falsified by reiterating that the holdfasts were different (Rohde & Watson 1995), but could be falsified by including more taxa in the analysis, as we have done herein.

Additionally, Rohde and coworkers (Rohde 1990, 1994a, 1996; Littlewood *et al.* 1999a) suggested that protonephridial characters should be given high weight in phylogenetic analyses of the Platyhelminthes. We considered six protonephridial characters in this study. Three of them (12, 13, 16) have character consistencies of 100%, character 15 has a character consistency of 50%, 17 has a character consistency of 33% and character 14 has a character consistency of 25%. The combined character consistencies for these traits is 68%, and their exclusion from the analysis produces the same tree topology as shown in Fig. 3(A) and increases the CI slightly. In addition, character 17 is one of the characters producing marked instability due to missing data. Reciprocal illumination thus tells us that protonephridial characters are, at best, no better than any other character.

Phylogeneticists expect that the analysis of a data set comprising incorrect homology assessments will produce a distinctive result — many MPTs with low CIs. This is not the case with the Brooks *et al.* (1985a) (see also Brooks 1989a,b; Brooks & McLennan 1993c) data sets, nor is it the case with the present data set. In the current study, 90% of the characters support the relationships indicated for the Revertospermata, and these results strongly corroborate previous analyses. In the past, these results have been rejected because we are dealing with parasites (Neodermata) and symbiotic ‘turbellaria’, and adaptation to a common lifestyle is ‘known’ to produce high degrees of correlated homoplasy (Rohde 1990, 1994a, 1996; Littlewood *et al.* 1999a). To correct for this problem, characters ‘known’ to be adaptations to parasitism/symbiosis have been discounted (eliminated from the analysis *a priori*). For example, Rieger & Tyler (1985) suggested that similar structures in taxa sharing similar environments (e.g. exposed to similar selection pressures) should be coded *a priori* as homoplasious, or ambiguous as in Littlewood *et al.* (1999a,b).

Such suggestions ignore the basic Darwinian notion that homologies can be adaptations and that adaptation need not produce homoplasy. In the past decade, a substantial amount of evidence has accumulated indicating that most similarities in structure, function and preferred environment are due to common ancestry (Wanntorp *et al.* 1990; Brooks & McLennan 1991; Harvey & Pagel 1991). There is thus no reason to exclude, or manipulate, any ‘adaptive’ character from any analysis (McLennan *et al.* 1988; Brooks & McLennan 1991, 1993c, 1994; McLennan 1993). In addition,

Ronquist’s (1994) study on the evolution of inquilinism in cynipid hymenopterans, for example, showed that removal of characters associated with a parasitic lifestyle did not alter the phylogenetic assessment that inquilinism had arisen only a single time in the group. Finally, Trouvé *et al.* (1998) showed that a suite of life history traits for free-living and parasitic platyhelminths did not differ, suggesting that neodermatans do not have a ‘parasitic mode of life’ so much as a ‘platyhelminth mode of life’ in a parasitic context.

In recent years, some authors have disparaged the morphological data upon which previous analyses of the Neodermata and their relatives had been performed because the results were not compatible with molecular data (Rohde 1990, 1994a, 1996; Littlewood *et al.* 1999a; Litvatis & Rohde 1999; Mollaret *et al.* 2000). It has also been suggested that the phylogenies based upon morphological data have been highly variable and greatly differ among each other (Littlewood *et al.* 1999a). This has not actually been the case. First, the relationships among the neodermatan groups have been the same in multiple studies using phylogenetic systematic methods beginning in 1985, with CI values remaining between 95% and 97% despite an increase in the number of characters used from 39 to 147. Second, differences in hypotheses of the sister group of the Neodermata have been based on differences in the taxa analysed; our analysis herein accommodates all previously proposed sister groups in a manner that is congruent with all previous hypotheses.

In addition, Kornakova & Joffe (1999) pointed out that molecular results have failed to reproduce the monophyly of several firmly established taxa (based on morphology) and suggest that we consider sampling and long-branch attraction as serious effects in molecular analyses. For example, molecular studies suggest various combinations of para- or even polyphyly for the Monogenea, whereas morphological studies consistently suggest that the group is monophyletic. Some authors take this as an indication that we should question all morphological traits used in phylogenetic studies of monogeneans (Rohde 1990, 1994a, 1996; Justine 1998b; Litvatis & Rohde 1999; Mollaret *et al.* 2000). Littlewood *et al.* (1999b) showed that a combination of sequence data and only 50 of the 89 characters used herein supported a monophyletic Monogenea, and accepted that grouping. Since the molecular data alone did not support Monogenean monophyly, the study by Littlewood *et al.* (1999b) provides evidence of insufficiencies in the sequence data as suggested by Kornakova & Joffe (1999).

This thinking needs to be carried through consistently in all future total evidence studies. Littlewood *et al.* (1999a) coded nine characters shared uniquely by *Urastoma*, the Fecampiidae and the Neodermata as ambiguous for *Urastoma* and Fecampiidae, presumably based on Rohde’s (1994a: 1104) assertion that ‘comparison of DNA sequences ... suggests that

the [fecampiids are] not a close relative of the Neodermata ... thus the morphological similarities of the two groups appear indeed to be due to convergent evolution'. Likewise, Littlewood *et al.* (1999a,b) made a number of *ad hoc* assumptions concerning *Udonella*. For example, the absence of larval hooks was coded *a priori* as apomorphic secondary loss, when the same absence of larval hooks in aspidogastreae and digeneans was coded as plesiomorphic absence. These added assumptions clearly demonstrate an *a priori* coding 'preference' for regarding *Udonella* as a Monogenean. Finally, Littlewood *et al.* (1999a,b) utilized only slightly more than half of the available morphological characters that had been summarized in Brooks & McLennan (1993c). Many of these traits were characterized by Rohde (1990, 1994a, 1996) (also Litvaitis & Rohde 1999) as exhibiting a low probability of being homologous. Our study herein does not support that characterization. In fact, the total morphological database provides very strong support not only for the monophyly of the Monogenea, which Littlewood *et al.* (1999b) accepted, but also for the Fecampiids + *Urastoma* as the sister group of the Neodermata and *Udonella* as the sister group of the Cercomeridea Brooks, O'Grady & Glen 1985 (Trematoda + Cercomeromorphae).

Finally, our study corroborates the hypothesis that the ancestor of the Trematoda + Cercomeromorphae had a two-host life cycle involving the addition of a vertebrate host to the plesiomorphic arthropod host direct life cycle (Brooks *et al.* 1985a; Brooks 1989b; Brooks & McLennan 1993c), contrary to the proposal by Littlewood *et al.* (1999b) that the original life cycle was a single vertebrate host direct cycle. This is the most parsimonious explanation even if *Udonella* is a monogenean. It supports the notion that vertebrate endoparasitism in this group originated through the predation of vertebrates on arthropods. It may also be an example supporting the hypothesis that the alternation of hosts is an adaptive response to avoid the evolutionary costs of overspecialization (Moran 1988, 1994; see also Kuris & Norton 1985).

## Conclusions

The morphological database for the Neodermata and close relatives is highly robust. This is partly due to the fact that the data themselves are numerous and unambiguous. More importantly, scientific hypotheses are more robust in proportion to the number of tests they have survived (Popper 1960, 1968a,b, 1972, 1976, 1992), and the current database reflects the efforts of a number of specialists to refute the hypothesis first proposed by Brooks *et al.* (1985a). The current study also shows that phylogenetic systematic analysis is capable of uncovering instances in which our *a priori* presumption of homology is not supported. Thus, the selective removal of characters *a priori* is not necessary and

indeed is counterproductive if our aim is always to produce the most robust hypothesis of phylogenetic relationships possible given all available evidence. The parasitic platyhelminths represent one of the most extensively studied animal groups, with a database assembled over the past 200 years that will soon exceed 2500 morphological character states. This represents historical continuity in our studies of flatworms, which comprises a formidable assemblage of knowledge about structure and biology. The results of the current study indicate that comparative morphology remains viable, tractable and powerful. Phylogenetic analyses using morphological data provide an excellent framework for assessing our young but growing molecular database. We suggest that future total evidence studies should make full use of the large and robust morphological database documented herein.

This study also highlights two other benefits of a phylogenetic systematic approach: the ability, through reciprocal illumination, to falsify previous hypotheses of character evolution, and the ability to highlight areas where further research would be the most immediately beneficial. In this case, we clearly need more studies on the enigmatic groups such as the Acholadidae, Pseudograffillinae, Hypoblephariniidae, Solenopharyngidae, Promesostomidae, Trigonostomidae and Kytorrhynchidae.

## Acknowledgements

We express particular thanks to Dr Lesley Fleming, University of New Brunswick, and Dr Robin Overstreet, Gulf Coast Research Laboratory, for providing critical information about some key characters in some key taxa. We thank Dr Walter Boeger, Dr Gregory Klassen, Dr Delane Kritsky, Dr Virginia León-Régagnon, Dr Gerardo Pérez Ponce de León and Dr Thomas Platt for helpful discussions about platyhelminth phylogeny throughout the long gestation of this study. Finally, we also thank Dr Fredrik Ronquist and Dr Sören Nylin for conceptual discussions on the study of parasitic organisms. Funds for this study were provided by operating grants from the Natural Sciences and Research Council of Canada to DRB and DAM.

## References

- Aken'ova, T. O. & Lester, R. J. G. (1996). *Udonella myliobati* n. comb. (Platyhelminthes: Udonellidae) and its occurrence in Eastern Australia. *Journal of Parasitology*, 82, 1017–1023.
- Bandoni, S. M. & Brooks, D. R. (1987a). Revision and phylogenetic analysis of the Gyrocotylidae Poche, 1926 (Platyhelminthes: Cercomeria: Cercomeromorphae). *Canadian Journal of Zoology*, 65, 2369–2389.
- Bandoni, S. M. & Brooks, D. R. (1987b). Revision and phylogenetic analysis of the Amphilinidae Poche, 1922 (Platyhelminthes: Cercomeria: Cercomeromorphae). *Canadian Journal of Zoology*, 65, 1110–1128.
- Baverstock, P. R., Fielke, R., Johnson, A. M., Bray, R. A. & Beveridge, I. (1991). Conflicting phylogenetic hypotheses for the

- parasitic platyhelminthes tested by partial sequencing of 18S ribosomal RNA. *International Journal for Parasitology*, 21, 329–339.
- Blair, D. (1993). The phylogenetic position of the Aspidogastrea within the parasitic flatworms inferred from ribosomal RNA sequence data. *International Journal for Parasitology*, 23, 169–178.
- Boeger, W. & Kritsky, D. C. (1993). Phylogeny and a revised classification of the Monogenea Bychowsky, 1937 (Platyhelminthes). *Systematic Parasitology*, 26, 1–32.
- Boeger, W. & Kritsky, D. C. (1997). Coevolution of the Monogenea (Platyhelminthes) based on a revised hypothesis of parasite phylogeny. *International Journal for Parasitology*, 27, 1495–1511.
- Brooks, D. R. (1982). Higher level classification of parasitic platyhelminths and fundamentals of cestode classification. In: D. F. Mettrick & S. S. Desser (Eds) *Parasites — Their World and Ours* (pp. 189–193). Amsterdam: Elsevier Biomedical.
- Brooks, D. R. (1989a). A summary of the database pertaining to the phylogeny of the major groups of parasitic platyhelminths, with a revised classification. *Canadian Journal of Zoology*, 67, 714–720.
- Brooks, D. R. (1989b). The phylogeny of the Cercomeria (Platyhelminthes: Rhabdocoela) and general evolutionary principles. *Journal of Parasitology*, 75, 606–616.
- Brooks, D. R., Bandoni, S. M., Macdonald, C. A. & O'Grady, R. T. (1989). Aspects of the phylogeny of the Trematoda Rudolphi, 1808 (Platyhelminthes: Cercomeria). *Canadian Journal of Zoology*, 67, 2609–2624.
- Brooks, D. R., Hoberg, E. P. & Weekes, P. J. (1991). Preliminary phylogenetic systematic analysis of the Eucestoda Southwell, 1930 (Platyhelminthes: Cercomeria). *Proceedings of the Biological Society of Washington*, 104, 651–668.
- Brooks, D. R. & McLennan, D. A. (1991). *Phylogeny, Ecology and Behavior: A Research Program in Comparative Biology*. Chicago: University of Chicago Press.
- Brooks, D. R. & McLennan, D. A. (1993a). Macroevolutionary trends in the morphological diversification among the parasitic flatworms (Platyhelminthes: Cercomeria). *Evolution*, 47, 495–509.
- Brooks, D. R. & McLennan, D. A. (1993b). Comparative study of adaptive radiations with an example using parasitic flatworms (Platyhelminthes: Cercomeria). *American Naturalist*, 142, 755–778.
- Brooks, D. R. & McLennan, D. A. (1993c). *Parascript: Parasites and the Language of Evolution*. Washington, DC: Smithsonian Institution University Press.
- Brooks, D. R. & McLennan, D. A. (1994). Historical ecology as a research programme: scope, limitations and the future. In: P. Eggleton & R. Vane-Wright (Eds) *Phylogenetics and Ecology*, Linnean Society Symposium Series no. 17 (pp. 1–27). London: Academic Press.
- Brooks, D. R., O'Grady, R. T. & Glen, D. R. (1985a). The phylogeny of the Cercomeria Brooks, 1982 (Platyhelminthes). *Proceedings of the Helminthological Society of Washington*, 52, 1–20.
- Brooks, D. R., O'Grady, R. T. & Glen, D. R. (1985b). Phylogenetic analysis of the Digenea (Platyhelminthes: Cercomeria) with comments on their adaptive radiation. *Canadian Journal of Zoology*, 63, 411–443.
- Brooks, D. R. & Overstreet, R. M. (1978). The family Liolopidae (Digenea) including a new genus and two new species from crocodilians. *International Journal for Parasitology*, 8, 267–273.
- Bullock, T. H. (1965). Chapter 9: Platyhelminthes. In: T. H. Bullock & G. A. Horridge (Eds) *Structure and Function in the Nervous System of Invertebrates* (pp. 535–576). San Francisco: W. H. Freeman.
- Cannon, L. R. G. (1982). Endosymbiotic Umagillids (Turbellaria) from Holothurians of the Great Barrier Reef. *Zoologica Scripta*, 11, 173–188.
- Cannon, L. R. G. (1987). Two new rhabdocoel turbellarians, *Umagilla pacifica* sp. n. & *U. karlingi* sp. n. (Umagillidae), endosymbiotic with holothurians (Echinodermata) from the Great Barrier Reef; and a discussion of sclerotic structures in the female system of the Umagillidae. *Zoologica Scripta*, 16, 297–303.
- Ching, H. & Leighton, B. J. (1993). The presence of *Udonella ophiodontis* in Washington and of *U. caligorum* in British Columbia. *Journal of the Helminthological Society of Washington*, 60, 137–140.
- Christensen, A. M. (1976). On the morphology and biology of *Kronborgia spiralis* (Baylis, 1949) (Turbellaria, Neorhabdocoela), with a note on its systematic position. *Ophelia*, 15, 77–97.
- Christensen, A. M. & Kannevorff, B. (1965). Life history and biology of *Kronborgia amphipodicola* Christensen and Kannevorff (Turbellaria, Neorhabdocoela). *Ophelia*, 2, 237–252.
- Cohn, L. (1902). Zwei neue Distomen. *Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheit, und Hygiene*, 32, 877–882.
- DeClerk, G. & Schockaert, E. (1995). Two peculiar new genera of Typhloplanoida from Western Indian Ocean. *Hydrobiologia*, 305, 3–9.
- de Pinna, M. C. C. (1991). Concepts and tests of homology in the cladistic paradigm. *Cladistics*, 7, 367–394.
- Ehlers, U. (1984). Phylogenetisches System der Plathelminthes. *Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg*, 27, 291–294.
- Ehlers, U. (1985a). *Das Phylogenetischen System der Plathelminthes*. Stuttgart: Gustav Fischer-Verlag.
- Ehlers, U. (1985b). Phylogenetic relationships within the Platyhelminthes. In: S. Conway Morris, J. D. George, R. Gibson & H. M. Platt (Eds) *The Origins and Relationships of Lower Invertebrates* (pp. 143–158). Oxford: Oxford University Press.
- Ehlers, U. (1986). Comments on a phylogenetic system of the Platyhelminthes. *Hydrobiologia*, 132, 1–12.
- Ehlers, U. (1995). The basic organization of the Platyhelminthes. *Hydrobiologia*, 305, 21–26.
- Ehlers, U. & Sopott-Ehlers, B. (1993). The caudal duo-gland adhesive system of *Jensenia angulata* (Platyhelminthes: Dalyelliidae): ultrastructure and phylogenetic significance (with comments on the phylogenetic position of the Temnocephalida and the polyphyly of the Cercomeria). *Microfauna Marina*, 8, 65–76.
- Eldredge, N. & Cracraft, J. (1980). *Phylogenetic Patterns and the Evolutionary Process*. New York: Columbia University Press.
- Fleming, L. (1986). Occurrence of symbiotic turbellarians in the oyster, *Crassostrea virginica*. *Hydrobiologia*, 132, 311–315.
- Fleming, L., Burt, M. D. B. & Bacon, G. B. (1981). On some commensal Turbellaria of the Canadian East Coast. *Hydrobiologia*, 84, 131–137.
- Gould, S. J. & Vrba, E.-S. (1982). Exaptation — a missing term from the science of form. *Paleobiology*, 8, 4–15.
- Hagen, D. W. (1967). Isolating mechanisms in three-spined sticklebacks (*Gasterosteus aculeatus*). *Journal of the Fisheries Research Board of Canada*, 24, 1637–1692.
- Harvey, P. H. & Pagel, M. (1991). *The Comparative Method in Evolutionary Biology*. Oxford: Oxford University Press.
- Hennig, W. (1966). *Phylogenetic Systematics*. Urbana, Illinois: University of Illinois Press.
- Hoberg, E. P., Mariaux, J. & Brooks, D. R. (in press). Phylogeny



- among orders of the Eucestoda (Cercomeromorphae): integrating morphology, molecules and total evidence. In: D. T. J. Littlewood & R. A. Bray (Eds) *Interrelationships of the Platyhelminthes* (in press). London: Taylor & Francis.
- Hoberg, E. P., Mariaux, J., Justine, J.-L., Brooks, D. R. & Weekes, P. J., Jr. (1997). Phylogeny of the orders of the Eucestoda (Cercomeromorphae) based on comparative morphology: historical perspectives and a new working hypothesis. *Journal of Parasitology*, *83*, 1128–1147.
- Hyman, L. H. (1951). *The Invertebrates: Platyhelminthes and Rhyncocoela*, Vol. II. New York: McGraw-Hill.
- Ivanov, A. V. (1952). Morphology of *Udonella caligorum* Johnston, 1835, and the position of Udonellidae in the systematics of the platyhelminthes. *Parasitological Collection of the Institute of Zoology, Academy of Sciences, USSR, XIV*, 112–163.
- Joffe, B. I. & Kornakova, E. E. (1998). *Notentera ivanovi* Joffe et al., 1997: a contribution to the question of phylogenetic relationships between 'turbellarians' and the parasitic Platyhelminthes (Neodermata). *Hydrobiologia*, *383*, 245–250.
- Jondelius, U. (1991). Unciliated body surface in 3 species of the Umagillidae (Dalyelloidea, Platyhelminthes). *Hydrobiologia*, *227*, 299–305.
- Jondelius, U. (1992). Sperm morphology in the Pterastericolidae (Platyhelminthes, Rhabdoceola): phylogenetic implications. *Zoologica Scripta*, *21*, 223–230.
- Jondelius, U. & Tholleson, M. (1993). Phylogeny of the Rhabdoceola (Platyhelminthes): a working hypothesis. *Canadian Journal of Zoology*, *71*, 298–308.
- Justine, J.-L. (1990). Phylogeny of parasitic platyhelminthes: a critical study of synapomorphies proposed on the basis of the ultrastructure of spermiogenesis and spermatozoa. *Canadian Journal of Zoology*, *69*, 1421–1440.
- Justine, J.-L. (1991). Cladistic study in the Monogenea (Platyhelminthes), based upon a parsimony analysis of spermiogenetic and spermatozoal ultrastructural characters. *International Journal for Parasitology*, *21*, 821–838.
- Justine, J.-L. (1993). Phylogenie des monogenes basee sur une analyse de parcimone des caracteres de l'ultrastructure de la spermiogenese et des spermatozoides incluant les resultats recents. *Bulletin Francaise de la Peche et Pisciculture*, *328*, 137–155.
- Justine, J.-L. (1995). Spermatozoal ultrastructure and phylogeny in the parasitic platyhelminthes. In: B. G. M. Jamieson, J. Ausio & J.-L. Justine (Eds) *Advances in Spermatozoal Phylogeny and Taxonomy, Memoires du Museum National d'Histoire Naturelle*, *166*, 55–86.
- Justine, J.-L. (1998a). Spermatozoa as phylogenetic characters for the Eucestoda. *Journal of Parasitology*, *84*, 385–408.
- Justine, J.-L. (1998b). Non-monophyly of the Monogeneans? *International Journal for Parasitology*, *28*, 1653–1657.
- Kanneworff, B. & Christensen, A. M. (1966). *Kronboria cardiacola* sp. nov. and endoparasitic turbellarian from North Atlantic shrimps. *Ophelia*, *3*, 65–80.
- Kluge, A. G. (1989). A concern for evidence and a phylogenetic hypothesis of relationships among *Epicrates* (Boidae, Serpentes). *Systematic Zoology*, *38*, 315–328.
- Kluge, A. G. (1997). Testability and the refutation and corroboration of cladistic hypotheses. *Cladistics*, *13*, 81–96.
- Kluge, A. G. (1998a). Total evidence or taxonomic congruence: cladistics or consensus classification. *Cladistics*, *14*, 151–158.
- Kluge, A. G. (1998b). Sophisticated falsification and research cycles: consequences for differential character weighting in phylogenetic systematics. *Zoologica Scripta*, *26*, 349–360.
- Kluge, A. G. (1999). The science of phylogenetic systematics: explanation, prediction, and test. *Cladistics*, *15*, 429–436.
- Koie, M. & Bresciani, J. (1973). On the ultrastructure of the larva of *Kronborgia amphipodcola* Christensen and Kanneworff, 1964 (Turbellaria, Neorhabdoceola). *Ophelia*, *12*, 171–203.
- Kornakova, E. E. & Joffe, B. I. (1999). A new variant of the neodermatan-type spermiogenesis in a parasitic 'turbellarian', *Notentera ivanovi* (Platyhelminthes) and the origin of the Neodermata. *Acta Zoologica (Stockholm)*, *80*, 135–151.
- Kuris, A. M. & Norton, S. F. (1985). Evolutionary importance of overspecialization: insect parasitoids as an example. *American Naturalist*, *126*, 387–391.
- Lee, D. L. (1972). The structure of the helminth cuticle. *Advances in Parasitology*, *10*, 347–379.
- Lindsey, C. C. (1962). Experimental study of meristic variation in a population of threespine sticklebacks, *Gasterosteus aculeatus*. *Canadian Journal of Zoology*, *40*, 271–312.
- Littlewood, D. T. J., Bray, R. A. & Clough, K. A. (1998). A phylogeny of the platyhelminthes: towards a total-evidence solution. *Hydrobiologia*, *383*, 155–160.
- Littlewood, D. T. J., Rohde, K. & Clough, K. A. (1999a). The interrelationships of all major groups of platyhelminthes: phylogenetic evidence from morphology and molecules. *Biological Journal of the Linnean Society*, *66*, 75–114.
- Littlewood, D. T. J., Rohde, K., Bray, R. A. & Herniou, E. A. (1999b). Phylogeny of the Platyhelminthes and the evolution of parasitism. *Biological Journal of the Linnean Society*, *68*, 257–287.
- Litvaitis, M. K. & Rohde, K. (1999). A molecular test of platyhelminth phylogeny: inferences from partial 28S rDNA sequences. *Invertebrate Biology*, *118*, 42–56.
- Looss, A. (1893). Ist der Laurer'sche Kanal der Trematoden eine Vagina? *Zentralblatt für Bakteriologie*, *13*, 808–819.
- Lundin, K. (2000). Phylogeny of the Nemertodermatida (Acoelomorpha, Platyhelminthes). A cladistic analysis. *Zoologica Scripta*, *29*, 65–74.
- McLennan, D. A. (1993). Phylogenetic relationships in the Gasterosteidae: an updated tree based on behavioral characters with a discussion of homoplasy. *Copeia*, *1993*, 318–326.
- McLennan, D. A. & Brooks, D. R. (in press). Phylogenetic systematics: five steps to enlightenment. In: J. Adrain, J. Edgecombe & B. Lieberman (Eds) *Fossils, Phylogeny, and Form: an Analytical Approach* (in press). London: Plenum Press.
- McLennan, D. A., Brooks, D. R. & McPhail, J. D. (1988). The benefits of communication between comparative ethology and phylogenetic systematics: a case study using gasterosteid fishes. *Canadian Journal of Zoology*, *66*, 2177–2190.
- Maddison, W. P. (1993). Missing data versus missing characters in phylogenetic analysis. *Systematic Biology*, *42*, 581–587.
- Mariaux, J. (1998). A molecular phylogeny of the Eucestoda. *Journal of Parasitology*, *84*, 114–124.
- Mollaret, I., Jamieson, B. G. M. & Justine, J.-L. (2000). Phylogeny of the Monopisthocotylea and Polypisthocotylea (Platyhelminthes) inferred from 28S rDNA sequences. *International Journal for Parasitology*, *30*, 171–185.
- Moran, N. A. (1988). The evolution of host-plant alternation in aphids: evidence for specialization as a dead end. *American Naturalist*, *132*, 681–706.

- Moran, N. A. (1994). Adaptation and constraint in the complex life cycles of animals. *Annual Review of Ecology and Systematics*, 25, 573–600.
- Nixon, K. C. & Davis, J. I. (1991). Polymorphic taxa, missing values and cladistic analysis. *Cladistics*, 7, 233–241.
- Noble, E., Noble, G., Schad, G. & MacInnes, A. (1989). *Parasitology: The Biology of Animal Parasites*. Philadelphia: Lea and Febiger.
- Noury-Srairi, N., Justine, J.-L. & Euzet, L. (1989a). Ultrastructure comparee de la spermiogenese et du spermatozoide de trois especes de *Paravortex* (Rhabdocoela, 'Dalyellioida', Graffillidae), Turbellaries parasites intestinaux de Mollusques. *Zoologica Scripta*, 18, 161–174.
- Noury-Srairi, N., Justine, J.-L. & Euzet, L. (1989b). Implications phylogenetiques de l'ultrastructure de la spermatogenese, du spermatozoide et de l'ovogenese du Turbellarie *Urastoma cyprinae* ('Prolecithophora', Urastomidae). *Zoologica Scripta*, 18, 175–185.
- Platnick, N. I., Griswold, C. E. & Coddington, J. A. (1991). On missing entries in cladistic analysis. *Cladistics*, 7, 337–343.
- Popper, K. R. (1960). *The Poverty of Historicism*. London: Routledge & Kegan Paul.
- Popper, K. R. (1968a). *The Logic of Scientific Discovery*. New York: Harper & Row.
- Popper, K. R. (1968b). *Conjectures and Refutations*. New York: Harper & Row.
- Popper, K. R. (1972). *Objective Knowledge: an Evolutionary Approach*. Oxford: Clarendon Press.
- Popper, K. R. (1976). *Unended Quest: an Intellectual Autobiography*. La Salle, Illinois: Open Court Publishers Co.
- Popper, K. R. (1992). *Realism and the Aim of Science*. London: Routledge.
- Remane, A. (1952). *Die Grundlagen des natuerlichen Systems, der vergleichenden Anatomie und der Phylogenetik*. Konigstein: Otto Koeltz.
- Rieger, R. & Tyler, S. (1985). Das homologiethorem in der Ultrastrukturforschung. In: J. A. Orr, G. P. Wagner & F. M. Wuketits (Eds) *Evolution, Ordnung und Erkenntnis* (pp. 101–122). Berlin: Parey.
- Rohde, K. (1986a). Ultrastructure of the flame cells and protonephridial capillaries of *Temnocephala*; implications for the phylogeny of parasitic platyhelminthes. *Zoologischer Anzeiger*, 216, 39–47.
- Rohde, K. (1986b). Ultrastructural studies of *Austrampbilina elongata* (Cestoda, Amphilinidea). *Zoomorphology*, 106, 91–102.
- Rohde, K. (1987). Ultrastructure of the flame cells and protonephridial capillaries of *Craspedella* and *Didymorchis* (Platyhelminthes, Rhabdocoela). *Zoomorphology*, 106, 346–351.
- Rohde, K. (1989). At least eight types of sense receptors in an endoparasitic flatworm: a counter-trend to sacculinization. *Naturwissenschaften*, 76, 383–385.
- Rohde, K. (1990). Phylogeny of the platyhelminthes, with special reference to parasitic groups. *International Journal for Parasitology*, 20, 979–1007.
- Rohde, K. (1991). The evolution of protonephridia of the platyhelminthes. *Hydrobiologia*, 227, 315–321.
- Rohde, K. (1994a). The origins of parasitism in the platyhelminthes. *International Journal for Parasitology*, 24 (8), 1099–1115.
- Rohde, K. (1994b). The minor groups of parasitic platyhelminthes. *Advances in Parasitology*, 33, 145–233.
- Rohde, K. (1996). Robust phylogenies and adaptive radiations: a critical examination of methods used to identify key innovations. *American Naturalist*, 148, 481–500.
- Rohde, K. (1998). Ultrastructure of the protonephridial system of the oncomiracidium of *Encotyllabe chironemi* (Platyhelminthes, Monopisthocotylea). *Microscopy Research and Technique*, 42, 212–217.
- Rohde, K., Cannon, L. R. G. & Watson, N. (1987b). Ultrastructure of epidermis, spermatozoa and flame cells of *Gyratix* and *Odontorhynchus* (Rhabdocoela, Kalyptorhynchia). *Journal of Submicroscopic Cytology*, 19, 585–594.
- Rohde, K., Johnson, A. M., Baverstock, P. R. & Watson, N. (1995). Aspects of the phylogeny of platyhelminthes based on 18S ribosomal DNA and protonephridial ultrastructure. *Hydrobiologia*, 305, 27–35.
- Rohde, K., Justine, J.-L. & Watson, N. (1989b). Ultrastructure of the flame bulbs of the monopisthocotylean Monogenea *Loimosina wilsoni* (Loimoidae) and *Calceostoma herculeana* (Calceostomidae). *Annales de Parasitologie Humaine et Comparee*, 64, 433–442.
- Rohde, K., Noury-Srairi, N., Watson, N., Justine, J.-L. & Euzet, L. (1990). Ultrastructure of the flame bulbs of *Urastoma cyprinae* (Platyhelminthes, 'Prolecithophora', Urastomidae). *Acta Zoologica*, 71, 211–216.
- Rohde, K. & Watson, N. (1993). Spermatogenesis in *Udonella* (Platyhelminthes, Udonellidea) and the phylogenetic position of the genus. *International Journal for Parasitology*, 23, 725–735.
- Rohde, K. & Watson, N. (1995). Comparative ultrastructural study of the posterior suckers of four species of symbiotic Platyhelminthes, *Temnocephala* sp. (Temnocephalida), *Udonella caligorum* (Udonellidea), *Anoplodiscus cirrusspiralis* (Monogenea: Monopisthocotylea), and *Philophthalmus* sp. (Trematoda: Digenea). *Folia Parasitologica*, 42, 11–28.
- Rohde, K., Watson, N. & Cannon, L. R. G. (1987a). Ultrastructure of the protonephridia of *Mesostoma* sp. (Rhabdocoela, 'Typhloplanoida'). *Journal of Submicroscopic Cytology*, 19, 107–112.
- Rohde, K., Watson, N. & Chisholm, L. A. (1999). Ultrastructure of the eyes of the larva of *Neobeterocotyle rbinobatidis* (platyhelminthes, Monopisthocotylea), and phylogenetic implications. *International Journal for Parasitology*, 29, 511–519.
- Rohde, K., Watson, N. & Jondelius, U. (1992). Ultrastructure of the protonephridia of *Syndisyrinx punicea* (Hickman, 1956) (Rhabdocoela: Umagillidae) and *Pterastericola pellucida* (Rhabdocoela: Pterastericolidae). *Australian Journal of Zoology*, 40, 385–399.
- Rohde, K., Watson, N. & Roubal, F. (1989a). Ultrastructure of flame bulbs, sense receptors, tegument and sperm of *Udonella* (Platyhelminthes) and the phylogenetic position of the genus. *Zoologischer Anzeiger*, 222, 143–157.
- Ronquist, F. (1994). Evolution of parasitism among closely related species: phylogenetic relationships and the origin of inquilinism in gall wasps (Hymenoptera, Cynipidae). *Evolution*, 48, 241–266.
- Shinn, G. L. & Christensen, A. M. (1985). *Kronborgia pugettensis* sp. nov. (Neorhabdocoela: Fecampiidae), an endoparasitic turbellarian infesting the shrimp *Heptacarpus kincaidi* (Rathbun), with notes on its life-history. *Parasitology*, 91, 431–447.
- Smyth, J. D. (1994). *Introduction to Animal Parasitology*. Cambridge: Cambridge University Press.
- Sopot-Ehlers, B. (1991). Comparative morphology of photoreceptors in free-living platyhelminthes — a survey. *Hydrobiologia*, 227, 231–239.
- Sopot-Ehlers, B. (1996). First evidence of mitochondrial lensing in two species of the 'Typhloplanoida' (Platyhelminthes,

- Rhabdozoa): phylogenetic implications. *Zoomorphology*, 116, 95–101.
- Sopott-Ehlers, B. (1998). Development and submicroscopic anatomy of male gametes in *Halammocortex nigrifrons* (Platyhelminthes, Rhabdozoa, 'Dalyellioida'): phylogenetic implications and functional aspects. *Zoomorphology*, 117, 213–222.
- Sopott-Ehlers, B. (2000). First evidence of unpigmented rhabdomeric photoreceptors in a Microstomum species (Platyhelminthes, Macrostomorpha, Microstomidae) and ultrastructure of photoreceptor-like ciliary aggregations. *Zoomorphology*, 119, 223–229.
- Sopott-Ehlers, B. & Ehlers, U. (1998). Characteristics of spermiogenesis and mature spermatozoa in *Gyratrix hermaphroditus* (Platyhelminthes, Rhabdozoa, Kalyptorhynchia). *Zoomorphology*, 118, 169–176.
- Sopott-Ehlers, B. & Ehlers, U. (1995). Modified sperm ultrastructure and some data on spermiogenesis in *Provortex tiferus* (Platyhelminthes, Rhabdozoa): phylogenetic implications for the Dalyellioida. *Zoomorphology*, 115, 41–49.
- Sopott-Ehlers, B. & Ehlers, U. (1997). Electronmicroscopical investigations of male gametes in *Ptychopera westbladi* (Platyhelminthes, Rhabdozoa, 'Typhloplanoida'). *Microfauna Marina*, 11, 193–208.
- Trouvé, S., Sasal, P., Jourdan, J., Renaud, F. & Morand, S. (1998). The evolution of life-history traits in parasitic and free-living platyhelminths: a new perspective. *Oecologia*, 115, 370–387.
- Wanntorp, H.-E., Brooks, D. R., Nilsson, T., Nylin, S., Ronqvist, F., Stearns, S. C. & Weddell, N. (1990). Phylogenetic approaches in ecology. *Oikos*, 57, 119–132.
- Watson, N. (1997). Proximo-distal fusion of flagella during spermiogenesis in the 'turbellarian' platyhelminth *Urastoma cyprinae*, and phylogenetic implications. *Invertebrate Reproduction and Development*, 32, 107–117.
- Watson, N. (1998a). Spermiogenesis and eyes with lenses in two Kalyptorhynch flatworm species, *Toia calceiformis* and *Nannorhynchides herdaensis* (Eukalyptorhynchia, Platyhelminthes). *Invertebrate Biology*, 117, 9–19.
- Watson, N. (1998b). Characteristics of protonephridial terminal organs and the eyes of four species of Trigonostominae (Platyhelminthes: Rhabdozoa). *Australian Journal of Zoology*, 46, 251–265.
- Watson, N. & Jondelius, U. (1995). Comparative ultrastructure of spermiogenesis and sperm in *Maebrethalia* sp. & *Bresslauilla relicta* (Platyhelminthes, Rhabdozoa). *Invertebrate Reproduction and Development*, 28, 103–112.
- Watson, N. & L'Hardy, J.-P. (1995). Origin of uniflagellate spermatozoon of *Baltoplana magna* (Platyhelminthes, Kalyptorhynchia). *Invertebrate Reproduction and Development*, 28, 185–192.
- Watson, N. & Rohde, K. (1994a). Ultrastructure of sperm and spermiogenesis in the Monocotyloid Monogeneans *Monocotyle helicophallus* and *Calicotyle australiensis* (Platyhelminthes). *International Journal for Parasitology*, 24, 1019–1030.
- Watson, N. & Rohde, K. (1994b). Ultrastructure of spermiogenesis and spermatozoa in *Phaenocora anomalozoa* (Platyhelminthes, Typhloplanida, Phaenocorinae). *Invertebrate Reproduction and Development*, 25, 237–246.
- Watson, N. & Rohde, K. (1995a). Re-examination of spermiogenesis of *Multicotyle purvisi* (Platyhelminthes, Aspidogastrea). *International Journal for Parasitology*, 25, 579–586.
- Watson, N. & Rohde, K. (1995b). Sperm and spermiogenesis of the 'Turbellaria' and implications for the phylogeny of the phylum platyhelminthes. In: B. G. M. Jamieson, J. Ausio & J.-L. Justine (Eds) *Advances in Spermatozoal Phylogeny and Taxonomy, Memoires du Museum National d'Histoire Naturelle*, 166, 37–54.
- Watson, N. & Rohde, K. (1995c). Ultrastructure of spermiogenesis and spermatozoa in the platyhelminthes *Actinodactylella blanchardi* (Temnocephalida, Actinodactylellidae), *Didymorchis* sp. (Temnocephalida, Didymorchidae) and *Gieystoria* sp. (Dalyelliida, Dalyelliidae), with implications for the phylogeny of the Rhabdozoa. *Invertebrate Reproduction and Development*, 27, 145–158.
- Watson, N., Rohde, K. & Sewell, K. B. (1995). Ultrastructure of spermiogenesis and spermatozoa of *Decadidymus gulosus*, *Temnocephala dendyi*, *T. minor*, *Craspedella* sp., *C. spenceri* and *Diceratocephala boschmai* (Platyhelminthes, Temnocephalida, Temnocephalidae), with emphasis on the intercentriolar body and zone of differentiation. *Invertebrate Reproduction and Development*, 27, 131–143.
- Watson, N., Rohde, K. & Williams, J. B. (1992). Ultrastructure of the protonephridial system of larval *Kronborgia isopodocola* (Platyhelminthes). *Journal of Submicroscopic Cytology and Pathology*, 24, 43–49.
- Watson, N. & Schockaert, E. (1996). Spermiogenesis and sperm ultrastructure in *Thylacorhynchus ambronensis* (Schizorhynchia, Kalyptorhynchia, Platyhelminthes). *Invertebrate Biology*, 115, 263–272.
- Watson, N. & Schockaert, E. (1997). Divergent protonephridial architecture within the Kalyptorhynchia (Platyhelminthes) and implications for the phylogeny of the Rhabdozoa. *Belgian Journal of Zoology*, 127, 139–158.
- Wiley, E. O. (1981). *Phylogenetics: the Theory and Practice of Phylogenetic Systematics*. New York: Wiley-Interscience.
- Wiley, E. O., Siegel-Causey, D., Brooks, D. R. & Funk, V. A. (1991). *The Complete Cladist: A Primer of Phylogenetic Procedures*. Lawrence, Kansas: University of Kansas Museum of Natural History Press.
- Wiley, E. O., Siegel-Causey, D., Brooks, D. R. & Funk, V. A. (in press). *The Complete Cladist: A Primer of Phylogenetic Procedures*. Lawrence, Kansas: University of Kansas Museum of Natural History Press.
- Wilkinson, M. (1995). Coping with abundant missing entries in phylogenetic inference using parsimony. *Systematic Biology*, 44, 501–514.
- Williams, J. B. (1993). Epidermal ultrastructure of the adult parasitic phase female and encapsulated larva of *Kronborgia isopodocola* (Platyhelminthes, Fecampiidae): phylogenetic implications. *International Journal for Parasitology*, 23, 1027–1037.
- Wirth, U. (1984). Die Struktur der Metazoen-Spermien und ihre Bedeutung für die Phylogenetik. *Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg*, 27, 295–362.
- Xylander, W. E. R. (1986). Ultrastrukturelle Befunde zur Stellung von *Gyrocotyle* im System der parasitischen Plathelminthen. *Verhandlungen der Deutschen Zoologischen Gesellschaft*, 79, 193.
- Xylander, W. E. R. (1987a). Ultrastructure of the lycophore larva of *Gyrocotyle urna* (Cestoda, Gyrocotylidae). I. Epidermis, neodermis anlage and body musculature. *Zoomorphology*, 106, 352–360.
- Xylander, W. E. R. (1987b). Ultrastructural studies on the reproductive system of Gyrocotylidae and Amphilinidea (Cestoda). II. Vitellaria, vitellocyte development and vitellocyte of *Gyrocotyle urna*. *Zoomorphology*, 107, 293–297.
- Xylander, W. E. R. (1987c). Ultrastructure of the lycophora larva of *Gyrocotyle urna* (Cestoda, Gyrocotylidae). II. Receptors and nervous system. *Zoologischer Anzeiger*, 219, 239–255.

- Xylander, W. E. R. (1987d). Das Protonephridialsystem der Cestoda: evolutiv Veränderungen und ihre mögliche funktionelle Bedeutung. *Verhandlungen der Deutschen Zoologischen Gesellschaft*, 80, 257–258.
- Xylander, W. E. R. (1988a). Ultrastructural studies on Udonellidae: evidence for a position within the Neodermata. *Fortschritte der Zoologie*, 36, 51–57.
- Xylander, W. E. R. (1988b). Ultrastructural studies on the reproductive system of Gyrocotylidea and Amphilinidea (Cestoda). *Parasitology Research*, 74, 363–370.
- Xylander, W. E. R. (1989). Ultrastructural studies on the reproductive system of Gyrocotylidea and Amphilinidea (Cestoda): spermatogenesis, spermatozoa, testes, and vas deferens of *Gyrocotyle*. *International Journal for Parasitology*, 19, 897–905.
- Xylander, W. E. R. (1990). Ultrastructure of the lycophore larva of *Gyrocotyle urna* (Cestoda, Gyrocotylidea). *Zoomorphology*, 109, 319–328.