

3-27-2015

A new species of *Neoregostoma* Monné and Giesbert, 1992 from Costa Rica and Panama (Coleoptera, Cerambycidae, Cerambycinae)

José Rafael Esteban Durán

Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria, esteban@inia.es

Antonio Santos-Silva

Universidade de São Paulo, toncriss@uol.com.br

Follow this and additional works at: <http://digitalcommons.unl.edu/insectamundi>

Esteban Durán, José Rafael and Santos-Silva, Antonio, "A new species of *Neoregostoma* Monné and Giesbert, 1992 from Costa Rica and Panama (Coleoptera, Cerambycidae, Cerambycinae)" (2015). *Insecta Mundi*. 908.

<http://digitalcommons.unl.edu/insectamundi/908>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0413

A new species of *Neoregostoma* Monné and Giesbert, 1992 from
Costa Rica and Panama (Coleoptera, Cerambycidae, Cerambycinae)

José Rafael Esteban Durán
Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria
Ministerio de Economía y Competitividad
Madrid, Spain

Antonio Santos-Silva
Museu de Zoologia
Universidade de São Paulo
São Paulo, São Paulo, Brazil

Date of Issue: March 27, 2015

José Rafael Esteban Durán and Antonio Santos-Silva
A new species of *Neoregostoma* Monné and Giesbert, 1992 from Costa Rica and
Panama (Coleoptera, Cerambycidae, Cerambycinae)
Insecta Mundi 0413: 1–4

ZooBank Registered: urn:lsid:zoobank.org:pub:88505B31-8856-45E1-9F4F-5577D08DFC08

Published in 2015 by

Center for Systematic Entomology, Inc.
P. O. Box 141874
Gainesville, FL 32614-1874 USA
<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Chief Editor: Paul E. Skelley, e-mail: insectamundi@gmail.com
Head Layout Editor: Eugenio H. Nearn
Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas
Review Editors: Listed on the **Insecta Mundi** webpage

Manuscript Preparation Guidelines and Submission Requirements available on the **Insecta Mundi** webpage at: <http://centerforsystematicentomology.org/insectamundi/>

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia
Museu de Zoologia, São Paulo, Brazil
Agriculture and Agrifood Canada, Ottawa, ON, Canada
The Natural History Museum, London, UK
Muzeum i Instytut Zoologii PAN, Warsaw, Poland
National Taiwan University, Taipei, Taiwan
California Academy of Sciences, San Francisco, CA, USA
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
Field Museum of Natural History, Chicago, IL, USA
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.
Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>
Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Eugenio H. Nearn

A new species of *Neoregostoma* Monné and Giesbert, 1992 from Costa Rica and Panama (Coleoptera, Cerambycidae, Cerambycinae)

José Rafael Esteban Durán

Instituto Nacional de Investigación y Tecnología Agrária y Alimentaria
Ministerio de Economía y Competitividad
Madrid, Spain
esteban@inia.es

Antonio Santos-Silva

Museu de Zoologia
Universidade de São Paulo
São Paulo, São Paulo, Brazil
toncriss@uol.com.br

Abstract. *Neoregostoma salazari*, a new species of Rhinotragini Thomson, 1861 from Alajuela (Costa Rica) and Panama (Panamá), is described, illustrated, and compared with *N. erythrocallum* (Lane, 1940) from Central Brazil. A key to *Neoregostoma* species is provided.

Key Words. Central America, key, Rhinotragini, taxonomy.

Resumo. *Neoregostoma salazari*, uma nova espécie de Rhinotragini Thomson, 1861 de Alajuela (Costa Rica) e Panamá (Panamá) é descrita, ilustrada e comparada com *N. erythrocallum* (Lane, 1940) do Brasil central. Uma chave para as espécies de *Neoregostoma* é fornecida.

Introduction

Nine species of *Neoregostoma* Monné and Giesbert, 1992 have been described from South America, and distributed in the sub-continent (Monné 2015) as shown below after the species name: *Neoregostoma spinipenne* (Fuchs, 1961) (French Guiana); *N. fasciatum* (Aurivillius, 1920) (Paraguay and Argentina); *N. erythrocallum* (Lane, 1940) [Brazil (Goiás)]; *N. coccineum* (Gory, 1831) [Brazil (Espírito Santo to Rio Grande do Sul)]; *N. discoideum* (Audinet-Serville, 1833) [Brazil (Rio de Janeiro)]; *N. luridum* (Klug, 1825) [Brazil (Goiás, São Paulo, Paraná), Paraguay, Argentina, and Uruguay]; *N. unicolor* (Aurivillius, 1920) (Brazil – known only from the holotype without mention of a precise locality); *N. giesberti* Clarke, 2007 [Bolivia (Santa Cruz)]; and *N. bettelai* Clarke, 2010 [Bolivia (Santa Cruz)]. The new species is the first of the genus known from Central America.

Interestingly, the specimens from Costa Rica, used to describe the new species, were collected on flowering trees by using long-handled entomological nets from a suspension bridge.

Materials

Collections and their acronyms are as follows:

ACMT – American Coleoptera Museum (James E. Wappes), San Antonio, Texas, USA

EDPC – José Rafael Esteban Durán, private collection, Madrid, Spain

MIUC – Museo de Insectos, Universidad de Costa Rica, San José, Costa Rica

***Neoregostoma salazari* sp. nov.**

(Fig. 1–6)

Description. Holotype female (Fig. 1, 3), total length 10.5 mm. General appearance elongate, compressed. Integument orangish-red, except some dark-brown spots, almost black. **Head:** Elongate, dorsally densely, coarsely punctate, with small curved grooves. Rostrum long; genae wide, somewhat cloven. Mandibles about one-third longer than gena; distal third black. Coronal suture distinct from about middle of inferior ocular lobes to margin of prothorax. Distance between lower eye lobes about equal to that between upper eye lobes. Antennae slightly longer than elytra, reaching base of elytral distal third; scape reddish, coarsely punctate; pedicel and antennomeres II–IV black; antennomeres V–XI slightly lighter; antennomere III longer than scape. **Thorax:** Prothorax barrel-like, about as long as wide, narrowed apically. Pronotum coarsely, densely, deeply punctate (punctures confluent in some areas); with short, sparse golden setae; with dark longitudinal band on each side of disk, slightly wider from anterior margin to middle, narrowed (mainly at distal fourth) to posterior margin. Scutellum sub-triangular; sides slightly curved inwards. Metepisterna partially black. **Elytra:** Laterally, sides narrowed at middle; surface coarsely, deeply, abundantly punctate; lateral carinae distinct from base to near apex; apex truncate, outer angle rounded, sutural angle slightly projected; sutural margins smooth, moderately thick, elevated; epipleural margins distinctly finer than sutural; the following parts black: humeri black, as is a narrow, longitudinal, sutural marking in basal 20 to 40 percent, and narrow band at apex. **Legs:** Robust; setae short, erect, sparse. Femora clavate, moderately short; profemora brown at apex; meso- and metafemora black at apex. Tibiae moderately enlarged towards apex. Tarsomere V darkened next to claws.

Male (Fig. 4–6). Distance between upper eye lobes equal to 0.70 times length of scape; distance between lower eye lobes, in frontal view, equal to 0.65 times length of scape. Antennae reaching distal third of elytra. Metepisterna very coarsely, abundantly punctate. Metasternum laterally very coarsely, sparsely punctate.

Variability. Paratype female (Fig. 2): antennomeres V–XI with orange basal ring; dark central spot of elytra, on suture, greatly reduced; an additional small dark spot on lateral side of elytra, below hind legs. Paratype male: lateral sides of prothorax with small, oblique, narrow dark-brown band; paler colored elytra; a large dark spot laterally on elytra, just below hind legs; apical dark bands on elytra larger.

Type material. Holotype female from COSTA RICA, *Alajuela*: Puentes de La Fortuna (near Arenal Volcano, Cantón de San Carlos), 11.V.2013 (on flowers of *Casearia aculeata* Jacq.), Marco A. Zumbado leg. (MIUC), length = 10.5 mm. Paratype female, same place as holotype, 20.V.2013 (on flowers of *Casearia aculeata* Jacq.), A. Hamer Salazar leg. (EDPC), length = 10.8 mm. Paratype male from PANAMA, *Panama*: Cerro Campana (2000'), 27–29.IV.2009, J. E. Wappes col. (ACMT), length = 9.9 mm.

Remarks. The general appearance of *Neoregostoma salazari* is similar to that of *N. erythrocallum* (Lane, 1940), but differs as follows: elytra with dark central band close to the suture (sometimes partly absent); pronotum with two dark bands; metafemora shorter, more distinctly thickened; basal third of metafemora orange; and metepisterna with large black area. In *N. erythrocallum* the elytra lack the dark central band close to the suture, the pronotum has a single, dark central band, the metafemora are longer and less thickened, the basal third of metafemora are dark-brown, and the metepisterna are entirely orange.

Etymology. The new species is named after Alberto Hamer Salazar Rodríguez, professor emeritus at the University of Costa Rica, for his dedication to the systematic investigation of the Class Insecta.

Key to species of *Neoregostoma*

1. Pronotum with a single dark central band. 2
- Pronotum with two dark bands or completely dark. 5

2(1).	Elytra with markings limited to a singular dark spot on humeri....	<i>N. erythrocallum</i> (Lane)	
–	Elytra with additional markings on elytra.		3
3(2).	Pronotal dark band incomplete.	<i>N. discoideum</i> (Audinet-Serville)	
–	Pronotal dark band complete from base to apex.		4
4(3).	Elytra with transverse dark band, near the middle, complete to suture.	<i>N. bettelai</i> Clarke	
–	Transverse dark band of elytra not complete to suture.	<i>N. coccineum</i> (Gory)	
5(1).	Pronotum completely dark.		6
–	Pronotum light colored with darker areas or entirely light.		7
6(5).	Elytra completely dark.	<i>N. unicolor</i> (Aurivillius)	
–	Elytra light colored with wide dark central band.	<i>N. fasciatum</i> (Aurivillius)	
7(5).	Elytra nearly all light colored.		8
–	Elytra with large dark areas.		9
8(7).	Elytra without dark central band close to the suture; elytral apex with spicule at outer angle; meso- and metafemora with middle of club dark.	<i>N. spinipenne</i> (Fuchs)	
–	Elytra with dark central band close to the suture; elytral apex without spicule at outer angle; meso- and metafemora dark at apex.	<i>N. salazari</i> sp. nov.	
9(7).	Elytral markings longitudinal; apices partially dark with light colored markings.	<i>N. luridum</i> (Klug)	
–	Elytral markings transverse; apices broadly dark.	<i>N. giesberti</i> Clarke	

Acknowledgments


To Alberto Hamer Salazar Rodríguez (University of Costa Rica, in Greece), and Marco Antonio Zumbado Echevarría (University of Costa Rica, in San Ramón) for the loan of study specimens. To James E. Wappes (ACMT) for the loan of study specimens, and for corrections to the manuscript. To Larry G. Bezark for corrections to the manuscript.

Literature Cited

Monné, M. A. 2015. Catalogue of the Cerambycidae (Coleoptera) of the Neotropical Region. Part I. Subfamily Cerambycinae. 1166 p. Available at: <http://www.cerambyxcat.com/> (Last accessed on January 10, 2015).

Received February 18, 2015; Accepted March 17, 2015.

Review Editor Paul Skelley.



Figures 1–6. *Neoregostoma salazari* sp. nov. 1) Holotype female, dorsal view. 2) Paratype female, dorsal view. 3) Holotype female, lateral view. 4) Paratype male, dorsal view. 5) Paratype male, ventral view. 6) Paratype male, lateral view.