

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Historical Materials from University of Nebraska-  
Lincoln Extension

Extension

---

2003

## NF03-580 Upper Safe Levels of Intake for Adults: Vitamins, Macrominerals, and Trace Minerals

Judy A. Driskell

University of Nebraska - Lincoln, [jdriskell1@unl.edu](mailto:jdriskell1@unl.edu)

Follow this and additional works at: <http://digitalcommons.unl.edu/extensionhist>



Part of the [Agriculture Commons](#), and the [Curriculum and Instruction Commons](#)

---

Driskell, Judy A., "NF03-580 Upper Safe Levels of Intake for Adults: Vitamins, Macrominerals, and Trace Minerals" (2003). *Historical Materials from University of Nebraska-Lincoln Extension*. 1770.

<http://digitalcommons.unl.edu/extensionhist/1770>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# Upper Safe Levels of Intake for Adults: Vitamins, Macrominerals, and Trace Minerals

By Judy A. Driskell, Professor of Nutritional Science and Dietetics

Our bodies need vitamins and essential minerals; however, if taken in large amounts, they can adversely affect our health. In fact large amounts of many of the vitamins and minerals can be toxic.

## Nutritional Status

Individuals can have deficient, adequate, or toxic intakes of any essential nutrient. A **deficiency** occurs when the intake of the essential nutrient is too low to meet a person's need for that specific nutrient. **Adequacy** occurs when a person gets enough but not too much of a nutrient. Nutrient **toxicity** occurs when a person gets an overdose of a given nutrient.

## Dietary Reference Intakes

The National Academy of Sciences, Institute of Medicine, has established **Dietary Reference Intakes** (DRIs), which can be used in planning and assessing diets of the healthy general population, as well as for other purposes. The Dietary Reference Intakes are a set of nutrient-based reference values: the **Estimated Average Requirement** (EAR), the **Recommended Dietary Allowance** (RDA), the **Adequate Intake** (AI), and the **Tolerable Upper Intake Level** (UL). These terms are defined in *Table 1*. An essential nutrient has either an EAR and RDA or an AI, for a specific gender:life stage group (such as female, 19 to 30 years) depending on whether sufficient data are available to set an EAR. If the data are insufficient an AI is given for that nutrient for a gender:life stage group. In that many individuals consume **fortified foods** (containing nutrients added during food processing) and **dietary supplements**, ULs have been set for many of the essential nutrients for a specific gender:life stage group. The UL is the highest level of daily nutrient intake

that is considered to be safe for almost all individuals in a gender:life stage group. These ULs are based on intakes from food (including fortified food), water, and dietary supplements unless otherwise stated. The UL is for daily use over a long period of time. ULs are not available for all essential nutrients for each gender:life stage grouping as data regarding adverse effects are frequently limited.

## Table 1. Dietary Reference Intakes

**Estimated Average Requirement** (EAR): the average daily nutrient intake level estimated to meet the requirement of half the healthy individuals in a particular life stage and gender group.

**Recommended Dietary Allowance** (RDA): the average daily dietary nutrient intake level sufficient to meet the nutrient requirement of nearly all (97 to 98 percent) healthy individuals in a particular life stage and gender group.

**Adequate Intake** (AI): the recommended average daily intake level based on observed or experimentally determined approximations or estimates of nutrient intake by a group (or groups) of apparently healthy people that are assumed to be adequate — used when an RDA cannot be determined.

**Tolerable Upper Intake Level** (UL): the highest average daily intake level that is likely to pose no risk of adverse health effects to almost all individuals in the general population. As intake increases above the UL, the potential risk of adverse effects may increase.

Directly quoted from Institute of Medicine, National Academy of Sciences, *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*, p. 3. Washington, DC: National Academy Press, 2000.

## Nutrient Toxicity

The UL is calculated from hazard identification (primarily evidence of adverse effects) and dose-response data. The

appropriate critical adverse effect(s) for the nutrient is considered in setting the **no-observed-adverse-effect level** (NOAEL). The NOAEL is the highest intake or dose of a nutrient at which no adverse effects have been observed in individuals studied. If no adequate data are available to determine the NOAEL, then a **lowest-observed-adverse-effect level** (LOAEL) is used. The LOAEL is the lowest intake or dose at which an adverse effect(s) has been demonstrated.

The Tolerable Upper Intake Levels, or ULs, of Vitamins and Essential Macro- and Trace Minerals for Adults are given in *Table II*. DRIs for the electrolyte minerals (sodium, chloride, and potassium) have not yet been established. The reasons given by the Institute of Medicine, National Academy of Sciences, for the UL and the basis for the UL are listed in *Table II*.

The current RDAs or AIs as well as the Daily Values for these essential nutrients are also given in *Table II*. **Daily Values** are set by the Food and Drug Administration for use in nutritional labeling of food products and dietary supplements. RDAs or AIs are not the same as Daily Values. Information from older releases of the RDAs were used in setting the Daily Values. The RDAs or AIs and Daily Values for these essential nutrients were included in *Table II* so that the reader can compare these values to the ULs. These values are also useful in planning and evaluating diets, as diets should meet the RDAs or AIs for an essential nutrient without exceeding the ULs.

Diets for healthy individuals should not routinely exceed the UL for any essential nutrient; however, intakes above the UL may be appropriate for investigation within

clinical research trials. Nutrient ULs are not meant to apply to individuals who may be given an essential nutrient under medical supervision.

## References

- Institute of Medicine, National Academy of Sciences, *Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride*. Washington, DC: National Academy Press, 1997.
- Institute of Medicine, National Academy of Sciences, *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B<sub>6</sub>, Folate, Vitamin B<sub>12</sub>, Pantothenic Acid, Biotin, and Choline*. Washington, DC: National Academy Press, 1998.
- Institute of Medicine, National Academy of Sciences, *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*. Washington, DC: National Academy Press, 2000.
- Institute of Medicine, National Academy of Sciences, *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*. Washington, DC: National Academy Press, 2001.
- Institute of Medicine, National Academy of Sciences, *Dietary Reference Intakes: Applications in Dietary Planning*. Washington, DC: National Academy Press, 2003.
- U.S. Food and Drug Administration, *A Food Labeling Guide: Reference Values for Nutrition Labeling*. On-line <http://vm.cfsan.fda.gov/~dms/flg-7a.html>, accessed April 18, 2003.

**Table II. Tolerable Upper Intake Levels (ULs) of Vitamins and Essential Macro- and Trace Minerals for Adults.**

| <i>Nutrient</i>          | <i>UL<sup>a</sup></i><br>(per day) | <i>Reasons given by the Institute of Medicine, National Academy of Sciences, for UL</i>  | <i>Basis for UL</i>   |   |  |
|--------------------------|------------------------------------|--|---|---|--|
|                          |                                    |  | (NOAEL or LOAEL <sup>b</sup> per day)                                       | <i>RDA or AI<sup>c</sup></i><br>(per day)<br><i>Men/Women</i> | <i>Daily Value<sup>d</sup></i><br>(per day)                            |
| Vitamin A (preformed)    | 3,000 mcg                          | The critical adverse effect is liver abnormalities except for women of childbearing age in which it is teratogenic. Other adverse effects include nausea, vomiting, headache, increased cerebrospinal fluid pressure, vertigo, blurred vision, muscular incoordination, bulging, fontanel in infants, nervous system changes, and bone and skin abnormalities. | LOAEL = 14,000 mcg, except for women of childbearing age, NOAEL = 3,000 mcg | 900/700 mcg   | 5000 IU (1500 mcg preformed vitamin A)                                 |
| Vitamin D                | 50 mcg                             | The critical adverse effect is hypercalcemia. Other adverse effects include anorexia, nausea, vomiting, increased thirst and urination, metastatic calcification of soft tissues (kidneys, blood vessels, heart, lungs), and renal disorders.  | NOAEL = 60 mcg  | 5, 10, or 15 mcg <sup>e</sup>                                 | 400 IU or 10 mcg   |
| Vitamin E <sup>f,g</sup> | 1,000 mg $\alpha$ -tocopherol      | No adverse effects from consumption of vitamin E naturally occurring in foods. The critical adverse effect of high intakes from fortified foods, dietary supplements, or pharmacologic agents is increased tendency to hemorrhage. Adults deficient in vitamin K, including those taking coumarin drugs, have increased risk of coagulation defects.           | LAOEL = 500 mg/kg   | 15/15 mg $\alpha$ -tocopherol                                 | 30 IU (22 mg of natural d- $\alpha$ -tocopherol acetate <sup>h</sup> ) |
| Vitamin K                | ND <sup>i</sup>                    | No adverse effects from consumption from foods or dietary supplements.   | NE <sup>j</sup>   | 120/90 mcg  | 80 mcg   |

| <i>Nutrient</i>  | <i>UL<sup>a</sup></i><br>(per day) | <i>Reasons given by the Institute of Medicine,<br/>National Academy of Sciences, for UL</i>   | <i>Basis for UL</i><br>(NOAEL or<br>LOAEL <sup>b</sup><br>per day) | <i>RDA or AI<sup>c</sup></i><br>(per day)<br><i>Men/Women</i> | <i>Daily<br/>Value<sup>d</sup></i><br>(per day) |
|--|------------------------------------|---|--|---|---|
| Vitamin C  | 2,000mg                            | The critical adverse effects are osmotic diarrhea and gastrointestinal disturbances. Other possible effects include increased oxalate excretion and kidney stone formation, uric acid excretion, pro-oxidant effects, rebound scurvy, increased iron absorption leading to iron overload, reduced vitamin B <sub>12</sub> and copper status, increased oxygen demand, and erosion of dental enamel. | LAOEL=<br>3000mg   | 90/75 mg  | 60 mg   |
| Thiamin<br>(vitamin B <sub>1</sub> )                         | ND                                 | No adverse effects from consumption from foods or dietary supplements. There have been occasional reports of anaphylaxis to parenteral thiamin as well as pruritus due to allergic sensitivity to thiamin injection.  | NE   | 1.2/1.1 mg  | 1.5 mg  |
| Riboflavin<br>(vitamin B <sub>2</sub> )                      | ND                                 | No adverse effects from consumption from foods or dietary supplements.  | NE   | 1.3/1.1 mg  | 1.7 mg  |
| Niacin <sup>e</sup><br>(nicotinic<br>acid &<br>nicotinamide) | 35 mg                              | No adverse effects from consumption of naturally occurring niacin in foods. The UL for nicotinic acid is based on vasodilation (flushing). Nicotinamide appears not to be associated with the flushing effects. There are gastrointestinal effects for patients treated with nicotinic acid. Hepatic toxicity has been reported in patients treated medically with the vitamin.                     | LOAEL=<br>50 mg  | 16/14 mg  | 20 mg   |
| Vitamin B <sub>6</sub>                                       | 100mg                              | No adverse effects from consumption from food sources. The critical adverse effect from high intake is neuropathy.  | NOAEL=<br>200mg  | 1.3, 1.5, or<br>1.7 mg <sup>k</sup>                           | 2.0 mg  |
| Folate <sup>g</sup>  | 1,000 mcg                          | No adverse effects have been associated with the consumption of folate levels in foods or fortified foods. Excess folate may precipitate or exacerbate the neurological damage of vitamin B <sub>12</sub> deficiency.   | LOAEL=<br>5 mg   | 400/400 mcg <sup>l</sup>                                      | 400 mcg   |
| Vitamin B <sub>12</sub>                                      | ND                                 | No adverse effects from consumption from foods or dietary supplements.  | NE   | 2.4/2.4 mcg <sup>m</sup>                                      | 6.0 mcg   |
| Pantothenic<br>Acid  | ND                                 | No adverse effects have been associated with high intakes.  | NE   | 5/5 mg  | 10 mg   |
| Biotin   | ND                                 | No adverse effects from foods or dietary supplements.   | NE   | 30/30 mcg   | 300 mcg   |
| Choline  | 3.5 g                              | The critical effect is hypotension and fishy body odor is a secondary consideration. Also nausea and diarrhea.  | LOAEL=<br>7.5 mg   | 550/425 mg  | NE  |
| Calcium  | 2.5 g                              | The critical effect is kidney stone formation or milk-alkali syndrome (hypercalcemia and renal insufficiency). Also affects absorption of iron, zinc, magnesium, and phosphorus.  | LOAEL=<br>5000mg   | 1,000 or<br>1,200 mg <sup>n</sup>                             | 1,000 mg  |
| Phosphorus   | 3 or 4 g <sup>o</sup>              | The critical effect is hyperphosphatemia. Other effects include hypocalcemia, adjustments in calcium-regulating hormones, and calcification of nonskeletal tissues (especially kidneys).  | NOAEL=<br>10,200 mg  | 700/700 mg  | 1,000 mg  |
| Magnesium  | 350mg                              | Adverse effects are from nonfood sources, such as magnesium salts used for pharmacologic purposes. The critical effect is osmotic diarrhea. Other effects include nausea, abdominal cramping, serious neurological and cardiac symptoms, and death.   | LOAEL=<br>360mg  | 310,320,400,<br>or 420 mg <sup>p</sup>                        | 400 mg  |
| Fluoride   | 10mg                               | The critical effects are enamel fluorosis and skeletal fluorosis.   | NOAEL=<br>10mg   | 4/3 mg  | NE  |

| <i>Nutrient</i> | <i>UL<sup>a</sup></i><br>(per day) | <i>Reasons given by the Institute of Medicine,<br/>National Academy of Sciences, for UL</i>   | <i>Basis for UL</i><br>(NOAEL or<br>LOAEL <sup>b</sup><br>per day) | <i>RDA or AI<sup>c</sup></i><br>(per day)<br><i>Men/Women</i> | <i>Daily<br/>Value<sup>d</sup></i><br>(per day) |
|-----------------|------------------------------------|---|--|---|---|
| Selenium        | 400mcg                             | The critical effects are hair and nail brittleness and loss. Other effects include gastrointestinal disturbances, skin rash, garlic breath odor, fatigue, irritability, and nervous system disorders.   | NOAEL=<br>800mcg   | 55/55 mcg   | 70 mcg  |
| Iron            | 45 mg                              | The critical adverse effect is gastrointestinal side effects. Other effects include impaired zinc absorption, increased risk for vascular disease and cancer, and systemic iron overload.   | LOAEL=<br>70 mg  | 8 or 18 mg <sup>e</sup>                                       | 18 mg   |
| Copper          | 10mg                               | The critical adverse effect is liver damage. Other effects include abdominal pain, cramps, nausea, diarrhea, and vomiting.  | NOAEL=<br>10mg   | 900/900 mcg   | 2.0mg   |
| Zinc            | 40mg                               | No adverse effects from consumption from foods. The critical adverse effect is the influence of excess zinc on copper metabolism. Other effects include epigastric pain, nausea, vomiting, loss of appetite, abdominal cramps, diarrhea, headaches, and immune response impairment.   | LOAEL=<br>60mg   | 11/8mg  | 15 mg   |
| Iodine          | 1,100mcg                           | The critical adverse effect is elevated serum thyroid stimulation hormone concentrations. Acute responses include burning of mouth, throat, and stomach, abdominal pain, fever, nausea, vomiting, diarrhea, weak pulse, cardiac irritability, coma, and cyanosis. Also can produce goiter, increased risk of thyroid papillary cancer, and iodemia. | LOAEL=<br>1,700mcg   | 150/150mcg  | 150mcg  |
| Manganese       | 11 mg                              | The critical adverse effects are elevated blood magnesium concentration and neurotoxicity.  | NOAEL=<br>11 mg  | 2.3/1.8 mg  | 2.0mg   |
| Chromium        | ND                                 | No adverse effects have been convincingly associated with excess intake of chromium from food or dietary supplements.   | NE   | 20,25,30,<br>or 35 mcg <sup>f</sup>                           | 120mcg  |
| Molybdenum      | 2mg                                | The UL is based upon impaired reproduction and fetal development in rats and mice. Molybdenum compounds appear to have low toxicity in humans, but inadequate data are available.   | NOAEL=<br>0.9 mg/kg<br>(from rat data)                             | 45/45 mcg   | 75 mcg  |

<sup>a</sup>UL = The maximum level of daily nutrient intake that is likely to pose no risk of adverse effects.

<sup>b</sup>NOAEL = No-observed-adverse-effect level; LOAEL = Lowest-observed-adverse-effect level.

<sup>c</sup>RDA = Recommended Dietary Allowance; AI = Adequate Intake.

<sup>d</sup>The Daily Value is used in nutritional labeling of food products and dietary supplements. The values given are for individuals age 4 years and above.

<sup>e</sup>5 mcg vitamin D for 31-50 year olds, 10 mcg for 51-70 year olds, and 15 mcg for those above 70 years.

<sup>f</sup>As  $\alpha$ -tocopherol; applies to any form of supplemental  $\alpha$ -tocopherol.

<sup>g</sup>The ULs for vitamin E, niacin, and folate apply to synthetic forms from supplements, fortified foods, or a combination of supplements and fortified foods.

<sup>h</sup>The conversion of IUs to mg  $\alpha$ -tocopherol depends on the form.

<sup>i</sup>ND = Not determinable due to lack of data of adverse effects.

<sup>j</sup>NE = Not established.

<sup>k</sup>1.3 mg vitamin B<sub>6</sub> for 19-50 year olds, 1.5 mg for women age 51 and above, and 1.7 mg for men age 51 and above.

<sup>l</sup>All women capable of becoming pregnant should consume 400 mcg from supplements or fortified foods in addition to intake of food folate from a varied diet.

<sup>m</sup>Adults over 50 years are advised to meet their RDA mainly by consuming foods fortified with vitamin B<sub>12</sub> or a supplement containing vitamin B<sub>12</sub>.

<sup>n</sup>1,000 mg calcium for 19-50 year olds and 1,200 mg for those 51 years and older.

<sup>o</sup>4 g phosphorus for adults 19-70 years old and 3 g for those above 70 years.

<sup>p</sup>310 mg magnesium for women 19-30 years, 320 mg for women 31 years and above, 400 mg for men 19-30 years, and 420 mg for men 31 years and above.

<sup>q</sup>8 mg for men and for women 51 years and older and 18 mg for women 19-50 years.

<sup>r</sup>20 mcg for women 31 years old and above, 25 mcg for women 19-30 years old, 30 mcg for men 31 years old and above, and 35 mcg for men 19-30 years old.

**Fileunder: FOODSANDNUTRITION  
C-20, Nutritive Value of Foods**

Issued October 2003

Extension is a Division of the Institute of Agriculture and Natural Resources at the University of Nebraska–Lincoln cooperating with the Counties and the United States Department of Agriculture.

University of Nebraska–Lincoln Extension educational programs abide with the nondiscrimination policies of the University of Nebraska–Lincoln and the United States Department of Agriculture.