

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Robert Katz Publications

Research Papers in Physics and Astronomy

November 1992

Katz Model Prediction of *Caenorhabditis elegans* Mutagenesis on STS-42

Francis A. Cucinotta

Langley Research Center, Hampton VA, francis.cucinotta@unlv.edu

John W. Wilson

Langley Research Center, Hampton VA

Robert Katz

University of Nebraska-Lincoln, rkatz2@unl.edu

Gautam Badhwar

Lyndon B. Johnson Space Center, Houston TX

Follow this and additional works at: <https://digitalcommons.unl.edu/physicskatz>



Part of the [Physics Commons](#)

Cucinotta, Francis A.; Wilson, John W.; Katz, Robert; and Badhwar, Gautam, "Katz Model Prediction of *Caenorhabditis elegans* Mutagenesis on STS-42" (1992). *Robert Katz Publications*. 16.

<https://digitalcommons.unl.edu/physicskatz/16>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Robert Katz Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Katz Model Prediction of *Caenorhabditis elegans* Mutagenesis on STS-42

Francis A. Cucinotta, *Langley Research Center, Hampton VA*

John W. Wilson, *Langley Research Center, Hampton VA*

Robert Katz, *University of Nebraska-Lincoln, Lincoln NE*

Gautam Badhwar, *Lyndon B. Johnson Space Center, Houston TX*

National Aeronautics and Space Administration Technical Memo NASA TM-4383

NASA Langley Research Center
Hampton, Virginia

November 1992

Summary: The Katz track structure model has been applied to describe recessive lethal mutagenesis in the nematode *Caenorhabditis elegans* after exposure to heavy ions. Based on models of the cosmic-ray environment and heavy-ion transport, mutation rates for the International Microgravity Laboratory 1 (IML-1) experiment on the Space Transportation System 42 (STS-42) are predicted and the results are discussed.

Katz Model Prediction of
Caenorhabditis elegans
Mutagenesis on STS-42

Francis A. Cucinotta and John W. Wilson
Langley Research Center
Hampton, Virginia

Robert Katz
University of Nebraska
Lincoln, Nebraska

Gautam D. Badhwar
Lyndon B. Johnson Space Center
Houston, Texas



National Aeronautics and
Space Administration

Office of Management

Scientific and Technical
Information Program

1992

N93-13023

Unclass

H1/52 0128394

(NASA-TM-4383) KATZ MODEL
PREDICTION OF CAENORHABDITIS
ELEGANS MUTAGENESIS ON STS-42
(NASA) 10 p

Acknowledgments

Thanks to Bill Atwell and Mark Weyland, Rockwell International, Houston, Texas, for providing the energy spectrum of trapped protons and the shielding distributions for the Space Shuttle.

Summary

The Katz track structure model has been applied to describe recessive lethal mutagenesis in the nematode *Caenorhabditis elegans* after exposure to heavy ions. Based on models of the cosmic-ray environment and heavy-ion transport, mutation rates for the International Microgravity Laboratory 1 (IML-1) experiment on the Space Transportation System 42 (STS-42) are predicted and the results are discussed.

Introduction

The nematode *Caenorhabditis elegans* (*C. elegans*) is being studied (refs. 1 to 3) as a model radiobiological system that can provide information on mutagenesis after exposure to galactic cosmic radiation (GCR). *C. elegans* is particularly well suited for study because this nematode has a relatively small, fixed number of cells with only six pairs of chromosomes; many of its genes have been identified and mapped. Nelson (refs. 2 and 3) has reviewed several properties of *C. elegans* that favor its use in understanding the biological effects of heavy charged particles, especially for genetic processes, and its mutagenesis rates have been measured for several heavy-ion beams (ref. 3). The International Microgravity Laboratory 1 (IML-1) mission on the Space Transportation System 42 (STS-42) carried an experimental package containing *C. elegans* to study the effects of microgravity and tested ground-based understanding of GCR effects in space conditions if significant exposures occurred. The STS-42 mission duration of about 7 days in an orbit of 57° by 163 n.mi. exposed the *C. elegans* package to a complicated mixture of particles that vary in composition with the amount and type of shielding surrounding the package. Major concerns were anomalous effects on radiation response from microgravity or physiological stress from spaceflight.

Several considerations could affect our extrapolations from ground-based to flight experiments: fluxes one or two orders of magnitude below those achieved using accelerators, protracted exposure, and the broad charge and velocity spectrum of the GCR. The track structure model of Katz et al. (refs. 4 to 9) utilizes a charge- and velocity-dependent fit to radiobiology experiments and is the only model that can predict cellular endpoint response for the complete GCR spectrum. Biological response at low fluence is described by an action cross section representing the probability that a single ion leads to the endpoint in question. Laboratory measurements at high dose for photons and with a few ions allow for a parametric determination of the cross section in terms of ion charge and velocity that can be applied to the

GCR spectrum. Compared with other radiobiological models, the zero initial slope for X rays assumed by the Katz track structure model for multitarget endpoints leads to the maximum prediction for GCR effectiveness.

In this report, we consider experimental data for ion (ref. 3) and photon (refs. 3 to 10) production of recessive lethal mutations in *C. elegans* for response parameters to describe this endpoint in the Katz track structure model. This is the first application of the track structure model to an animal system. We then consider the GCR and trapped particle spectrum that is expected for IML-1 and predict the expected mutation rates by using models of the cosmic-ray environment, accurate transport codes, and shielding models.

Track Structure Model

The track structure model of Katz et al. (ref. 4) attributes biological damage from energetic ions to the secondary electrons (delta rays) produced along the ion path. The deleterious effects of energetic ions are correlated with those of gamma rays by the assumption that the response in sensitive sites near the ion path is part of a larger system irradiated with gamma rays at the same dose. The response from ion effects is then determined by knowledge of the gamma-ray response and the delta-ray dose surrounding the ion path. For a multitarget response with target number m , the inactivation of cells by gamma rays is assumed to follow a Poisson distribution reflecting the random accumulation of sublethal damage with a radiosensitivity parameter D_0 .

When ions inactivate the cells, two modes are identified: "ion kill," which corresponds to intratrack effects, and "gamma kill," which corresponds to intertrack effects. Here, the ion-kill mode is unique to ions corresponding to single-particle inactivation of cells described by the cross section σ . The action cross section for a sensitive site is determined as

$$\sigma = \int_0^\infty 2\pi t dt [1 - \exp(-\bar{D}/D_0)]^m \quad (1)$$

where \bar{D} is the average dose at the sensitive site from the ion delta rays. The cell damage is divided by Katz et al. (ref. 4) into a grain-count regime, where inactivation occurs randomly along the path of the particle, and into a track-width regime, where many inactivations occur and are said to be distributed like a "hairy rope." In the grain-count regime, σ may be parameterized as

$$\sigma = \sigma_0 [1 - \exp(-Z^{*2}/\kappa\beta^2)]^m \quad (2)$$

where σ_0 is the plateau value of the cross section, β is the ion velocity, the effective charge number is given by

$$Z^* = Z \left[1 - \exp \left(-125\beta/Z^{2/3} \right) \right] \quad (3)$$

and κ is a parameter related to the radius of the sensitive site a_0 by

$$D_0 \frac{a_0^2}{\kappa} \approx 2 \times 10^{-11} \text{ Gy/cm} \quad (4)$$

The transition from the grain-count to the track-width regime (refs. 4 and 5) occurs at a value of $Z^{*2}/(\kappa\beta^2)$ on the order of 4. At lower values of Z^{*2} we are in the grain-count regime and, at higher values, the track-width regime.

In the grain-count regime, the fraction of the cells damaged in the ion-kill mode is $P = \sigma/\sigma_0$, where σ is the single-particle-inactivation cross section and P is the probability of damage. In the track-width regime, where $\sigma > \sigma_0$, it is assumed that $P = 1$. In the track model a fraction of the ion dose $(1 - P)$ is assumed to add cumulatively to the ions from other particles to inactivate cells in the gamma-kill mode. The surviving fraction of a cellular population N_0 , whose response parameters are m , D_0 , and κ or a_0 after irradiation by a fluence of particles F , is then written as

$$\frac{N}{N_0} = \Pi_i \Pi_\gamma \quad (5)$$

where

$$\Pi_i = \exp(-\sigma F) \quad (6)$$

is the ion-kill survival probability and

$$\Pi_\gamma = 1 - [1 - \exp(-D_\gamma/D_0)]^m \quad (7)$$

is the gamma-kill survival probability. The gamma-kill dose fraction is

$$D_\gamma = (1 - P)D \quad (8)$$

where D is the absorbed dose. The fraction of mutations produced by ions is evaluated as

$$M = 1 - \Pi_i \Pi_\gamma \quad (9)$$

when mutation response parameters are applied. The track structure model is applied next to estimate *C. elegans* mutation rates after exposure to charged particles.

Response Parameters

Testing procedures for several types of mutations in *C. elegans* have been developed (refs. 3 and 10)




Figure 1. Percentage of mutations in *C. elegans* versus gamma-ray dose. Data from references 3 and 10.

over the last decade. Now, mutation response parameters for recessive lethal mutations are being considered through fits to the data of Nelson et al. (ref. 3) based on results of ion beam experiments in the Lawrence Berkeley Laboratory Bevelac 0-degree Beam Spectrometer. The response to gamma rays has also been measured (refs. 3 and 10) from about 5 to 30 Gy. A linear fit to the gamma-ray data could be used; however, we obtain a good representation by using equation (7) with $m = 2$ and $D_0 = 68$ Gy. Figure 1 shows that the fit is consistent with the ion data. The plateau value of the action cross section for recessive lethal mutations is near $0.2 \mu\text{m}^2$ from the data presented in reference 3. We do not consider track-width effects too cumbersome for use in GCR transport codes at this time. We used the parameterization of equation (2) with $\sigma_0 = 0.25 \mu\text{m}^2$ and $\kappa = 700$ to compare the track model prediction with measured mutation rates. Our findings are shown in figures 2 to 4. The quality of the fit is quite good for the overall data set, as shown on a linear scale with all data reproduced to within a factor of 2 or better. In figure 4, we included target fragment effects extracted from reference 9 for the low linear energy transfer (LET) ions. At high fluence levels, target fragments have minimal effect, but will dominate the effect of low LET ions at lower fluences (ref. 9). Several other ions considered in reference 3 were closer to the Bragg peak than the plateau region and were not considered because fragmentation and other effects complicate the comparisons. Table I is a summary of the response parameters for *C. elegans* mutations. In figure 5, the resulting action cross section versus

Table I. Response Parameters for Recessive Lethal Mutations

	m	D_0 , Gy	σ_0 , cm ²	κ
<i>Caenorhabditis elegans</i>	2	68	2.5×10^{-9}	700




Figure 2. Percentage of mutations in *C. elegans* versus ion fluence for several ions. Data from Nelson et al. (ref. 3).




Figure 4. Percentage of mutations in *C. elegans* versus ion fluence for ions with low LET. Data from Nelson et al. (ref. 3). Solid line includes target fragmentation effects.




Figure 3. Percentage of mutations in *C. elegans* versus ion fluence for several ions. Data from Nelson et al. (ref. 3).




Figure 5. Action cross section versus LET for several ions.

LET is shown for several ions. The cross sections extend from the stopping region to energies above 100 GeV/nucleon. The radial dose distribution of

delta rays used in the track structure model does not reflect relativistic effects that may be important at the highest energies. In reference 11, Fermi has

suggested that relativistic effects may cancel such that a nonrelativistic calculation is accurate. The plateau in the cross section at the highest LET for each ion could be corrected for track width and thin-down effects by using equation (1). The cross section obtained there indicates a target size much smaller than that typical of cellular inactivation, approaching the target sizes in cell transformation and mutation studies (refs. 12 and 13).

Discussion

Solar maximum was achieved in 1990 and the next solar minimum is expected in about 1996; that solar minimum should approach the high fluxes of 1977. Because accurate solar modulation models are not available for transport codes, we present predictions based on 1990 and 1977 galactic cosmic radiation (GCR) spectra and actual measurements (ref. 14). The scaling assumed for the less abundant elements is taken from the Naval Research Laboratory model (ref. 15). Recent measurements (ref. 16) on the Space Transportation System 40 (STS-40) indicated that transport code predictions with the 1990 spectrum were about half those of the measurements. Several factors could lead to this disagreement: solar modulation of the GCR spectrum, the magnetic cutoffs used in the predictions, meson production (which is not included in the transport codes), and the nuclear fragmentation parameters used for heavy-ion transport. Predictions using solar minimum and maximum spectra are expected to bound the actual fluxes on STS-42. In figure 6, we show the unattenuated GCR spectrum for an orbit of 57° by 163 n.mi. for the solar minimum and maximum.




Figure 6. Unattenuated integral flux spectrum versus LET for solar minimum and maximum in an orbit of 57° by 163 n.mi.

The International Microgravity Laboratory 1 (IML-1) experiment was placed inside the Spacelab for its early 1992 mission. Therefore, we are particularly interested in assessing the fractions of mutagenesis in space behind the STS-42 shielding. Figure 7 shows the expected energy spectrum of trapped protons based on the Goddard Space Flight Center energy-space-particle model AP-8. The plot reflects our finding that very few high-energy protons had sufficient range to penetrate the STS-42 shielding.




Figure 7. Trapped proton energy spectrum for STS-42.

The corresponding shielding distribution is shown in figure 8. Also shown is the distribution for dosimeter location 2, which was in the least shielded location on STS-42. For the Spacelab, a significant fraction of the solid angle is above 20 g/cm² aluminum, and we expect that many heavy ions were fragmented with the large buildup of proton and neutron secondaries near *Caenorhabditis elegans* (*C. elegans*). The action cross section is plotted against range in figure 9 to illustrate how beam attenuation with shielding affects the biological damage expected for various ions. The low and medium charged ions should be drastically affected; the high *Z* particles with significant range should be only slightly changed in cross section; however, the use of equation (2) in the track-width regime will lead to an underestimate of cross section. The attenuated linear energy transfer (LET) values for the solar maximum and minimum spectra are shown in figures 10 and 11, respectively.

The Langley Research Center cosmic-ray transport code (ref. 17) was used to predict the resulting




Figure 8. Shielding distribution for Spacelab and dosimeter location 2.




Figure 9. Action cross section versus range for several ions.

mutation rates (ref. 7) for the solar maximum and minimum spectra. Mutation rates were obtained as a function of aluminum shield thickness and were averaged over the shield distributions in figure 8 and over the predictions in table II. It is clear from the table II predictions that few mutations are expected in even a large *C. elegans* population because of the short exposure period. Also shown in table II are predicted values for relative biological effectiveness (RBE) with energetic photons the reference radiation. The high values indicate that our reference photon radiation did not produce indicative numbers of mutations at low doses compared with the low exposure. The exposure could be improved, however, if we optimize

the exposure geometry (if possible within Spacelab constraints) and greatly increase the exposure period. The substantial Spacelab shielding promotes a trapped proton-induced mutation rate of less than 10^{-7} for the mission that is much smaller than the GCR contribution.

Part of the *C. elegans* population was stored at 2°C in Spacelab so that researchers could identify tracks through the surrounding plastic track detector (CR-39). The temperature differential may have altered repair mechanisms in that population; these mechanisms can be interpreted by a repair model that incorporates the track structure model discussed in reference 18.




Figure 10. Solar maximum integral flux spectrum versus LET at Spacelab and dosimeter location 2.




Figure 11. Solar minimum integral flux spectrum versus LET at Spacelab and dosimeter location 2.

Table II. Mutation Rates, RBE Values, and Dose Predictions for STS-42

Location	Dose, mrad		Mutation factor		RBE	
	Min	Max	Min	Max	Min	Max
Spacelab	57	34	0.77×10^{-6}	0.47×10^{-6}	102	131
Dosimeter 2	54	31	$.91 \times 10^{-6}$	$.48 \times 10^{-6}$	107	139

NASA Langley Research Center
Hampton, VA 23681-0001
September 30, 1992

References

1. Coohill, Thomas; Marshall, Tamara; Schubert, Wayne; and Nelson, Gregory: Ultraviolet Mutagenesis of Radiation-Sensitive (rad) Mutants of the Nematode *Caenorhabditis Elegans*. *Mutation Res.*, vol. 209, 1988, pp. 99-106.
2. Nelson, Gregory A.: *Radiation Biology of HZE Particles*. IAF-90-548, Oct. 1990.
3. Nelson, Gregory A.; Schubert, Wayne W.; Marshall, Tamara M.; Benton, Eric R.; and Benton, Eugene V.: Radiation Effects in *Caenorhabditis Elegans*, Mutagenesis by High and Low LET Ionizing Radiation. *Mutation Res.*, vol. 212, 1989, pp. 181-192.
4. Katz, R.; Ackerson, B.; Homayoonfar, M.; and Sharma, S. C.: Inactivation of Cells by Heavy Ion Bombardment. *Radiat. Res.*, vol. 47, 1971, pp. 402-425.
5. Katz, Robert; Sharma, S. C.; and Homayoonfar, M.: The Structure of Particle Tracks. *Topics in Radiation Dosimetry, Supplement 1*, F. H. Attix, ed., Academic Press, Inc., c.1972, pp. 317-383.
6. Waligorski, M. P. R.; Sinclair, G. L.; and Katz, R.: Radiosensitivity Parameters for Neoplastic Transformations in C3H10T1/2 Cells. *Radiat. Res.*, vol. 111, 1987, pp. 424-437.
7. Cucinotta, Francis A.; Katz, Robert; Wilson, John W.; Townsend, Lawrence W.; Nealy, John E.; and Shinn, Judy L.: *Cellular Track Model of Biological Damage to Mammalian Cell Cultures From Galactic Cosmic Rays*. NASA TP-3055, 1991.
8. Katz, Robert; and Cucinotta, F. A.: RBE vs. Dose for Low Doses of High-LET Radiations. *Health Phys.*, vol. 60, no. 5, May 1991, pp. 717-718.
9. Cucinotta, Francis A.; Katz, Robert; Wilson, John W.; Townsend, Lawrence W.; Shinn, Judy; and Hajnal, Ferenc: Biological Effectiveness of High-Energy Protons: Target Fragmentation. *Radiat. Res.*, vol. 127, 1991, pp. 130-137.
10. Rosenbluth, Raja E.; Cuddeford, Cheryl; and Baillie, David L.: Mutagenesis in *Caenorhabditis Elegans*—I. A Rapid Eukaryotic Mutagen Test System Using the Reciprocal Translocation, eTI(III;V). *Mutation Res.*, vol. 110, 1983, pp. 39-48.
11. Fermi, Enrico: *Nuclear Physics*, Revised ed. Univ. of Chicago Press, c.1950.
12. Yang, Tracy Chui-Hsu; Craise, Laurie M.; Mei, Man-Tong; and Tobias, Cornelius A.: Neoplastic Cell Transformation by Heavy Charged Particles. *Radiat. Res.*, vol. 104, 1985, pp. S-177-S-187.
13. Thacker, John; Stretch, Albert; and Stephens, Miriam A.: Mutation and Inactivation of Cultured Mammalian Cells Exposed to Beams of Accelerated Heavy Ions. II. Chinese Hamster V79 Cells. *Int. J. Biol.*, vol. 36, no. 2, 1979, pp. 137-148.
14. Badhwar, G. D.; and O'Neill, P. M.: An Improved Model of Galactic Cosmic Radiation for Space Exploration Missions. Johnson Space Center paper presented at the Second International Cosmic Ray Conference (Dublin, Ireland), Aug. 1991.
15. Adams, James H., Jr.: *Cosmic Ray Effects on Microelectronics, Part IV*. NRL Memo. Rep. 5901, U.S. Navy, Dec. 31, 1986. (Available from DTIC as AD A176 611.)
16. Badhwar, Gautam D.; Braby, Leslie A.; Cucinotta, Frank A.; and Atwell, William: Dose Rate, Dose-Equivalent Rate, and Quality Factor in SLS-1. *Int. J. Radiat. Appl. Instrum., Part D: Nucl. Tracks Radiat. Meas.*, vol. 20, no. 3, 1992, pp. 447-451.
17. Wilson, John W.; Townsend, Lawrence W.; Schimmerling, Walter; Khandelwal, Govind S.; Khan, Ferdous; Nealy, John E.; Cucinotta, Francis A.; Simonsen, Lisa C.; Shinn, Judy L.; and Norbury, John W.: *Transport Methods and Interactions for Space Radiations*. NASA RP-1257, 1991.
18. Wilson, John W.; and Cucinotta, Francis A.: *Cellular Repair/Misrepair Track Model*. NASA TP-3124, 1991.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY(Leave blank)		2. REPORT DATE November 1992		3. REPORT TYPE AND DATES COVERED Technical Memorandum
4. TITLE AND SUBTITLE Katz Model Prediction of <i>Caenorhabditis elegans</i> Mutagenesis on STS-42			5. FUNDING NUMBERS WU 199-04-16-11	
6. AUTHOR(S) Francis A. Cucinotta, John W. Wilson, Robert Katz, and Gautam Badhwar				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) NASA Langley Research Center Hampton, VA 23681-0001			8. PERFORMING ORGANIZATION REPORT NUMBER L-17090	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, DC 20546-0001			10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA TM-4383	
11. SUPPLEMENTARY NOTES Cucinotta and Wilson: Langley Research Center, Hampton, VA; Katz: University of Nebraska, Lincoln, NE; Badhwar: Lyndon B. Johnson Space Center, Houston, TX.				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified Unlimited Subject Category 52			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Response parameters that describe the production of recessive lethal mutations in <i>C. elegans</i> from ionizing radiation are obtained with the Katz track structure model. The authors used models of the space radiation environment and radiation transport to predict and discuss mutation rates for <i>C. elegans</i> on the IML-1 experiment aboard STS-42.				
14. SUBJECT TERMS Radiation mutagenesis; Track structure; International Microgravity Laboratory; Cosmic rays			15. NUMBER OF PAGES 7	
			16. PRICE CODE A02	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	