

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Insecta Mundi

Center for Systematic Entomology, Gainesville,
Florida

June 1997

Nomenclatural changes, reinstatements, new combinations, and new synonymies among American Cerambycids (Coleoptera)

Gerard Luc Tavakilian
French Guiana

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>



Part of the [Entomology Commons](#)

Tavakilian, Gerard Luc, "Nomenclatural changes, reinstatements, new combinations, and new synonymies among American Cerambycids (Coleoptera)" (1997). *Insecta Mundi*. 263.

<https://digitalcommons.unl.edu/insectamundi/263>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Nomenclatural changes, reinstatements, new combinations, and new synonymies among American Cerambycids (Coleoptera)

Gérard Luc Tavakilian

Entomologie Forestière

ORSTOM B.P. 165

97323 CAYENNE cedex, FRENCH GUIANA

Abstract: New synonyms: *Protorma scabrosa* Waterhouse, 1880 and *Protorma recurvatum* Williams, 1829 = *Strongylaspis costifer* Thomson, 1877; *Callophisma ruficollis* Bates, 1870 = *Eriphus croceicollis* White, 1855; *Odontocera cinctiventris* Bates, 1870 = *Odontocera simplex* White, 1855; *Tomopterus aurantiacosignatus* Zajciw, 1969 = *Tomopterus similis* Fisher, 1930; *Cerambyx spectabilis* Voet, 1778 = *Cerambyx velutinus* Fabricius, 1775; *Chion ochraceus* Bates, 1885 = *Lamia spinifera* Fabricius, 1792; *Lissonotus shepherdii* Pascoe, 1859 = *Callidium equestre* Fabricius, 1787; *Cerambyx pulverulentus* Olivier, 1790 = *Cerambyx farinosus* Linné, 1758; *Lamia crypta* Say, 1832 = *Saperda annulata* Fabricius, 1801; *Ptericoptus forsteri* Tippmann, 1960 = *Bisaltus buquetii* Thomson, 1868; *Ataxia flaviceps* Breuning, 1942 = *Stenocorus obscurus* Fabricius, 1801; *Cryptocranium gounellei* Breuning, 1980 = *Cryptocranium cazieri* Lane, 1958; *Epectasis grossepunctata* Breuning, 1942 = *Saperda juncea* Newman, 1840; *Paraclytemnestra gigantea* Breuning, 1974 = *Jamesia lineata* Fisher, 1926; *Hippopsis tremata* Galileo and Martins, 1988 = *Hippopsis meinerti* Aurivillius, 1900; *Lamia punctata* Fabricius, 1792 = *Cerambyx (Lamia) Daviesii* Swederus, 1787; *Leiopus (Oedopeza) pogonocheroides* Audinet-Serville, 1835 = *Lamia ocellator* Fabricius, 1801; *Colobothea velutina* Bates, 1865 = *Lamia macularis* Olivier, 1792; *Xylergates dorotheae* Gilmour, 1962 = *Xylergates elaineae* Gilmour, 1962; *Colobothea velutina* Bates, 1865 = *Lamia macularis* Olivier, 1792; *Pretilia telephoroides* Bates, 1866 = *Saperda tuberculata* Fabricius, 1801; *Guyanestola macrophthalma* Breuning, 1961 = *Estoloides (Estoloides) angustifrons* Breuning, 1943; *Drycothaea marmorata* Martins and Galileo, 1990 = *Guyanestola brasiliensis* Breuning, 1974.

The following species are reinstated: *Strongylaspis costifer* Thomson, 1877; *Oxymerus lineatus* Dupont, 1836; *Hebestola operaria* Erichson, 1848; *Hesycha lateralis* Thomson, 1868.

The genus *Hephialtes* Thomson, 1864 is reinstated.

New combinations: *Hephialtes ruber* (Thunberg, 1822); *Protorma costifera* (Thomson, 1877); *Knalliana cincta spinifera* (Fabricius, 1801) comb. nov.; *Ataxia operaria* (Erichson, 1848) comb. nov.; *Paraclytemnestra lineata* (Fisher, 1926) comb. nov.; *Oedopeza ocellator* (Fabricius, 1801) comb. nov.; *Toronaus incisus* (Bates, 1864) comb. nov.; *Pretilia tuberculata* (Fabricius, 1801) comb. nov. *Drycothaea turrialbae* (Breuning, 1943) comb. nov. *Drycothaea truncatipennis* new name for *Estola stictica* Breuning, 1942 (nec Bates, 1881)

Some host-plants are given.

Introduction

During my various visits through several institutions sheltering historical collections, I noticed some unknown synonymies which are interesting to publish. The acronyms used are:

MNHN for the Muséum d'Histoire Naturelle de Paris,
BMNH for the British Natural History Museum in
London,

AMNH for the American Museum of Natural History in
New York,

USNM for the United States National Museum in Wash-
ington D.C. (Smithsonian Institution),

ICCM for the Carnegie Museum in Pittsburgh, Pennsylv-
ania,

MNHU for the Museum für Naturkunde der Humboldt
Universität in Berlin,

ZMK for the Zoologisk Museum in Copenhagen,

MNRJ for the Museu Nacional da Quinta da Boa Vista
in Rio de Janeiro,

ZSMA for the Zoologische Staatssammlung des Bay-
erischen Staates in Munich,

NRS for the Naturhistoriska Riksmuseet in Stockholm.

Concerning host plant records, the scientific name is followed by the herbarium collection refer-

ence number sampled by a botanist; several parts are distributed throughout main botanical institutions of the world.

1. Reinstated species:

Subfamily Prioninae

Tribe Macrotomini

***Protorma costifera* (Thomson, 1877), reinstated and new combination.**

Strongylaspis costifer Thomson, 1877: 275

Protorma scabrosa Waterhouse, 1880: 289, **new syno-
nymy**

Stenodontes (Protorma) scabrosus; Lameere, 1903: 215

Protorma recurvatum Williams, 1929: 143, pl. V, fig. 1,
new synonymy

Protorma recurvata Blackwelder, 1946: 552 [Catalog]

Lameere, without having seen Thomson's material, simply synonymized *Strongylaspis costifer* with *Strongylaspis corticarius* (Erichson, 1848). All the following authors (Villiers, 1980: 143 and

others) accepted Lameere's judgement. Fortunately, Thomson's type of *Strongylaspis costifer* is visible in the MNHN, and in fact, belongs to the genus *Protorma*. Later on, I saw Williams' type in ICCM of *Protorma recurvatum* and came to the conclusion that Williams species is just a synonym of Thomson's species. Williams wrote "It was carefully compared by me with the type of *Protorma scabrosa* Waterhouse, from which it is quite distinct." In fact, the description of *Protorma recurvatum* was based on a single male specimen, and the type of *Protorma scabrosa* in the BMNH is a female. The three described taxa belong to the same species, and the valid name is *Protorma costifera* (Thomson, 1877).

Subfamily Cerambycinae

Tribe Trachyderini

Oxymerus lineatus Dupont, 1838 reinstated

Oxymerus lineatus Dupont, 1838: 41, pl. 211, fig. 1 [Brazil]

Oxymerus approximatus Dupont, 1838: 44, pl. 212, fig. 2 [Cayenne]

Oxymerus elongatus Dupont, 1838, 1838: 45, pl. 213, fig. 1 [Cayenne]

Oxymerus distinguendus Dupont, 1838: 48, pl. 214, fig. 2 [Demerary]

Oxymerus aculeatus subsp. *lineatus* Hüdepohl, 1979: 20, pl. I, 1(m)

Oxymerus aculeatus var. *lineatus* Rémy, 1988: 130

From November 1982 to January 1983 the species was particularly abundant around Kourou where I saw more than a thousand. I definitely do not agree with Hüdepohl synonymies and status changing (Hüdepohl, 1979: 20). The correct synonymy, after checking the types found in the MNHN, is listed above.

Subfamily Lamiinae

Tribe Pteropliini

Fabricius described (1775: 180) a *Stenocorus annulatus* from "America meridionali" belonging to the Banks collection (today in the BMNH). Fabricius later on (1801: 310) mentioned the synonymy with *Cerambyx annulatus* Olivier, 1792 and *Cerambyx hirtipes* Degeer, 1775 (belonging to the actual genus *Colobothea* Lepeletier and Audinet-Serville, 1825). In 1792: 314 he described a *Saperda lineata* from America meridionali. The two synonyms in the ZMK belong to Degeer's species, *Cerambyx hirtipes*.

In 1801: 326, he described *Saperda annulata* and gave the short following description in latin: thorace rotundato, subspinoso, antennarum articulis basi albis, which agrees with the type in Fabricius' collection in the ZMK (conspecific with Say's *Lamia crypta*); both the locality (America meridionali) and the synonymy (*Saperda lineata* 1792: 314) were wrongly associated with the above description. The confusion was obviously caused by an error in the printing of page 326 in Fabricius' Systema Eleutheratorum. It is clear that between the *Saperda nigricornis* (n° 47) and the *Saperda tristis* (n° 51) the synonymies were wrongly skipped. Bates in the supplement of Biologia Centrali Americana (1885: 347) wrote: *Ataxia Crypta* ... Said by Erichson to be the *Saperda lineata* of Fabricius (Ent. Syst. ii p. 314), and so registered in the Munich catalogue, but it does not agree with the description, which refers also, as Fabricius states, to a South-American (Brazilian ?) insect in Lunds collection. Erichson might have seen the type and Bates read the wrong description (Fabricius, 1792: 314). The name *Saperda annulata* being used several times by Fabricius, we decide to conserve Say's species name in order to avoid more confusion. The complete synonymy of *Ataxia crypta* is the following:

Ataxia crypta (Say, 1832)

Saperda annulata Fabricius, 1801(nec Fabricius, 1792): 326, **new synonymy**

Lamia crypta Say, 1832: 5

Amniscus ? cryptus Haldeman, 1847: 47

Stenosoma crypta LeConte, 1873: 302

Ataxia crypta LeConte, 1873: 344

Ataxia sordida Haldeman, 1847: 56

Stenosoma sordida LeConte, 1852: 158

Consequently *Hebestola operaria* Erichson, 1848 synonymized with *Saperda annulata* Fabricius, *Parysatis nigratarsis* Thomson, 1868b, and *Parysatis flavescens* Bates, 1880 becomes a valid taxon and we can write the following synonymy:

Ataxia operaria (Erichson, 1848), reinstated

Hebestola operaria Erichson, 1848: 574

Parysatis nigratarsis Thomson, 1868b: 120

Esthlogena operaria Lacordaire, 1872: 600

Ataxia lineata Gemminger and Harold, 1873: 3100 [Catalog]

Parysatis flavescens Bates, 1880: 112

Tribe Onciderini

Lampedusa lateralis (Thomson, 1868) reinstated

Hesycha lateralis Thomson, 1868a: 63 [Cayenne]
Lampedusa obliquator Dillon and Dillon (*nec* Fabricius, 1801), 1945: 114, pl. V, fig. 10

***Lampedusa obliquator* (Fabricius, 1801)**

Lamia obliquator Fabricius, 1801: 303
Hypsioma obliquator Thomson, 1868a: 50
Plerodia obliquator Aurivillius, 1923: 342 [Catalog]

The type of *Hesycha lateralis* Thomson, 1868 is different from the type of *Lamia obliquator* Fabricius, 1801. Dillon and Dillon wrongly interpreted Fabricius' species and consequently *Lampedusa lateralis* (Thomson, 1868) is reinstated. Both species were encountered in French Guiana and belong to the genus *Lampedusa* as described by Dillon and Dillon, 1945: 112. Following the nomenclatural code, *Lamia obliquator* remains the type species of the genus *Lampedusa* Dillon and Dillon, 1945.

2. Reinstated genus:

Subfamily Prioninae

Tribe Callipogonini

The genus *Hephialtes* Thomson, 1864 was created for *Hephialtes tricastatus* Thomson, 1864: 286 based on a female specimen from Brazil. A year later he described the male as *badium*. Both are synonyms of *Trachyderes ruber* Thunberg, 1822. Lameere (1904: 28) synonymized the genus with *Anacanthus* Audinet-Serville, 1832 which he considered as a subgenus of *Stictosomus* Audinet-Serville, 1832. Whilst reading the description of the genus *Stictosomus* (page 153) we note incompatible features with Thunberg's species "Corselet sans crénelures" and "mandibules allongées, étroites, aiguës, denticulées intérieurement". Considering the description of *Anacanthus* (page 165 of the same Audinet-Serville's paper) a name which means in Greek "without spines" we find again incompatible features (corselet mutique ... arrondi...). Neither the genus nor the subgenus retained by Lameere is correct.

Though the type species of the 3 concerned genera (*Stictosomus semicostatus* Audinet-Serville, 1832, *Anacanthus costatus* Audinet-Serville, 1832 and *Hephialtes tricastatus* Thomson, 1864) belong obviously to the same natural subdivision of the tribe Callipogonini, I think it is unreasonable to synonymize them under the same genus.

***Hephialtes* Thomson, 1864 reinstated**

Type species: *Hephialtes tricastatus* Thomson, 1864 (= *Trachyderes ruber* Thunberg, 1822) [Brazil] by original designation.

Hephialtes Thomson, 1864: 285

***Hephialtes ruber* (Thunberg, 1822), comb. nov.**

Trachyderes ruber Thunberg, 1822: 305
Orthosoma badium Dejean (nomen nudum), 1837: 342
Hephialtes tricastatus Thomson, 1864: 286
Hephialtes badium Thomson, 1865: 577
Hephialtes sulcatus Lacordaire, 1869: 147
Hephialtes badius Gemminger and Harold, 1872: 2774 [Catalog]
Hephialtes tricastatus Gahan, 1895: 84
Stictosomus (Anacanthus) tricastatus Lameere, 1904: 28
Stictosomus ruber Lameere, 1912: 164
Stictosomus (Anacanthus) ruber Lameere, 1913: 38 [Catalog]
Anacanthus ruber Villiers, 1980: 152, fig. 20

3. New synonyms and consequently new combinations:

Subfamily Cerambycinae

Tribe Rhinotragini

***Odontocera simplex* White, 1855**

Odontocera simplex White, 1855: 189
Odontocera cinctiventris Bates, 1870a: 321, new synonymy

White's type, in the BMNH, is represented by a male specimen and Bates' type, in the MNHN, a female, although the latter was doubtful concerning the gender of his *O. cinctiventris*. I personally obtained many individuals of both sexes by rearing them on their host plant. I was already alerted by the elytral similarity of the two supposed species before having them bred. They emerged from three different species of trees all belonging to the family of Papilionaceae: *Taralea oppositifolia* Aublet [Denis Loubry 1611]; *Dipteryx punctata* (Black) Amshoff [Denis Loubry 1797]; *Monopteryx inpa* Rodrigues [Denis Loubry 1739].

Tribe Callichromatini

Callichroma velutinum (Fabricius, 1775)
Cerambyx velutinus Fabricius, 1775: 167
Cerambyx spectabilis Voet, 1778: 12, pl. X, fig. 40, **new synonymy**
Cerambyx velutinus Olivier, 1795: 24, (67) pl. VI, fig. 41
Callichroma velutina Serville, 1833: 557
Callichroma velutinum White, 1853: 164

Callichroma velutinus Lacordaire, 1869: 16
Callichroma porphyrogenitum Bates, 1870: 333
Callichroma (Callichroma) velutinum Schmidt, 1924: 321

Considering that the illustration is sufficient, and that the locality is "Indes Occidentales", this leaves no doubt about the identity of this taxon. The epithet "*spectabilis*" still remains in all catalogs for obscure reasons.

Tribe Lissonotini

Lissonotus equestris (Fabricius, 1787)

Cerambyx biserratus Voet, 1778: 29, pl. XXV, fig. 137 [America]
Callidium equestre Fabricius, 1787: 153 [Cajennae]
Cerambyx unidentatus Olivier, 1795: 20, (n° 67) pl. XIX, fig. 145
Cerambyx cinctus Panzer in Voet, 1798: 2, pl. XXV, fig. 136
Cerambyx cingulatus Panzer in Voet, 1798: 2, pl. XXV, fig. 137
Lissonotus cinctus Dalman in Schönherr, 1817: 364 [Cayenna]
Lissonotus unidentatus Dalman in Schönherr, 1817: 364
Lissonotus equestris Dejean, 1821: 109
Lissonotus shepherdii Pascoe, 1859: 16 [Pará], **new synonymy**
Lissonotus shepherdii Blackwelder, 1946: 592 [Brazil] Catalog

I found in the MNHN at the Library of the French Entomological Society an example of Voet's book where the first pages are missing but *Cerambyx biserratus* was already described in 1778. Since the name *equestris* has been used for over a century I think it is better to conserve the well known name. On the other hand, *Lissonotus shepherdii* is exactly similar to Fabricius' type. To summarize my observation, the most common form with the red band complete seems to be a variety of *L. equestris*. I obtained this abundant species from several legume trees and vines. Both forms emerged from the same logs (interrupted or complete red band). Though the type locality is Cayenne the interrupted form is rarer in French Guiana.

Observed host plants: *Hymenaea courbaril* Linné (Caesalpiniaceae) [Sabatier and Prévost 3490]; *Bauhinia guianensis* Aublet (Caesalpiniaceae) [Scott Mori and al. 23641, 23678]; *Macrolobium* sp. (Caesalpiniaceae) [Scott Mori and al. 23467, 23409].

Tribe Trachyderini

Subtribe Bothriospilina

The identity of *Lamia spinifera* Fabricius, 1792 is not yet resolved. Martins and Moure (1973: 79), giving an illustration of the species, came to the conclusion that it might certainly be an Oriental or Ethiopian species, maybe belonging to the genus *Plocaederus* Thomson.

Knulliana cincta spinifera (Fabricius, 1792) new combination

Lamia spinifera Fabricius, 1792: 275; Fabricius, 1801: 292; Aurivillius, 1923: 606 [Catalog]; Blackwelder, 1946: 627 [Catalog] *incertae sedis*; Zimsen, 1964: 169 [Types]; Martins and Moure, 1973: 79 figs 1-2 (male).
Chion ochraceus Bates, 1885 243, **new synonymy**
Chion cinctus var. *ochraceus* Schaeffer, 1908: 331
Chion cinctum v. *ochraceum* Blackwelder, 1946: 562 [Catalog]
Cerasphorus cinctum ochraceum Cazier and Lacey, 1952: 10 fig. 1
Knulliana cincta ochracea Linsley, 1962: 111 fig. 35 (distribution map)

Subtribe Ancylocerina

Callancyla croceicollis (White, 1855)

Eriphus croceicollis White, 1855: 292; Bates, 1870b: 429.
Callopisma ruficollis Bates, 1870b: 419, **new synonymy**
Ancylocera ruficollis Gemminger and Harold, 1872: 2959 [Catalog]
Callancyla ruficollis Aurivillius, 1912: 446 [Catalog]

Bates wrote (1870: 429), concerning *Eriphus croceicollis* White, 1855, "Although taken by me, I do not find the species among my own reserved collection of Amazonian Longicorns". He would never have been able to guess White's misplacement whose type (BMNH) is a female. Bates, though uncertain about the sex of his *Callopisma ruficollis* supposed the type (MNHN) to be a male.

Subfamily Lamiinae

Tribe Lamiini

Taeniotes farinosus (Linné, 1758)

Cerambyx farinosus Linné, 1758: 390
Cerambyx farinosus DeGeer, 1775: 108, pl. XIV, fig. 1 [Suriname]
Cerambyx pulverulentus Olivier, 1790: 302, **new synonymy**

Cerambyx surinamensis maculosus Panzer, 1794: 15, pl. V, fig. 8 [Suriname]
Cerambix pulverulentus Olivier, 1795: 50, (n° 67) pl. VII, fig. 46 b
Stenochorus pulverulentus Schönherr, 1817: 407 [Catalogue, Suriname]
Monochamus pulverulentus Dejean, 1821: 106
Taeniotus pulverulentus Audinet-Serville, 1835: 91 [Cayenne]
Monochamus (Taeniotus) farinosus Castelnau, 1840: 479 [Brazil]
Monochamus (Taeniotus) pulverulentus Castelnau, 1840: 479 [Brazil]
Taeniotus farinosus Bates, 1865b: 110 [Brazil: Amazon, Tapajos]
Taeniotus guttularis Schwarzer, 1929: 364, fig. 16-17
Taeniotus pulverulentus m. *guttularis* Breuning, 1943b: 248
Taeniotus pulvurentus Bosq, 1943: 109
Taeniotus farinosus Dillon and Dillon, 1941 pl. I, fig. 6
Taeniotus farinosa Blackwelder, 1946: 594 [Catalog]
Taeniotus pulverulenta Blackwelder, 1946: 594 [Catalog]

Olivier's drawing representing *Cerambyx pulverulentus* closely resembles the species commonly collected in Cayenne. Linné's *farinosus*, which is neither in the London Linnean Society nor in Gustave-Adolphe's collection in Uppsala, has in fact no type. Linné mentioned after his laconic description the corresponding illustration in Sybille Merian's book published in 1719 (table 24, f. *infima*) and another reference published by Degeer before the tenth edition of Linné's *Systema Naturae*. The iconotype belongs to the prelinnean literature; I ignore what is to be done in this case. I consulted this very rare book with colored plates and I am certain that *C. pulverulentus* is a synonym of *C. farinosus* and their localities (Suriname and Cayenne) confirm the synonymy. The species is very common on the introduced breadfruit tree *Artocarpus altilis* (S. Parkinson) Fosberg (Moraceae). I do not understand why these two taxa were considered as different by all authors, except for Bates who had already reported the synonymy (1865b: 110).

Type Apomecynini

Bisaltes buqueti Thomson, 1868

Bisaltes buqueti Thomson, 1868b: 111 [Cayenne]
Bisaltes buqueti Gemminger and Harold, 1873: 3102 [Catalog]

Bisaltes sp. Bates, 1885: 346, pl. IX, fig. 21a
Bisaltes batesi Aurivillius, 1923: 609 [Catalog]
Ptericoptus forsteri Tippmann, 1960: 152, pl. IX, fig. 21a, new synonymy
Bisaltes (Bisaltes) buqueti Breuning, 1960: 177 [Catalog]
Bisaltes (s.str.) forsteri Breuning, 1971: 284

Tippmann's type is in the USNM and the type of Thomson in the MNHN. The species was reared on a Convolvulaceae vine belonging to the genus *Maripa* [Scott Mori *et al.* 23592].

Tethystola brasiliensis Breuning, 1940

Tethystola brasiliensis Breuning, 1940: 42
Tethystola flavoapicalis Breuning, 1942: 141, new synonymy

Both types are at the MNHN and were collected in the Brazilian state of Pernambuco. The type of *T. flavoapicalis* is a big specimen.

Tribe Pteropliini

Ataxia obscura (Fabricius, 1801)

Stenocorus obscurus Fabricius, 1801: 307 [America meridionali]
Esthlogena sulcata Bates, 1866: 290
Parysatis collaris Thomson, 1868: 119
Parysatis sulcata Aurivillius, 1922: 293 (Catalog)
Parysatis obscura Aurivillius, 1922: 292 (Catalog)
Ataxia flaviceps Breuning, 1942: 137, new synonymy
Ataxia obscura Breuning, 1961: 53, fig. 7

Cryptocranium cazieri Lane, 1958

Cryptocranium cazieri Lane, 1958: 6, fig. 2
Ornatodesisa pulchra Breuning, 1961: 19
Cryptocranium gounellei Breuning, 1980: 68, new synonymy

Lane's type (a female from Peru) belongs to the AMNH and Breuning's type (a male with flattened legs from the Brazilian Goiás) belongs to the MNHN. The recent synonymy published by Monné and Giesbert (1992: 252) for a specimen already described by Breuning from the Brazilian state of Pará, and the presence of this species in French Guiana, lead us to consider that this very rare species is widely distributed in northern South America.

Epectasis juncea (Newman, 1840)

Saperda juncea Newman, 1840: 13
Epectasis attenuata Bates, 1866: 294
Epectasis grossepunctata Breuning, 1942: 175, new synonymy

Epectasis juncea m. attenuata Breuning, 1961c: 25

Breuning's type belongs to the USNM, the type of *E. attenuata* is in the MNHN.

Tribe Onciderini

***Paraclytemnestra lineata* (Fisher, 1926), new combination.**

Jamesia lineata Fisher, 1926: 14

Paraclytemnestra gigantea Breuning, 1974a: 240, **new synonymy**

This incredible species from the Antilles merits its own genus. Breuning's type belongs to the MNHU, though Fisher's type belongs to the USNM.

Tribe Agapanthiini

***Hippopsis meinerti* Aurivillius, 1900**

Hippopsis meinerti Aurivillius, 1900: 416; Aurivillius, 1923: 357 [Catalog]

Hippopsis meinerti Blackwelder, 1946: 606 [Catalog]; Galileo and Martins, 1988a: 185 figs 17-20; Galileo and Martins, 1988b: 205

Hippopsis freyi Breuning, 1955: 661

Hippopsis (Hippopsis) lemniscata m. meinerti Breuning, 1961b: 197 [Catalog]; Breuning, 1962: 10

Hippopsis (Hippopsis) freyi Breuning, 1961b: 197 [Catalog]; 1962: 18

Hippopsis (Hippopsis) lemniscata tobagoensis Breuning, 1962: 10

Hippopsis tremata Galileo and Martins, 1988a: 184, figs. 13-16, **new synonymy**

Aurivillius' type, in the NRS is a male, and the abdominal depression is absolutely identical to the drawing given in the paper of Galileo and Martins to characterize their new species *Hippopsis tremata*. I compared Aurivillius' type with my own material of both sexes from French Guiana and identified by Martins as *H. tremata*.

Tribe Acanthoderini

***Acanthoderes daviesi* (Swederus, 1787)**

Cerambyx (Lamia) daviesii Swederus, 1787: 195, pl. VIII, fig. 6

Cerambyx daviesii Gmelin, 1790: 1837

Lamia punctata Fabricius, 1792: 272 [Cajennae], **new synonymy**

Lamia daviesii Schönherr, 1817: 380 [America, Cayenna]

Acanthoderes daviesii Dejean, 1835: 336

Acanthoderes daviesii Audinet-Serville, 1835: 29 [Cayenne]

Acanthoderes swederi White, 1855: 360, pl. IX, fig. 6 [Para]

Acanthoderes daviesi Thomson, 1864: 17

Acanthoderes daviesi var. *swederi* Gemminger and Harold, 1873: 3146 [Catalog]

Acanthoderes daviesi Bodkin, 1919: 268

Acanthoderes (Acanthoderes) daviesi Aurivillius, 1923: 384 [Catalog]

Acanthoderes (Acanthoderes) punctata Aurivillius, 1923: 385 [Catalog]

Acanthoderes (Acanthoderes) daviesi Buck, 1959: 604

The common and widely distributed epiphyte *Clusia grandiflora* Splitgerber (Clusiaceae) [Denis Loubry 1827] breeds *A. daviesi*.

Tribe Acanthocinini

***Oedopeza ocellator* (Fabricius, 1801), new combination**

Lamia ocellator Fabricius, 1801: 287 [America meridionali]

Leipus (Oedopeza) pogonocheroides Audinet-Serville, 1835: 88, **new synonymy**

Leipus pogonocheroides Castelnau, 1840: 464 [Cayenne]

Oedopeza pogonocheroides White, 1855: 394

Ozineus ocellator Aurivillius, 1923: 398 [Catalog]

I do not understand why this very common species was not recognized by Aurivillius. The type of Fabricius is a male with a characteristic swollen protarsal segment. I obtained this species from 16 different tree species, mainly belonging to the legume group: *Couma guianensis* Aublet (Apocynaceae) [Christian Feuillet 2311]; *Simarouba amara* Aublet (Simaroubaceae) [Christian Feuillet 2375]; *Theobroma cacao* Linné (Sterculiaceae) [Jean-Jacques de Granville det.]; *Swartzia lomatophylla* (Benth.) Pittier (Caesalpiniaceae) [Denis Loubry, 1585]; *Swartzia panacoco* (Aublet) Cowan (Caesalpiniaceae) [Denis Loubry, 1804]; *Machaerium* sp. (Papilionaceae) [Denis Loubry, 1815]; *Taralea oppositifolia* Aublet (Papilionaceae) [Denis Loubry, 1611]; *Abarema curvicaarpa* (Irwin) Barneby and Grimes (Mimosaceae) [Grimes and Sabatier 3312]; *Abarema jupunba* (Willdenow) Barneby and Grimes (Mimosaceae) [Christian Feuillet 2307, Denis Loubry 1875 and Scott Mori 23688]; *Balizia pedicellaris* (de Candolle) Barneby and Grimes (Mimosaceae) [Christian Feuillet 1341]; *Enterolobium schomburgkii* Benth. (Mimosaceae) [Daniel Sabatier 2367]; *Enterolobium* sp. (Mimosaceae) [Denis Loubry 1763]; *Hydrochorea corymbosa* (L.C.

Richard) Barneby and Grimes (Mimosaceae) [Denis Loubry 1300]; *Inga melinonis* Sagot (Mimosaceae) [Denis Loubry 1780]; *Inga* sp. (Mimosaceae) [Scott Mori 23456]; *Parkia nitida* Miquel (Mimosaceae) [Daniel Sabatier 1511 and Denis Loubry 1564].

***Toronaeus incisus* (Bates, 1864), new combination.**

Nyssodrys incisus Bates, 1864: 155

Nyssodrystes incisus Gilmour, 1965: 599 [Brazil] Catalogue

Stenolis incisus Monné, 1985: 535.

The rearing of this rare species demonstrates clearly its greater affinity with the genus *Toronaeus* (where majority were bound to Burseraceae trees), confirmed by its taxonomic features. It was obtained on a still unidentified Burseraceae [Denis Loubry 1318]

***Xylergates elaineae* Gilmour, 1962**

Astynomus hiliaris: Prudhomme, 1906: 38

Xylergates sp. (*hiliaris* Dejean?) Duffy, 1960: 258, figs. 158-159 (Larva)

Xylergates elaineae Gilmour, 1962: 277, pl. III, fig. 3

Xylergates dorotheae Gilmour, 1962: 279, pl. III, fig. 4, **new synonymy**

Gilmour's two types belong to the American Museum of Natural History in New York. The species commonly attacks fallen trees belonging to the family of Lecythidaceae. I found this species from seven different host plants in French Guiana: *Lecythis poiteaui* O.C. Berg [Scott Mori det. 1995]; *Lecythis holcogyne* (Sandwith) Mori [D. Sabatier and M.F. Prévost 2108]; *Eschweilera sagotiana* Miers [Daniel Sabatier 1762]; *Eschweilera micrantha* (O.C. Berg) Miers [Scott Mori 23479]; *Eschweilera coriacea* (A.P. de Candolle) Martius ex O.C. Berg [Denis Loubry 1846, 1341]; *Eschweilera congestiflora* (R. Benoist) Eyma [Scott Mori 23478]; *Couratari guianensis* Aublet [Daniel Sabatier 1771 and Denis Loubry 1337].

Tribe Colobothini

***Colobothaea macularis* (Olivier, 1792)**

Lamia macularis Olivier, 1792: 474 [Suriname]

Cerambyx macularis Olivier, 1795: (67) 98 pl. XX, fig. 154 [Suriname]

Colobothaea velutina Bates, 1865a: 216, **new synonymy**

Colobothaea macularis: Aurivillius, 1923: 454 [Catalog]

I had the opportunity to consult and to photograph the original paintings of Olivier which served to illustrate his famous "Entomologie..." still conserved in his family, six generations later. These paintings reveal more details than the printed and post-colored plates found in the remaining examples of this very rare book and thus permit the above synonymy. I obtained several adults from *Hydrochorea corymbosa* (L.C. Richard) Barneby and Grimes (Mimosaceae) [Denis Loubry 1300]

Tribe Pretiliini

***Pretilia tuberculata* (Fabricius, 1801), new combination.**

Saperda tuberculata Fabricius, 1801: 331 [America meridionalis]

Pretilia telephoroides Bates, 1866: 302; Lacordaire, 1872: 910; Gilmour, 1962: 125; Martins and Galileo, 1990b: 707 figs 1-11, **new synonymy**

Eupogonius tuberculatus Aurivillius, 1923: 312 [Catalog]; Blackwelder, 1946: 600 [Catalog]

Tribe Calliini

***Drycothaea angustifrons* (Breuning, 1943)**

Estoloides (Estoloides) angustifrons Breuning, 1943a: 56 [Guyane: Maroni]; Breuning, 1974b: 59.

Guyanestola macrophthalma Breuning, 1961a: 248, **new synonymy**

Guyanestola macrophthalma Breuning, 1963: 507 [Catalog]

Drycothaea angustifrons Tavakilian, 1991: 450; Galileo and Martins, 1991: 250

Drycothaea macrophthalma Galileo and Martins, 1991: 251

In 1990, the blocked situation concerning Frey's collection made impossible the examination of Breuning's type during my journey to Munich. The situation is now normal since the collection was transferred from Frey's house in Tutzing to the Staatliches Museum in Munich (ZSMA). I am grateful to Patrick Bleuzen who sent me the picture of *Guyanestola macrophthalma* and checked the claws, confirming that they are appendiculated as Galileo and Martins and I already guessed. This illustrates how uncomfortable can be the position of an entomologist revising a group of which the types are still in private collections.

I obtained this species from *Cecropia palmata* Willdenow (Cecropiaceae) [Denis Loubry 1570].

***Drycothaea brasiliensis* (Breuning, 1974)**

Guyanestola brasiliensis Breuning, 1974b: 49 [Brazil: Amazonas]

Drycothaea marmorata Martins and Galileo, 1990a: 611, fig. 4, **new synonymy**

Drycothaea brasiliensis Galileo and Martins, 1991: 251

Breuning's type is in the ZSMA. The host plant is still *Cecropia palmata* Willdenow (Cecropiaceae) [Denis Loubry 1570].

***Drycothaea truncatipennis* new name for *Estola stictica* Breuning, 1942 (nec Bates, 1881)**

Estola stictica Breuning, 1942: 163

The type belongs to the USNM (ex Tippmann's collection n° 41605) and is a male collected in the Brazilian state of Santa Catarina. The structure of the claws transfers this species from the tribe Desmiphorini to Calliini in the genus *Drycothaea*. Unfortunately another *Drycothaea stictica* Bates, 1881 already exists from Guatemala. I propose the specific name *D. truncatipennis* which fits better with the elytral aspect of that misplaced species.

***Drycothaea turrialbae* (Breuning, 1943), new combination**

Estoloides (Estoloides) turri-albae Breuning, 1943: 56

Estoloides (Estoloides) turrialbae Breuning, 1963: 507

Estoloides turrialbae Chemsak, Linsley and Noguera, 1992: 119 (Catalog)

The type belongs to the USNM (ex Tippmann's collection n° 41596) and is a female labelled as *Parestola turrialbae*! As seen above, this species is transferred from Desmiphorini to the tribe Calliini for the same reason.

References

- Audinet-Serville, J. G.** 1832. Nouvelle classification de la famille des longicornes. Annales de la Société Entomologique de France. Paris 1 (1): 118-201
- Audinet-Serville, J. G.** 1835. Nouvelle classification de la famille des Longicornes 3ème Tribu, Lamiaries, Lamiariae. Annales de la Société Entomologique de France. Paris 4 (1): 5-100.
- Aurivillius, C.** 1900. Verzeichniss der von Dr F. Meinert im Jahre 1891 in Venezuela gesammelten Cerambyciden. Öfversigt af Kongl. Svenska Vetenskaps-Akademiens Föreläsningar 57 (4): 409-421, 1 fig.
- Aurivillius, C.** 1912. Cerambycidae: Cerambycinae. Coleopterorum Catalogus 22 (39): 1-574. W. Junk et S. Schenkling, Berlin.
- Aurivillius, C.** 1923. Lamiinae II. Coleopterorum Catalogus 23 (74): 323-704. W. Junk et S. Schenkling, Berlin.
- Bates, H. W.** 1864. XIX. Contributions to an Insect Fauna of the Amazon Valley. Coleoptera: Longicornes. The Annals and Magazine of Natural History. London 13 (3): 144-164.
- Bates, H. W.** 1865a. XXV. Contributions to an Insect Fauna of the Amazons Valley. Coleoptera: Longicornes. The Annals and Magazine of Natural History. London 15 (3): 213-225.
- Bates, H. W.** 1865b. X. Contributions to an Insect Fauna of the Amazons Valley. Coleoptera: Longicornes. The Annals and Magazine of Natural History. London 16 (3): 101-113.
- Bates, H. W.** 1866. XXXIII. Contributions to an Insect Fauna of the Amazons Valley. Coleoptera: Longicornes. The Annals and Magazine of Natural History. London 17 (3): 288-303.
- Bates, H. W.** 1870a. XVI. Contributions to an Insect Fauna of the Amazon Valley (Coleoptera, Cerambycidae). The Transactions of the Entomological Society of London, 1870 (part III): 243-335.
- Bates, H. W.** 1870b. XIX. Contributions to an Insect Fauna of the Amazon Valley (Coleoptera, Cerambycidae). The Transactions of the Entomological Society of London, 1870 (part IV) December: 391-444.
- Bates, H. W.** 1880. Biologia Centrali-Americana, Insecta, Coleoptera, Cerambycidae, Lamiidae 5: 17-152, pls. III-XI.
- Bates, H. W.** 1885. Biologia Centrali-Americana, Insecta, Coleoptera, Cerambycidae Supplement to Longicornia. 5: 249-436, pls. XVII-XXIV.
- Blackwelder, R. E.** 1946. Checklist of the coleopterous insects of Mexico, Central America, the West Indies and South America. Part 4. Bulletin. United States National Museum 185 (4): 551-763.
- Bodkin, G. E.** 1919. Notes on the Coleoptera of British Guiana. The Entomologist's Monthly Magazine. London 55: 264-272.
- Bosq, J. M.** 1943. Agregados al Catálogo de los Longicornes de la República Argentina. Revista Argentina de Zoogeografía. Buenos Aires 3 (3): 103-112.
- Breuning, S.** 1940. Novae species Cerambycidarum VIII. Folia Zoologica et Hydrobiologica. Riga 10: 37-85.
- Breuning, S.** 1942. Novae species Cerambycidarum XI. Folia Zoologica et Hydrobiologica. Riga 11: 113-175.
- Breuning, S.** 1943 a. Novae species Cerambycidarum XII. Folia Zoologica et Hydrobiologica. Riga 12: 12-66.
- Breuning, S.** 1943 b. Études sur les Lamiaries (Coleop., Cerambycidae). Douzième tribu: Agniini Thomson. Novitates Entomologicae, Paris 3 (89-106): 137-280, figs 1-157.

- Breuning, S.** 1955. Neue Cerambyciden von der Insel Trinidad. *Entomologische Arbeiten aus dem Museum G. Frey, Tutzing bei München* 6 (2): 659-661.
- Breuning, S.** 1960. Catalogue des Lamiaires du Monde (Col. Céramb.). Verlag des Museums G. Frey, Tutzing bei München (3) juli 1960: 109-182.
- Breuning, S.** 1961a. Neue Lamiiden aus dem Museum G. Frey (Col. Ceramb.). *Entomologische Arbeiten aus dem Museum G. Frey, Tutzing bei München* 12(1): 245-250.
- Breuning, S.** 1961b. Catalogue des Lamiaires du Monde (Col. Céramb.). Verlag des Museums G. Frey, Tutzing bei München (4) Jänner 1961: 183-284.
- Breuning, S.** 1961c. Révision des Pteropliini (Col. Cerambycidae). *Pesquisas, Porto Alegre (Zoologia)* 5 (9): 5-61, 7 figs.
- Breuning, S.** 1962. Révision des Agapanthiini Muls. Américains (Col., Cerambycidae). *Pesquisas, Porto Alegre (Zoologia)* 6 (13): 5-48, 9 figs.
- Breuning, S.** 1963. Catalogue des Lamiaires du Monde (Col. Céramb.). Verlag des Museums G. Frey, Tutzing bei München (7) April 1963: 463-555.
- Breuning, S.** 1971. Révision des espèces américaines de la tribu des Apomecynini Lac. (Coleoptera, Cerambycidae). *Entomologische Abhandlungen Staatliches Museum für Tierkunde in Dresden* 37 (3): 209-335, 21 figs.
- Breuning, S.** 1974a. Neue Lamiinae (Coleoptera, Cerambycidae). *Mitteilungen aus dem Zoologischen Museum in Berlin* 50 (2): 237-241.
- Breuning, S.** 1974b. Révision des Rhodopinini Américains (Coleoptera, Cerambycidae). *Studia Entomologica, Petrópolis* 17 (1-4): 1-210, 26 figs.
- Breuning, S.** 1980. Nouveaux Cerambycidae Lamiinae néotropicaux. *Bulletin de la Société Entomologique de France, Paris* 85 (3-4): 67-71.
- Buck, P.** 1959. Cerambycidae in der Sammlung des Instituto Anchieta de Pesquisas. *Pesquisas, Porto Alegre* 3: 577-609.
- Cazier, M., and Lacey, L.** 1952. The Cerambycidae of the Bahama Islands, British West Indies (Coleoptera). *American Museum Novitates*, New York 1588: 1-55.
- Chevrolat L. A. A.** 1862. Coléoptères de l'Île de Cuba. Notes, synonymies et descriptions d'espèces nouvelles. Familles des cérambycides et des parandrides. *Annales de la Société Entomologique de France, Paris* 4 (2): 245-280.
- Dalman, J. W.** In Schönher 1817. *Synonymia insectorum*, oder: Versuch einer Synonymie aller bisher bekannte Insekten; nach Fabricii Systema Eleutheratorum geordnet. Eleutherata oder Käfer 1 (3): xi + 1-506. Stockholm.
- Degeer, C.** 1775. *Mémoires pour servir à l'histoire des insectes* 5: 1-448, 16 pls. Stockholm.
- Dejean, P. F. M. A.** 1821. Catalogue des coléoptères de la collection de M. le baron Dejean. 136 pp. Paris.
- Dejean, P. F. M. A.** 1835. Catalogue des coléoptères de la collection de M. le Comte Dejean. livr. 4, 257-360. Paris.
- Dejean, P. F. M. A.** 1837. Catalogue des coléoptères de la collection de M. le Comte Dejean. livr. 5, 361-443. [Seconde Edition] Paris.
- Dillon, L. S. and Dillon, E. S.** 1941. The tribe Monochamini in the Western Hemisphere (Coleoptera: Cerambycidae). Scientific Publications. Reading Public Museum and Art Gallery. Reading, Pennsylvania. 1: 1-135, 5 pls.
- Dillon, L. S., and Dillon, E. S.** 1945. The tribe Onciderini (Coleoptera: Cerambycidae). Part I. Scientific Publications. Reading Public Museum and Art Gallery. Reading, Pennsylvania. 5: i-xv + 1-186.
- Dupont, H.** 1838. Monographie des trachydérides de la famille des longicornes. *Magasin de Zoologie, Paris* 8: i-xiii + 29-59, pls. 186-200 and 204-224.
- Erichson, W. F.** 1848. Die Insekten in Schomburg's Reisen in Britisch-Guiana. Leipzig 3: 570-574.
- Fabricius, J. C.** 1775. *Systema entomologiae, sistens insectorum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus, observationibus*. Korte; Flensburgi et Lipsiae 30 + 832 pp.
- Fabricius, J. C.** 1787. *Mantissa insectorum sistens eorum species nuper detectas adjectis characteribus genericis, differentiis specificis, emendationibus, observationibus*. Proft, Hafniae 1: xx + 348 pp.
- Fabricius, J. C.** 1792. *Entomologia systematica, emendata et aucta. Secundum classes ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus*. Proft, Hafniae 1: xx + 538 pp.
- Fabricius, J. C.** 1801. *Systema eleutheratorum secundum ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus*. Bibliopoli Academici Novi, Kiliae 2: 1-687.
- Fisher, W. S.** 1926. Descriptions of new West Indian longicorn beetles of the subfamily Lamiinae. *Proceedings of United States National Museum, Washington* 68 (22) 2623: 1-40.
- Fisher, W. S.** 1930. Notes on the Rhinotragine beetles of the family cerambycidae, with descriptions of new species. *Proceedings of United States National Museum, Washington* 77(19):1-20.
- Gahan, C.** 1895. On the longicorn Coleoptera of the West India Islands. *Transactions of the Entomological Society of London* [1895]: 79-140.
- Galileo, M. H. M. and Martins, U. R.** 1988a. Notas sobre Agapanthiini (Coleoptera, Cerambycidae, Lamiinae). I. Hippopsis do grupo solangeae. *Revista Brasileira de Entomologia, São Paulo* 32 (2): 179-185, 20 figs.
- Galileo, M. H. M., and Martins, U. R.** 1988b. Notas sobre Agapanthiini (Coleoptera, Cerambycidae, Lamiinae). V. Revisão do gênero Hippopsis Lepeletier and A.-Serville, 1825. *Revista Brasileira de Entomologia, São Paulo* 32 (2): 199-207, 9 figs.
- Galileo, M. H. M., and Martins, U. R.** 1991. Revisão da tribo Calliini (Coleoptera, Cerambycidae, Lamiinae).

- nae). *Giornale italiano de Entomologia*, 5: 243-262, illus.
- Gemminger, M., and Harold, E.** 1872. *Catalogus coleopterorum hucusque descriptorum synonymicus et systematicus*. Sumptu E. H. Gummi (G. Beck) Monachii 9: 2669-2988.
- Gemminger, M., and Harold, E.** 1873. *Catalogus coleopterorum hucusque descriptorum synonymicus et systematicus*. Monachii 10: 2989-3232 and index.
- Gilmour, E. F.** 1962. Synopsis of the tribe Aerenicini (Coleoptera, Cerambycidae, Lamiinae). *Revista de Biologia Tropical, Costa Rica* 10 (2): 123-147, 11 figs.
- Gilmour, E. F.** 1962. On the Neotropical Acanthocinini (Coleoptera, Cerambycidae, Lamiinae). Some new Genera and Generic Revisions. *Beiträge zur neotropischen Fauna, Jena* 2 (4): 249-293, 4 pls.
- Gilmour, E. F.** 1965. *Catalogue des Lamiaires du Monde (Col. Ceramb.)*. Verlag des Museums G. Frey Tutzing bei München 8: 559-655.
- Gmelin, J. F.** 1790. *Caroli a Linné Systema Naturae*. Edit. 13, 1 (4): 1517-2224.
- Haldeman, S. S.** 1847. Materials toward a study of the Coleoptera Longicornia of the United States. *Transactions of the American Philosophical Society* 10: 27-66.
- Hüdepohl, K. E.** 1979. Revision der Gattung *Oxymerus* Serville (Coleoptera, Cerambycidae, Cerambycinae, Trachyderini). *Entomologische Arbeiten aus dem Museum G. Frey, Tutzing bei München* 28: 9-74, 13 figs, 2 pls, 5 cartes.
- Lacordaire, J. T.** 1869. *Genera des Coléoptères*. (Contenant les familles des Tricentenotomides et des longicornes). Paris, Librairie Encyclopédique de Roret 8: 1-552.
- Lacordaire, J. T.** 1872. *Genera des Coléoptères*. Famille LXVIII. Longicornes. (suite). Sous-famille III. LAMIIDES. Paris, Librairie Encyclopédique de Roret 9 (2): 411-930.
- Lameere, A. A. L.** 1903. Révision des prionides. (Sixième mémoire - Basitoxus). *Annales de la Société Entomologique de Belgique, Bruxelles* 47: 213-224.
- Lameere, A. A. L.** 1904. Révision des prionides. (Neuvième mémoire - Callipogonines). *Annales de la Société Entomologique de Belgique, Bruxelles* 48 (1): 7-78.
- Lameere, A. A. L.** 1912. Révision des Prionides. Vingt-deuxième Mémoire. - Addenda et Corrigenda. *Mémoires de la Société Entomologique de Belgique, Bruxelles* 21: 113-188.
- Lameere, A. A. L.** 1913. Cerambycidae: Prioninae. *Coleopterum Catalogus* 22 (52): 1-108. W. Junk und S. Schenkling, Berlin.
- Lane, F.** 1958. The Neotropical Genus *Cryptocranium* Serville, with the description of a new species (Coleoptera, Lamiidae). *American Museum Novitates*, New York 1916: 1-10, 2 figs.
- Laporte, F. L. dit Castelnau** 1840. *Histoire Naturelle des Insectes Coléoptères*. 2: 1-563, 36 pls. P. Duménil, Paris.
- Leconte, J. L.** 1852. An attempt to classify the longicorn Coleoptera of the part of America, north of Mexico. *Journal of the Academy of Natural Sciences of Philadelphia* (2) 2: 139-178.
- Leconte, J. L.** 1873. Classification of the Coleoptera of North America. Part II. *Smithsonian Miscellaneous Collections, Washington* 11 (265): 279-348.
- Linné, C.** 1758. *Systema naturae per regna tria naturae secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. ed. 10, Laur. Salvius 1: i-iii + 824 pp. Holmiae.
- Linsley, E. G.** 1962. The Cerambycidae of North America. Part III. Taxonomy and Classification of the subfamily Cerambycinae, Tribes Opsimini through Megaderini. University of California. *Publications in Entomology* 20: 1-188, 56 figs.
- Martins, U. R., and Galileo, M. H. M.** 1990 a. Notas sobre Calliini (Coleoptera, Cerambycidae, Lamiinae). IV. As espécies sul-americanas do gênero *Drycothaea*. *Revista Brasileira de Entomologia, São Paulo* 34(3): 607-613, 5 figs.
- Martins, U. R., and Galileo, M. H. M.** 1990 b. Lamiinae (Coleoptera, Cerambycidae) com garras tarsais apendiculadas e descrição de Pretiliini, trib. n. *Revista Brasileira de Entomologia, São Paulo* 34(4): 703-708, 11 figs.
- Martins, U. R., and Moure, J. S.** 1973. Notas sobre Cerambycidae (Coleoptera), VII. *Revista Brasileira de Entomologia, São Paulo* 17 (2): 77-84, 2 figs.
- Merian, M.S.** 1719. *Dissertatio de generatione et metamorphosis insectorum Surinamensium Amstelædami*.
- Monné, M. A.** 1985. Sinopse do gênero *Nyssodrysternum* Gilmour, 1960, com notas sobre o complexo *Nyssodryus*. (Coleoptera, Cerambycidae, Lamiinae, Acanthocinini). *Revista Brasileira de Entomologia, São Paulo* 29 (3-4): 535-548, 9 figs.
- Monné, M. A., and Giesbert, E. F.** 1992. Nomenclatural notes on Western Hemisphere Cerambycidae (Coleoptera). *Insecta Mundi* 6 (1-2): 249-255.
- Newman, E.** 1840. *Entomological notes*. *Entomologist* 1: 1-16.
- Olivier, A. G.** 1790. *Entomologie, ou histoire naturelle des insectes, avec leurs caractères génériques et spécifiques, leur description, leur synonymie, et leur figure enluminée*. Coléoptères, vol. 2. Paris.
- Olivier, A. G.** 1792. *Encyclopédie méthodique. Histoire naturelle. Insectes*, vol. 6 (pt. 2), 369-704.
- Olivier, A. G.** 1795. *Entomologie, ou histoire naturelle des insectes, avec leurs caractères génériques et spécifiques, leur description, leur synonymie, et leur figure enluminée*. Coléoptères 4: 1-519, 72 pls. Baudouin, Paris.
- Panzer, G. W. F.** 1794. *Voets Beschreibungen und Abbildungen hartsschaaliger Insecten* pt. 2, Edition 3. Erlangen [1793] 112 pp.

- Panzer, G. W. F.** 1798. Voets Beschreibungen und Abbildungen hartsschaaliger Insecten pt. 4, 112 pp., pls. 25-48. Nörnberg.
- Pascoe, F. P.** 1859. On New Genera and Species of Longicorn Coleoptera. Part IV. The Transactions of the Entomological Society of London, ser. 2, vol. 5, 12-61, 1 pl.
- Say, T.** 1832. New species of North American insects, found by Joseph Barabino, chiefly in Louisiana. New Harmony, Indiana, 16 pp.
- Schaeffer, C. F. A.** 1908. List of the longicorn Coleoptera collected on the Museum expedition to Brownsville, Texas and the Huachuca Mts., Arizona with descriptions of new genera and species and notes on known species. Science Bulletin of the Museums Brooklyn Institute of Arts and Sciences 1 (12): 325-352.
- Schönherr, C. J.** 1817. Synonymia insectorum. Skara. Lewerentzischen Buchdruckerey 1 (3): xi + 1-506.
- Schwarzer, B.** 1929. Neue Cerambyciden des Senckenberg-Museums (Ins. Col.). Senckenbergiana, Frankfurt am Mein 11 (5-6): 353-379, 35 figs.
- Swederus, N. S.** 1787. Et nytt genus, och femtio nya species af Insekter beskrefne. Svenska Vetenskaps Akademien Nya Handlingar 8: 181-201.
- Tavakilian, G. L.** 1991. Notas sinonímicas e novas combinações em longicórneos sul-americanos (Coleoptera, Cerambycidae). Revista Brasileira de Entomologia, São Paulo, 35 (2): 439-453, 14 figs.
- Thomson, J.** 1860. Essai d'une classification de la famille des cérambycides et matériaux pour servir à une monographie de cette famille. 404 pages, illus. Paris.
- Thomson, J.** 1864. Systema cerambycidarum ou exposé de tous les genres compris dans la famille des cérambycides et familles limitrophes. Mémoires de la Société Royale des Sciences de Liège 19: 1-540.
- Thomson, J.** 1865. Diagnoses d'espèces nouvelles qui seront décrites dans l'appendix du systema cerambycidarum. Mémoires de la Société Royale des Sciences de Liège 19: 541-578.
- Thomson, J.** 1868 a. Révision du groupe des oncidérites (lamites, cérambycides, coléoptères). Physis Recueil d'Histoire Naturelle, Paris 2 (5): 41-92.
- Thomson, J.** 1868 b. Matériaux pour servir à une Révision des desmiphorites (lamites, cérambycides, coléoptères). Physis Recueil d'Histoire Naturelle, Paris 2 (6): 101-146.
- Thomson, J.** 1877. Typi cerambycidarum musei Thomsoniani. Revue et Magasin de Zoologie, Paris (3) 5: 249-279.
- Thunberg, C. P.** 1822. Trachyderes insecti genus ulterius examinatum et auctum sex novis speciebus. Mémoires de l'Académie Impériale des Sciences de Saint-Petersbourg 8: 303-307, 1 pl.
- Tippmann, F. F.** 1960. Studien über neotropische Longicornier - III (Coleoptera: Cerambycidae). Koleopterologische Rundschau, Wien 37/38 (1959/60) (1-6): 82-218, 14 pls, 35 figs.
- Villiers, A.** 1980. Coléoptères Cerambycidae des Antilles Françaises I. Parandrinae, Prioninae, Lepturinae. Annales de la Société Entomologique de France. Paris (Nouvelle Série) 16 (1): 133-157, 22 figs.
- Voet, J. E.** 1778. Catalogus systematicus coleopterum. La Haye, Backhuysen 2: 1-254, 50 pls.
- Waterhouse, C. O.** 1880. New South-American Coleoptera, chiefly from Ecuador. The Annals and Magazine of Natural History. London 5(5): 285-302.
- White, A.** 1855. Catalogue of coleopterous insects in the collection of the British Museum, pt. VIII. Longicornia II, 175-412, pls. 5-10. London.
- Williams, S. H.** 1929. A list of prionid beetles taken at Kartabo, Bartica district, British Guiana, with the description of a new species. The Annals of the Carnegie Museum 19: 139-148, pls 4-5.
- Zajciw, D.** 1969. Três espécies novas do gênero *Tomopterus* Serville, 1833 (Col., Cerambycidae, Rhinotragini). Studia Entomologica, Petrópolis 12 (1-4): 411-416, 2 figs.
- Zimsen, E.** 1964. The type material of I.C. Fabricius. 656 pp. Munskgaard, Copenhagen.

Book Review

Biologie des Coleopteres Chrysomelides. Pierre Jolivet. Boubee Publisher, Paris, 1997, 280 pp. (USA \$57.00)

The main biological aspects of this economically important group of beetles, the Chrysomelidae, are summarized in this new work by the internationally known author, Dr. Pierre Jolivet. Around 37,000 species are now described and more remain to be named. The size and diversity of the family are less than those of the weevils (Curculionidae), but probably there will be a total of 40 to 50 thousand species once the canopy fauna is better known. For instance, we still know nothing about the biology and the host plants of the Madagascan leaf beetles and surely many more species from that region await description.

The importance of the group is mostly agricultural as might be expected from a group almost entirely leaf, stem, and root feeders. Members range in length from 1 mm, or less, up to 27 mm. Many species are brightly colored.

Probably the family is polyphyletic, *i.e.*, composed of morphologically similar but phylogenetically distinct groups, and with Bruchidae as its sister group, all have Cerambycoid ancestors. Unquestioned chrysomelid fossils date from the Jurassic.

The book covers the palaeontology, evolution, development, and adaptations of the family. Mimicry, defensive reactions, pathogens and parasites, and every aspect of the biology and ecology of the beetles are discussed.

The classification adopted for this book is not that of cladists. Jolivet recognizes 20 extant subfamilies, and one extinct subfamily, probably the direct Jurassic ancestor of the Aulacoscelinae. The larvae of several subfamilies were still unknown at the time this book was written, such as the those of the Megascelinae and Aulacoscelinae, but recently, I am told by the author, these larvae have been obtained and are now being described.

The author believes that the recently proposed monophyletic classifications do not reflect the real relationships within the family. He disagrees with the fusion of Alticinae and Galerucinae, of Hispinae and Cassidinae, and Synetinae and Megascelinae, included with the Eumolpinae, as well as the elevation to family rank of some subfamilies. *Syneta*, for instance, has its wings and male genitalia completely different from those of the Eumolpinae. In the book he argues strongly that the relationships recognized within the Chrysomelidae be based on their food-plant selection when these are known, wing venation, and male genitalia, and use both physiological and mechanical characters of the plants.

Line drawings illustrate the book and six colored plates show the beetles in various tropical settings in both the Old and New World. Three other plates include some water color drawings of spectacular species.

The author, with several collaborators, has written five specialized books on the biology of Chrysomelidae prior to this volume. Here, the author presents a new synthesis of the topic, including many aspects not found in the previous volumes. The collected works were noted on pages 260 and 316 of volume 10, 1996, of *Insecta Mundi*.

R. H. Arnett, Jr.