

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Kenneth Bloom Publications

Research Papers in Physics and Astronomy

---

11-28-2008

## Search for Neutral Higgs Bosons in Multi-b-Jet Events in $pp$ Collisions at $\sqrt{s} = 1.96$ TeV

V. M. Abazov

*Joint Institute for Nuclear Research, Dubna, Russia*

Kenneth A. Bloom

*University of Nebraska-Lincoln, kenbloom@unl.edu*

Gregory Snow

*University of Nebraska-Lincoln, gsnow1@unl.edu*

D0 Collaboration

Follow this and additional works at: <https://digitalcommons.unl.edu/physicsbloom>



Part of the [Physics Commons](#)

---

Abazov, V. M.; Bloom, Kenneth A.; Snow, Gregory; and Collaboration, D0, "Search for Neutral Higgs Bosons in Multi-b-Jet Events in  $pp$  Collisions at  $\sqrt{s} = 1.96$  TeV" (2008). *Kenneth Bloom Publications*. 263.  
<https://digitalcommons.unl.edu/physicsbloom/263>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Kenneth Bloom Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# Search for Neutral Higgs Bosons in Multi-*b*-Jet Events in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

V. M. Abazov,<sup>36</sup> B. Abbott,<sup>75</sup> M. Abolins,<sup>65</sup> B. S. Acharya,<sup>29</sup> M. Adams,<sup>51</sup> T. Adams,<sup>49</sup> E. Aguilo,<sup>6</sup> S. H. Ahn,<sup>31</sup> M. Ahsan,<sup>59</sup> G. D. Alexeev,<sup>36</sup> G. Alkhazov,<sup>40</sup> A. Alton,<sup>64,\*</sup> G. Alverson,<sup>63</sup> G. A. Alves,<sup>2</sup> M. Anastasoae,<sup>35</sup> L. S. Ancu,<sup>35</sup> T. Andeen,<sup>53</sup> S. Anderson,<sup>45</sup> B. Andrieu,<sup>17</sup> M. S. Anzelc,<sup>53</sup> M. Aoki,<sup>50</sup> Y. Arnoud,<sup>14</sup> M. Arov,<sup>60</sup> M. Arthaud,<sup>18</sup> A. Askew,<sup>49</sup> B. Åsman,<sup>41</sup> A. C. S. Assis Jesus,<sup>3</sup> O. Atramentov,<sup>49</sup> C. Avila,<sup>8</sup> F. Badaud,<sup>13</sup> A. Baden,<sup>61</sup> L. Bagby,<sup>50</sup> B. Baldin,<sup>50</sup> D. V. Bandurin,<sup>59</sup> P. Banerjee,<sup>29</sup> S. Banerjee,<sup>29</sup> E. Barberis,<sup>63</sup> A.-F. Barfuss,<sup>15</sup> P. Bargassa,<sup>80</sup> P. Baringer,<sup>58</sup> J. Barreto,<sup>2</sup> J. F. Bartlett,<sup>50</sup> U. Bassler,<sup>18</sup> D. Bauer,<sup>43</sup> S. Beale,<sup>6</sup> A. Bean,<sup>58</sup> M. Begalli,<sup>3</sup> M. Begel,<sup>73</sup> C. Belanger-Champagne,<sup>41</sup> L. Bellantoni,<sup>50</sup> A. Bellavance,<sup>50</sup> J. A. Benitez,<sup>65</sup> S. B. Beri,<sup>27</sup> G. Bernardi,<sup>17</sup> R. Bernhard,<sup>23</sup> I. Bertram,<sup>42</sup> M. Besançon,<sup>18</sup> R. Beuselinck,<sup>43</sup> V. A. Bezzubov,<sup>39</sup> P. C. Bhat,<sup>50</sup> V. Bhatnagar,<sup>27</sup> C. Biscarat,<sup>20</sup> G. Blazey,<sup>52</sup> F. Blekman,<sup>43</sup> S. Blessing,<sup>49</sup> D. Bloch,<sup>19</sup> K. Bloom,<sup>67</sup> A. Boehnlein,<sup>50</sup> D. Boline,<sup>62</sup> T. A. Bolton,<sup>59</sup> E. E. Boos,<sup>38</sup> G. Borissov,<sup>42</sup> T. Bose,<sup>77</sup> A. Brandt,<sup>78</sup> R. Brock,<sup>65</sup> G. Brooijmans,<sup>70</sup> A. Bross,<sup>50</sup> D. Brown,<sup>81</sup> N. J. Buchanan,<sup>49</sup> D. Buchholz,<sup>53</sup> M. Buehler,<sup>81</sup> V. Buescher,<sup>22</sup> V. Bunichev,<sup>38</sup> S. Burdin,<sup>42,†</sup> S. Burke,<sup>45</sup> T. H. Burnett,<sup>82</sup> C. P. Buszello,<sup>43</sup> J. M. Butler,<sup>62</sup> P. Calfayan,<sup>25</sup> S. Calvet,<sup>16</sup> J. Cammin,<sup>71</sup> W. Carvalho,<sup>3</sup> B. C. K. Casey,<sup>50</sup> H. Castilla-Valdez,<sup>33</sup> S. Chakrabarti,<sup>18</sup> D. Chakraborty,<sup>52</sup> K. Chan,<sup>6</sup> K. M. Chan,<sup>55</sup> A. Chandra,<sup>48</sup> F. Charles,<sup>19,\*\*</sup> E. Cheu,<sup>45</sup> F. Chevallier,<sup>14</sup> D. K. Cho,<sup>62</sup> S. Choi,<sup>32</sup> B. Choudhary,<sup>28</sup> L. Christofek,<sup>77</sup> T. Christoudias,<sup>43</sup> S. Cihangir,<sup>50</sup> D. Claes,<sup>67</sup> J. Clutter,<sup>58</sup> M. Cooke,<sup>80</sup> W. E. Cooper,<sup>50</sup> M. Corcoran,<sup>80</sup> F. Couderc,<sup>18</sup> M.-C. Cousinou,<sup>15</sup> S. Crépe-Renaudin,<sup>14</sup> D. Cutts,<sup>77</sup> M. Ćwiok,<sup>30</sup> H. da Motta,<sup>2</sup> A. Das,<sup>45</sup> G. Davies,<sup>43</sup> K. De,<sup>78</sup> S. J. de Jong,<sup>35</sup> E. De La Cruz-Burelo,<sup>64</sup> C. De Oliveira Martins,<sup>3</sup> J. D. Degenhardt,<sup>64</sup> F. Déliot,<sup>18</sup> M. Demarteau,<sup>50</sup> R. Demina,<sup>71</sup> D. Denisov,<sup>50</sup> S. P. Denisov,<sup>39</sup> S. Desai,<sup>50</sup> H. T. Diehl,<sup>50</sup> M. Diesburg,<sup>50</sup> A. Dominguez,<sup>67</sup> H. Dong,<sup>72</sup> L. V. Dudko,<sup>38</sup> L. Dufloot,<sup>16</sup> S. R. Dugad,<sup>29</sup> D. Duggan,<sup>49</sup> A. Duperrin,<sup>15</sup> J. Dyer,<sup>65</sup> A. Dyshkant,<sup>52</sup> M. Eads,<sup>67</sup> D. Edmunds,<sup>65</sup> J. Ellison,<sup>48</sup> V. D. Elvira,<sup>50</sup> Y. Enari,<sup>77</sup> S. Eno,<sup>61</sup> P. Ermolov,<sup>38</sup> H. Evans,<sup>54</sup> A. Evdokimov,<sup>73</sup> V. N. Evdokimov,<sup>39</sup> A. V. Ferapontov,<sup>59</sup> T. Ferbel,<sup>71</sup> F. Fiedler,<sup>24</sup> F. Filthaut,<sup>35</sup> W. Fisher,<sup>50</sup> H. E. Fisk,<sup>50</sup> M. Fortner,<sup>52</sup> H. Fox,<sup>42</sup> S. Fu,<sup>50</sup> S. Fuess,<sup>50</sup> T. Gadfort,<sup>70</sup> C. F. Galea,<sup>35</sup> E. Gallas,<sup>50</sup> C. Garcia,<sup>71</sup> A. Garcia-Bellido,<sup>82</sup> V. Gavrilov,<sup>37</sup> P. Gay,<sup>13</sup> W. Geist,<sup>19</sup> D. Gelé,<sup>19</sup> C. E. Gerber,<sup>51</sup> Y. Gershtein,<sup>49</sup> D. Gillberg,<sup>6</sup> G. Ginther,<sup>71</sup> N. Gollub,<sup>41</sup> B. Gómez,<sup>8</sup> A. Goussiou,<sup>82</sup> P. D. Grannis,<sup>72</sup> H. Greenlee,<sup>50</sup> Z. D. Greenwood,<sup>60</sup> E. M. Gregores,<sup>4</sup> G. Grenier,<sup>20</sup> Ph. Gris,<sup>13</sup> J.-F. Grivaz,<sup>16</sup> A. Grohsjean,<sup>25</sup> S. Grünendahl,<sup>50</sup> M. W. Grünewald,<sup>30</sup> F. Guo,<sup>72</sup> J. Guo,<sup>72</sup> G. Gutierrez,<sup>50</sup> P. Gutierrez,<sup>75</sup> A. Haas,<sup>70</sup> N. J. Hadley,<sup>61</sup> P. Haefner,<sup>25</sup> S. Hagopian,<sup>49</sup> J. Haley,<sup>68</sup> I. Hall,<sup>65</sup> R. E. Hall,<sup>47</sup> L. Han,<sup>7</sup> K. Harder,<sup>44</sup> A. Harel,<sup>71</sup> J. M. Hauptman,<sup>57</sup> R. Hauser,<sup>65</sup> J. Hays,<sup>43</sup> T. Hebbeker,<sup>21</sup> D. Hedin,<sup>52</sup> J. G. Hegeman,<sup>34</sup> A. P. Heinson,<sup>48</sup> U. Heintz,<sup>62</sup> C. Hensel,<sup>22,§</sup> K. Herner,<sup>72</sup> G. Hesketh,<sup>63</sup> M. D. Hildreth,<sup>55</sup> R. Hirosky,<sup>81</sup> J. D. Hobbs,<sup>72</sup> B. Hoeneisen,<sup>12</sup> H. Hoeth,<sup>26</sup> M. Hohlfield,<sup>22</sup> S. J. Hong,<sup>31</sup> S. Hossain,<sup>75</sup> P. Houben,<sup>34</sup> Y. Hu,<sup>72</sup> Z. Hubacek,<sup>10</sup> V. Hynek,<sup>9</sup> I. Iashvili,<sup>69</sup> R. Illingworth,<sup>50</sup> A. S. Ito,<sup>50</sup> S. Jabeen,<sup>62</sup> M. Jaffré,<sup>16</sup> S. Jain,<sup>75</sup> K. Jakobs,<sup>23</sup> C. Jarvis,<sup>61</sup> R. Jesik,<sup>43</sup> K. Johns,<sup>45</sup> C. Johnson,<sup>70</sup> M. Johnson,<sup>50</sup> A. Jonckheere,<sup>50</sup> P. Jonsson,<sup>43</sup> A. Juste,<sup>50</sup> E. Kajfasz,<sup>15</sup> J. M. Kalk,<sup>60</sup> D. Karmanov,<sup>38</sup> P. A. Kasper,<sup>50</sup> I. Katsanos,<sup>70</sup> D. Kau,<sup>49</sup> V. Kaushik,<sup>78</sup> R. Kehoe,<sup>79</sup> S. Kermiche,<sup>15</sup> N. Khalatyan,<sup>50</sup> A. Khanov,<sup>76</sup> A. Kharchilava,<sup>69</sup> Y. M. Kharzheev,<sup>36</sup> D. Khatidze,<sup>70</sup> T. J. Kim,<sup>31</sup> M. H. Kirby,<sup>53</sup> M. Kirsch,<sup>21</sup> B. Klima,<sup>50</sup> J. M. Kohli,<sup>27</sup> J.-P. Konrath,<sup>23</sup> A. V. Kozelov,<sup>39</sup> J. Kraus,<sup>65</sup> D. Krop,<sup>54</sup> T. Kuhl,<sup>24</sup> A. Kumar,<sup>69</sup> A. Kupco,<sup>11</sup> T. Kurča,<sup>20</sup> V. A. Kuzmin,<sup>38</sup> J. Kvita,<sup>9</sup> F. Lacroix,<sup>13</sup> D. Lam,<sup>55</sup> S. Lammers,<sup>70</sup> G. Landsberg,<sup>77</sup> P. Lebrun,<sup>20</sup> W. M. Lee,<sup>50</sup> A. Leflat,<sup>38</sup> J. Lellouch,<sup>17</sup> J. Leveque,<sup>45</sup> J. Li,<sup>78</sup> L. Li,<sup>48</sup> Q. Z. Li,<sup>50</sup> S. M. Lietti,<sup>5</sup> J. G. R. Lima,<sup>52</sup> D. Lincoln,<sup>50</sup> J. Linnemann,<sup>65</sup> V. V. Lipaev,<sup>39</sup> R. Lipton,<sup>50</sup> Y. Liu,<sup>7</sup> Z. Liu,<sup>6</sup> A. Lobodenko,<sup>40</sup> M. Lokajicek,<sup>11</sup> P. Love,<sup>42</sup> H. J. Lubatti,<sup>82</sup> R. Luna,<sup>3</sup> A. L. Lyon,<sup>50</sup> A. K. A. Maciel,<sup>2</sup> D. Mackin,<sup>80</sup> R. J. Madaras,<sup>46</sup> P. Mättig,<sup>26</sup> C. Magass,<sup>21</sup> A. Magerkurth,<sup>64</sup> P. K. Mal,<sup>82</sup> H. B. Malbouisson,<sup>3</sup> S. Malik,<sup>67</sup> V. L. Malyshev,<sup>36</sup> H. S. Mao,<sup>50</sup> Y. Maravin,<sup>59</sup> B. Martin,<sup>14</sup> R. McCarthy,<sup>72</sup> A. Melnitchouk,<sup>66</sup> L. Mendoza,<sup>8</sup> P. G. Mercadante,<sup>5</sup> M. Merkin,<sup>38</sup> K. W. Merritt,<sup>50</sup> A. Meyer,<sup>21</sup> J. Meyer,<sup>22,§</sup> T. Millet,<sup>20</sup> J. Mitrevski,<sup>70</sup> R. K. Mommsen,<sup>44</sup> N. K. Mondal,<sup>29</sup> R. W. Moore,<sup>6</sup> T. Moulík,<sup>58</sup> G. S. Muanza,<sup>20</sup> M. Mulhearn,<sup>70</sup> O. Mundal,<sup>22</sup> L. Mundim,<sup>3</sup> E. Nagy,<sup>15</sup> M. Naimuddin,<sup>50</sup> M. Narain,<sup>77</sup> N. A. Naumann,<sup>35</sup> H. A. Neal,<sup>64</sup> J. P. Negret,<sup>8</sup> P. Neustroev,<sup>40</sup> H. Nilsen,<sup>23</sup> H. Nogima,<sup>3</sup> S. F. Novaes,<sup>5</sup> T. Nunnemann,<sup>25</sup> V. O'Dell,<sup>50</sup> D. C. O'Neil,<sup>6</sup> G. Obrant,<sup>40</sup> C. Ochando,<sup>16</sup> D. Onoprienko,<sup>59</sup> N. Oshima,<sup>50</sup> N. Osman,<sup>43</sup> J. Osta,<sup>55</sup> R. Otec,<sup>10</sup> G. J. Otero y Garzón,<sup>50</sup> M. Owen,<sup>44</sup> P. Padley,<sup>80</sup> M. Pangilinan,<sup>77</sup> N. Parashar,<sup>56</sup> S.-J. Park,<sup>22,§</sup> S. K. Park,<sup>31</sup> J. Parsons,<sup>70</sup> R. Partridge,<sup>77</sup> N. Parua,<sup>54</sup> A. Patwa,<sup>73</sup> G. Pawloski,<sup>80</sup> B. Penning,<sup>23</sup> M. Perfilov,<sup>38</sup> K. Peters,<sup>44</sup> Y. Peters,<sup>26</sup> P. Pétroff,<sup>16</sup> M. Petteni,<sup>43</sup> R. Piegaia,<sup>1</sup> J. Piper,<sup>65</sup> M.-A. Pleier,<sup>22</sup> P. L. M. Podesta-Lerma,<sup>33,‡</sup> V. M. Podstavkov,<sup>50</sup> Y. Pogorelov,<sup>55</sup> M.-E. Pol,<sup>2</sup> P. Polozov,<sup>37</sup> B. G. Pope,<sup>65</sup> A. V. Popov,<sup>39</sup> C. Potter,<sup>6</sup> W. L. Prado da Silva,<sup>3</sup> H. B. Prosper,<sup>49</sup> S. Protopopescu,<sup>73</sup> J. Qian,<sup>64</sup> A. Quadt,<sup>22,§</sup> B. Quinn,<sup>66</sup> A. Rakitine,<sup>42</sup> M. S. Rangel,<sup>2</sup> K. Ranjan,<sup>28</sup> P. N. Ratoff,<sup>42</sup> P. Renkel,<sup>79</sup> S. Reucroft,<sup>63</sup> P. Rich,<sup>44</sup> J. Rieger,<sup>54</sup> M. Rijssenbeek,<sup>72</sup> I. Ripp-Baudot,<sup>19</sup> F. Rizatdinova,<sup>76</sup> S. Robinson,<sup>43</sup> R. F. Rodrigues,<sup>3</sup>

M. Rominsky,<sup>75</sup> C. Royon,<sup>18</sup> P. Rubinov,<sup>50</sup> R. Ruchti,<sup>55</sup> G. Safronov,<sup>37</sup> G. Sajot,<sup>14</sup> A. Sánchez-Hernández,<sup>33</sup>  
M. P. Sanders,<sup>17</sup> B. Sanghi,<sup>50</sup> A. Santoro,<sup>3</sup> G. Savage,<sup>50</sup> L. Sawyer,<sup>60</sup> T. Scanlon,<sup>43</sup> D. Schaile,<sup>25</sup> R. D. Schamberger,<sup>72</sup>  
Y. Scheglov,<sup>40</sup> H. Schellman,<sup>53</sup> T. Schliephake,<sup>26</sup> C. Schwanenberger,<sup>44</sup> A. Schwartzman,<sup>68</sup> R. Schwienhorst,<sup>65</sup>  
J. Sekaric,<sup>49</sup> H. Severini,<sup>75</sup> E. Shabalina,<sup>51</sup> M. Shamim,<sup>59</sup> V. Shary,<sup>18</sup> A. A. Shchukin,<sup>39</sup> R. K. Shivpuri,<sup>28</sup> V. Siccaldi,<sup>19</sup>  
V. Simak,<sup>10</sup> V. Sirotenko,<sup>50</sup> P. Skubic,<sup>75</sup> P. Slattey,<sup>71</sup> D. Smirnov,<sup>55</sup> G. R. Snow,<sup>67</sup> J. Snow,<sup>74</sup> S. Snyder,<sup>73</sup>  
S. Söldner-Rembold,<sup>44</sup> L. Sonnenschein,<sup>17</sup> A. Sopczak,<sup>42</sup> M. Sosebee,<sup>78</sup> K. Soustruznik,<sup>9</sup> B. Spurlock,<sup>78</sup> J. Stark,<sup>14</sup>  
J. Steele,<sup>60</sup> V. Stolin,<sup>37</sup> D. A. Stoyanova,<sup>39</sup> J. Strandberg,<sup>64</sup> S. Strandberg,<sup>41</sup> M. A. Strang,<sup>69</sup> E. Strauss,<sup>72</sup> M. Strauss,<sup>75</sup>  
R. Ströhmer,<sup>25</sup> D. Strom,<sup>53</sup> L. Stutte,<sup>50</sup> S. Sumowidagdo,<sup>49</sup> P. Svoisky,<sup>55</sup> A. Sznajder,<sup>3</sup> P. Tamburello,<sup>45</sup> A. Tanasijczuk,<sup>1</sup>  
W. Taylor,<sup>6</sup> J. Temple,<sup>45</sup> B. Tiller,<sup>25</sup> F. Tissandier,<sup>13</sup> M. Titov,<sup>18</sup> V. V. Tokmenin,<sup>36</sup> T. Toole,<sup>61</sup> I. Torchiani,<sup>23</sup> T. Trefzger,<sup>24</sup>  
D. Tsybychev,<sup>72</sup> B. Tuchming,<sup>18</sup> C. Tully,<sup>68</sup> P. M. Tuts,<sup>70</sup> R. Unalan,<sup>65</sup> L. Uvarov,<sup>40</sup> S. Uvarov,<sup>40</sup> S. Uzunyan,<sup>52</sup> B. Vachon,<sup>6</sup>  
P. J. van den Berg,<sup>34</sup> R. Van Kooten,<sup>54</sup> W. M. van Leeuwen,<sup>34</sup> N. Varelas,<sup>51</sup> E. W. Varnes,<sup>45</sup> I. A. Vasilyev,<sup>39</sup> M. Vaupel,<sup>26</sup>  
P. Verdier,<sup>20</sup> L. S. Vertogradov,<sup>36</sup> M. Verzocchi,<sup>50</sup> F. Villeneuve-Seguié,<sup>43</sup> P. Vint,<sup>43</sup> P. Vokac,<sup>10</sup> E. Von Toerne,<sup>59</sup>  
M. Voutilainen,<sup>68,||</sup> R. Wagner,<sup>68</sup> H. D. Wahl,<sup>49</sup> L. Wang,<sup>61</sup> M. H. L. S. Wang,<sup>50</sup> J. Warchol,<sup>55</sup> G. Watts,<sup>82</sup> M. Wayne,<sup>55</sup>  
G. Weber,<sup>24</sup> M. Weber,<sup>50</sup> L. Welty-Rieger,<sup>54</sup> A. Wenger,<sup>23,¶</sup> N. Worms,<sup>22</sup> M. Wetstein,<sup>61</sup> A. White,<sup>78</sup> D. Wicke,<sup>26</sup>  
G. W. Wilson,<sup>58</sup> S. J. Wimpenny,<sup>48</sup> M. Wobisch,<sup>60</sup> D. R. Wood,<sup>63</sup> T. R. Wyatt,<sup>44</sup> Y. Xie,<sup>77</sup> S. Yacoub,<sup>53</sup> R. Yamada,<sup>50</sup>  
M. Yan,<sup>61</sup> T. Yasuda,<sup>50</sup> Y. A. Yatsunenko,<sup>36</sup> K. Yip,<sup>73</sup> H. D. Yoo,<sup>77</sup> S. W. Youn,<sup>53</sup> J. Yu,<sup>78</sup> C. Zeitnitz,<sup>26</sup> T. Zhao,<sup>82</sup>  
B. Zhou,<sup>64</sup> J. Zhu,<sup>72</sup> M. Zielinski,<sup>71</sup> D. Zieminska,<sup>54</sup> A. Zieminski,<sup>54,\*\*</sup> L. Zivkovic,<sup>70</sup> V. Zutshi,<sup>52</sup> and E. G. Zverev<sup>38</sup>

(D0 Collaboration)

<sup>1</sup>Universidad de Buenos Aires, Buenos Aires, Argentina<sup>2</sup>LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil<sup>3</sup>Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil<sup>4</sup>Universidade Federal do ABC, Santo André, Brazil<sup>5</sup>Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil<sup>6</sup>University of Alberta, Edmonton, Alberta, Canada,

Simon Fraser University, Burnaby, British Columbia, Canada,

York University, Toronto, Ontario, Canada,

and McGill University, Montreal, Quebec, Canada

<sup>7</sup>University of Science and Technology of China, Hefei, People's Republic of China<sup>8</sup>Universidad de los Andes, Bogotá, Colombia<sup>9</sup>Center for Particle Physics, Charles University, Prague, Czech Republic<sup>10</sup>Czech Technical University, Prague, Czech Republic<sup>11</sup>Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic<sup>12</sup>Universidad San Francisco de Quito, Quito, Ecuador<sup>13</sup>LPC, Univ Blaise Pascal, CNRS/IN2P3, Clermont, France<sup>14</sup>LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, France<sup>15</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France<sup>16</sup>LAL, Univ Paris-Sud, IN2P3/CNRS, Orsay, France<sup>17</sup>LPNHE, IN2P3/CNRS, Universités Paris VI and VII, Paris, France<sup>18</sup>DAPNIA/Service de Physique des Particules, CEA, Saclay, France<sup>19</sup>IPHC, Université Louis Pasteur et Université de Haute Alsace, CNRS/IN2P3, Strasbourg, France<sup>20</sup>IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France<sup>21</sup>III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany<sup>22</sup>Physikalisches Institut, Universität Bonn, Bonn, Germany<sup>23</sup>Physikalisches Institut, Universität Freiburg, Freiburg, Germany<sup>24</sup>Institut für Physik, Universität Mainz, Mainz, Germany<sup>25</sup>Ludwig-Maximilians-Universität München, München, Germany<sup>26</sup>Fachbereich Physik, University of Wuppertal, Wuppertal, Germany<sup>27</sup>Panjab University, Chandigarh, India<sup>28</sup>Delhi University, Delhi, India<sup>29</sup>Tata Institute of Fundamental Research, Mumbai, India<sup>30</sup>University College Dublin, Dublin, Ireland<sup>31</sup>Korea Detector Laboratory, Korea University, Seoul, Korea<sup>32</sup>SungKyunKwan University, Suwon, Korea<sup>33</sup>CINVESTAV, Mexico City, Mexico<sup>34</sup>FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands

- <sup>35</sup>*Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands*  
<sup>36</sup>*Joint Institute for Nuclear Research, Dubna, Russia*  
<sup>37</sup>*Institute for Theoretical and Experimental Physics, Moscow, Russia*  
<sup>38</sup>*Moscow State University, Moscow, Russia*  
<sup>39</sup>*Institute for High Energy Physics, Protvino, Russia*  
<sup>40</sup>*Petersburg Nuclear Physics Institute, St. Petersburg, Russia*  
<sup>41</sup>*Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden, and Uppsala University, Uppsala, Sweden*  
<sup>42</sup>*Lancaster University, Lancaster, United Kingdom*  
<sup>43</sup>*Imperial College, London, United Kingdom*  
<sup>44</sup>*University of Manchester, Manchester, United Kingdom*  
<sup>45</sup>*University of Arizona, Tucson, Arizona 85721, USA*  
<sup>46</sup>*Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA*  
<sup>47</sup>*California State University, Fresno, California 93740, USA*  
<sup>48</sup>*University of California, Riverside, California 92521, USA*  
<sup>49</sup>*Florida State University, Tallahassee, Florida 32306, USA*  
<sup>50</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*  
<sup>51</sup>*University of Illinois at Chicago, Chicago, Illinois 60607, USA*  
<sup>52</sup>*Northern Illinois University, DeKalb, Illinois 60115, USA*  
<sup>53</sup>*Northwestern University, Evanston, Illinois 60208, USA*  
<sup>54</sup>*Indiana University, Bloomington, Indiana 47405, USA*  
<sup>55</sup>*University of Notre Dame, Notre Dame, Indiana 46556, USA*  
<sup>56</sup>*Purdue University Calumet, Hammond, Indiana 46323, USA*  
<sup>57</sup>*Iowa State University, Ames, Iowa 50011, USA*  
<sup>58</sup>*University of Kansas, Lawrence, Kansas 66045, USA*  
<sup>59</sup>*Kansas State University, Manhattan, Kansas 66506, USA*  
<sup>60</sup>*Louisiana Tech University, Ruston, Louisiana 71272, USA*  
<sup>61</sup>*University of Maryland, College Park, Maryland 20742, USA*  
<sup>62</sup>*Boston University, Boston, Massachusetts 02215, USA*  
<sup>63</sup>*Northeastern University, Boston, Massachusetts 02115, USA*  
<sup>64</sup>*University of Michigan, Ann Arbor, Michigan 48109, USA*  
<sup>65</sup>*Michigan State University, East Lansing, Michigan 48824, USA*  
<sup>66</sup>*University of Mississippi, University, Mississippi 38677, USA*  
<sup>67</sup>*University of Nebraska, Lincoln, Nebraska 68588, USA*  
<sup>68</sup>*Princeton University, Princeton, New Jersey 08544, USA*  
<sup>69</sup>*State University of New York, Buffalo, New York 14260, USA*  
<sup>70</sup>*Columbia University, New York, New York 10027, USA*  
<sup>71</sup>*University of Rochester, Rochester, New York 14627, USA*  
<sup>72</sup>*State University of New York, Stony Brook, New York 11794, USA*  
<sup>73</sup>*Brookhaven National Laboratory, Upton, New York 11973, USA*  
<sup>74</sup>*Langston University, Langston, Oklahoma 73050, USA*  
<sup>75</sup>*University of Oklahoma, Norman, Oklahoma 73019, USA*  
<sup>76</sup>*Oklahoma State University, Stillwater, Oklahoma 74078, USA*  
<sup>77</sup>*Brown University, Providence, Rhode Island 02912, USA*  
<sup>78</sup>*University of Texas, Arlington, Texas 76019, USA*  
<sup>79</sup>*Southern Methodist University, Dallas, Texas 75275, USA*  
<sup>80</sup>*Rice University, Houston, Texas 77005, USA*  
<sup>81</sup>*University of Virginia, Charlottesville, Virginia 22901, USA*  
<sup>82</sup>*University of Washington, Seattle, Washington 98195, USA*  
(Received 27 May 2008; published 26 November 2008)

Data recorded by the D0 experiment at the Fermilab Tevatron Collider are analyzed to search for neutral Higgs bosons produced in association with  $b$  quarks. This production mode can be enhanced in the minimal supersymmetric standard model (MSSM). The search is performed in the three  $b$  quark channel using multijet triggered events corresponding to an integrated luminosity of  $1 \text{ fb}^{-1}$ . No statistically significant excess of events with respect to the predicted background is observed and limits are set in the MSSM parameter space.



Supersymmetry (SUSY) [1] is a popular extension of the standard model (SM), requiring the presence of at least two Higgs doublets. In the minimal supersymmetric standard model (MSSM) five Higgs bosons remain after electroweak symmetry breaking: three neutral ( $h$ ,  $H$ , and  $A$ , collectively denoted as  $\phi$ ) and two charged ( $H^\pm$ ). The Higgs sector can be parameterized by  $\tan\beta$ , the ratio of the vacuum expectation values of the two Higgs doublets, and  $m_A$ , the mass of the pseudoscalar  $A$ . For large values of  $\tan\beta$  two of the three neutral Higgs bosons have approximately the same mass and couplings to down-type quarks, which are enhanced by a factor  $\tan\beta$  relative to the SM ones, while the couplings to up-type quarks are suppressed. More precisely, the three Higgs boson couplings to bottom quarks follow the sum rule  $g_{hbb}^2 + g_{Hbb}^2 + g_{Abb}^2 \approx 2\tan^2\beta g_{h_{SM}}^2$  where  $g_{h_{SM}}^2$  is the respective SM coupling. As well as increased production, the  $\tan\beta$  enhancement means that the main decay mode is  $\phi \rightarrow b\bar{b}$ , with a branching fraction,  $\mathcal{B}(\phi \rightarrow b\bar{b})$ , of  $\approx 90\%$ . Consequently, in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV at the Fermilab Tevatron Collider, the final state with at least three  $b$  jets is an important channel in the search for MSSM Higgs bosons at large  $\tan\beta$ . However, the very large multijet background at a hadron collider makes the search for this topology challenging.

LEP excluded  $m_{h,A} < 93$  GeV/ $c^2$  for all  $\tan\beta$  values [2]. CDF [3,4] and D0 [5,6] have extended the MSSM Higgs boson searches to higher masses for high  $\tan\beta$  values. The result presented in this Letter supersedes our previous published result [5]. In addition to including more data, this analysis benefits from improved signal and background modeling and an improved limit setting procedure, which uses only the shape, and not the normalization, of the final discriminating variable.

The D0 detector is described in Ref. [7]. Dedicated triggers designed to select events with at least three jets are used in this analysis. Typical trigger requirements are at least two jets with transverse momenta  $p_T > 25$  GeV/ $c$  and at least one additional jet with  $p_T > 15$  GeV/ $c$ . Algorithms for identifying  $b$  jets at the trigger level are also employed in about 70% of the integrated luminosity used for this analysis. After data quality requirements the total data sample corresponds to  $1.02 \pm 0.06$  fb $^{-1}$  [8].

Signal samples are generated for Higgs boson masses from 90–220 GeV/ $c^2$  using the leading order PYTHIA event generator [9] to generate associated production of  $\phi$  and a  $b$  quark in the 5-flavor scheme,  $gb \rightarrow \phi b$ . Weights, calculated with MCFM [10], are applied to the signal samples as a function of  $p_T$  and  $\eta$  of the highest  $p_T$   $b$  jet which is not from the decay of the Higgs boson to correct the cross section and experimental acceptance to next-to-leading order. Multijet background events from the  $b\bar{b}$ ,  $b\bar{b}j$ ,  $b\bar{b}jj$ ,  $c\bar{c}$ ,  $c\bar{c}j$ ,  $c\bar{c}jj$ ,  $b\bar{b}c\bar{c}$ , and  $b\bar{b}b\bar{b}$  processes (where  $j$  denotes a light parton:  $u$ ,  $d$ ,  $s$  quark or gluon) are generated with the ALPGEN [11] event generator. The con-

tributions from other processes, such as  $t\bar{t}$ ,  $Zb\bar{b}$ , and single top production, are found to be negligible. The ALPGEN samples are processed through PYTHIA for showering and hadronization. All samples are then processed through a GEANT-based [12] simulation of the D0 detector and the same reconstruction algorithms as the data. A parameterized trigger simulation is used to model the effects of the trigger requirements on the simulated events.

Jets are reconstructed from energy deposits in calorimeter towers using the midpoint cone algorithm [13] with radius = 0.5. Jet reconstruction and energy scale determination are described in detail in Ref. [14]. All calorimeter jets are required to pass a set of quality criteria and have at least two reconstructed tracks within  $\Delta\mathcal{R}(\text{track}, \text{jet-axis}) = \sqrt{(\Delta\eta)^2 + (\Delta\varphi)^2} < 0.5$  (where  $\eta$  is the pseudorapidity and  $\varphi$  the azimuthal angle), for a total per-jet efficiency of 83%–93%.

We select signal events for which the  $p\bar{p}$  interaction vertex is reconstructed well within the geometric acceptance of the silicon detector. We further require at least three and at most five jets with  $p_T > 20$  GeV/ $c$  and  $|\eta| < 2.5$ . A neural network based  $b$ -tagging algorithm [15], with lifetime based information involving the track impact parameters and secondary vertices as inputs, is used to identify  $b$  jets. Each event must have at least three jets satisfying a tight  $b$ -tag neural network requirement. This requirement provides  $\approx 50\%$  efficiency for tagging a single  $b$ -jet at  $\approx 0.4\%$  mistag rate of light jets ( $u$ ,  $d$ ,  $s$  quarks or gluons). The events with at least two tight  $b$  tags are also kept and used to model the background, since the relative amount of signal is negligible in this double tagged sample. Simulated events are weighted based on their tagging and fake rate probabilities determined from data. Finally, the transverse momenta of the two highest  $p_T$  jets which are also  $b$  tagged are required to be above 25 GeV/ $c$ . To further increase the sensitivity, the analysis is split into separate three-, four-, and five-jet channels. After the event selection 3224 events remain in the exclusive three-jet sample, 2503 and 704 events in the four- and five-jet samples, respectively. The signal efficiencies for Higgs boson masses between 100 and 200 GeV/ $c^2$  range from 0.3%–1.2% in the three-jet channel (0.2%–0.6% and 0.03%–0.12% in the four- and five-jet channels).

The background composition is determined separately for each jet multiplicity. By considering eight different  $b$ -tagging criteria, each applied to three subsamples (exactly one, two and three  $b$ -tagged jets, respectively), a linear system of equations can be solved to determine the relative contribution of the different processes since the  $b$ ,  $c$  and light jet tagging efficiencies are known. The double  $b$ -tagged (2Tag) sample is found to be dominated by  $b\bar{b}j$  while the triple  $b$ -tagged (3Tag) sample consists of a mix of  $\approx 50\%$   $b\bar{b}b$ ,  $\approx 30\%$   $b\bar{b}j$ , and  $\approx 20\%$   $b\bar{b}c + bc\bar{c}$ . An alternative method to determine the background, based on

fitting simulated  $H_T = \sum p_{T\text{jet}}$  shape templates to the data, confirms the composition of the background.

For every event the two jet pairs with the largest summed transverse momenta are considered as possible Higgs boson candidates. To remove discrepancies between data and simulation originating from gluon splitting ( $g \rightarrow b\bar{b}$ ), only jet pairs with  $\Delta\mathcal{R} > 1.0$  are considered in the final analysis. The agreement between data and simulation was verified in several control samples after the selection criteria.

The following six variables separate the Higgs boson jet pair from the background jet pairs and are well modeled by the simulation: the difference in pseudorapidity between the two jets in the pair; the azimuthal angular difference between the two jets in the pair; the angle between the leading jet in the pair and the total momentum of the pair;  $|p_1 - p_2|/|p_1 + p_2|$ , the momentum balance, where  $p_i$  ( $i = 1, 2$ ) are the momenta of the jets in the pair; the combined rapidity of the jet pair; and the event sphericity [16], defined as  $\frac{3}{2}(\epsilon_1 + \epsilon_2)$ , where  $\epsilon_n$  is the  $n$ th eigenvalue of the normalized momentum tensor calculated using all jets in the event. Based on these kinematic variables, a likelihood discriminant  $\mathcal{D}$ , is calculated according to:

$$\mathcal{D}(x_1, \dots, x_6) = \frac{\prod_{i=1}^6 P_i^{\text{sig}}(x_i)}{\prod_{i=1}^6 P_i^{\text{sig}}(x_i) + \prod_{i=1}^6 P_i^{\text{bkg}}(x_i)}, \quad (1)$$

where  $P_i^{\text{sig}}$  ( $P_i^{\text{bkg}}$ ) refers to the signal (background) probability density function (PDF) for variable  $x_i$ , and  $(x_1, \dots, x_6)$  is the set of measured kinematic variables for the jet pair. The PDFs are obtained from the 3Tag signal and background simulation. Two likelihoods are built combining simulated samples in the 90–130 GeV/ $c^2$  (“low-mass”) and 130–220 GeV/ $c^2$  (“high-mass”) mass ranges, providing discrimination at low and high masses, respectively. Studies show that this division of the mass range gives the best discrimination.

Several multijet processes contribute to the background and the uncertainty on the cross sections is large. The  $b\bar{b}b$  component may also contain a contribution that is indistinguishable from a signal and cannot be normalized from the data. To model the background we therefore rely on a combination of data and simulation. The distribution of the expected background in the 3Tag sample is obtained, in the two-dimensional  $\mathcal{D}$  and invariant mass ( $M_{bb}$ ) plane, by:

$$S_{3\text{Tag}}^{\text{exp}}(\mathcal{D}, M_{bb}) = R^{\text{MC}}(\mathcal{D}, M_{bb}) S_{2\text{Tag}}^{\text{data}}(\mathcal{D}, M_{bb}), \quad (2)$$

where  $R^{\text{MC}} = S_{3\text{Tag}}^{\text{MC}}/S_{2\text{Tag}}^{\text{MC}}$ ,  $S_{n\text{Tag}}^{\text{MC}}$  is the simulated  $n$ Tag background shape, and  $S_{2\text{Tag}}^{\text{data}}$  is the 2Tag data shape. Many uncertainties affecting the simulation cancel in the ratio  $R^{\text{MC}}$ . Figure 1 shows the low-mass likelihood for data and background in the three-jet channel.

The selection cuts on  $\mathcal{D}$ ,  $b$  tagging, and the number of jet-pair combinations per event are optimized by maximizing the expected sensitivity. The optimal cuts for the likelihood vary between 0.25 and 0.60 depending on the jet




FIG. 1 (color online). Comparison of the low-mass likelihood distributions for the 3Tag data and background (bkg exp.) defined by Eq. (2). Every event has two entries, one for each jet pair. Black crosses refer to data, the solid line shows the total background estimate, and the shaded region represents the heavy flavor component ( $b\bar{b}b$ ,  $b\bar{b}c$ , and  $c\bar{c}b$ ). The distribution for a Higgs boson of mass 100 GeV/ $c^2$  is also shown.

multiplicity and Higgs boson mass. The agreement of the data and the background expectation is verified in a control region where the impact of any Higgs boson signal is limited, defined by  $\mathcal{D} < 0.25$ . Figure 2 shows the invariant mass for the optimized high-mass likelihood cuts.

The number of signal events depends on several measurements which introduce a systematic uncertainty: integrated luminosity (6.1%), theoretical uncertainty (12%–14%), trigger efficiency (2%–5%), jet identification (0.3%–0.5%), jet energy calibration (3%–5%), jet energy resolution (0.1%–0.6%), and  $b$  jet identification (8%–9%). The size of these uncertainties depends on the Higgs boson mass and the number of jets. Several sources of systematic uncertainties affecting the background shape through the ratio  $R^{\text{MC}}$  in Eq. (2) are parametrized as a function of  $M_{bb}$ . The dominant uncertainty, due to the background composition, is estimated by varying the ratio of  $b\bar{b}j$  and  $b\bar{b}b$  events in the sample corresponding to the uncertainties from the background composition fit. The uncertainty from the kinematic dependence of the  $b$  tagging of jets is evaluated by varying the light,  $b$ , and  $c$  jet tag efficiencies within their uncertainties. The uncertainty from the  $b$ -jet energy resolution is obtained by smearing the simulated  $b$  and  $c$  jets by an additional factor of half the uncertainty of the light jet energy resolution. The effect due to the uncertainties in the kinematic modeling of  $b\bar{b}b$  and  $b\bar{b}j$  is estimated by replacing the  $b\bar{b}j$  Monte Carlo events with the 2Tag data and taking half the resulting shape difference as the uncertainty. Finally, the small shape difference between 3Tag and 2Tag data in the turn-on of the trigger level  $b$  tag is accounted for as a systematic uncertainty.

The modified frequentist method [17] is used to estimate  $1 - CL_b$ , the probability for a sample of background only




FIG. 2 (color online). Invariant mass for the high-mass likelihood region for the exclusive (a) three-jet (b) four-jet, and (c) five-jet channels. Black crosses refer to data, the solid line shows the total background estimate, and the shaded region represents the heavy flavor component ( $b\bar{b}b$ ,  $b\bar{b}c$ , and  $c\bar{c}b$ ). The lower panels show the difference between the data and the background expectation. Only statistical errors are shown.

to appear more signal-like than the observed data, as well as to derive limits at the 95% confidence level on the cross section times branching ratio as a function of  $m_A$ . As the absolute level of the multijet background cannot be reliably determined, only the shapes (not the normalization) of the  $M_{bb}$  distributions are used in the limit setting.

The systematic uncertainties on the signal and on the background shape are included in the limit calculation. Table I shows the limits and the  $1 - CL_b$  values obtained versus the hypothesized Higgs boson mass, assuming the width of  $\phi$  to be negligible relative to the experimental resolution ( $\approx 20\%$ ). The low  $1 - CL_b$  values around a Higgs boson mass of 180  $\text{GeV}/c^2$  are due to a slight excess over the expected SM background.

The results of this search can be used to set limits on the parameters of the MSSM. As a consequence of the enhanced couplings to  $b$  quarks at large  $\tan\beta$  the total width of the neutral Higgs bosons also increases with  $\tan\beta$ . This

can have an impact on our search if the width is comparable to or larger than the experimental resolution of the reconstructed invariant mass of a di-jet system. To take this effect into account, the width of the Higgs boson is calculated with FEYNHIGGS [18] and included in the simulation as a function of the mass and  $\tan\beta$  by convoluting a relativistic Breit-Wigner function with the next-to-leading order cross section. In the MSSM the masses and couplings of the Higgs bosons depend, in addition to  $\tan\beta$  and  $m_A$ , on the SUSY parameters through radiative corrections. Limits on  $\tan\beta$  as a function of  $m_A$  are derived for two particular scenarios assuming a  $CP$ -conserving Higgs sector [19]: the  $m_h^{\text{max}}$  scenario [20] and the no-mixing scenario [21]. Since the results depend considerably upon the Higgs sector bilinear coupling  $\mu$ , its two possible signs are also probed.

Figure 3 shows the results obtained in the present analysis interpreted in these different MSSM scenarios. Substantial areas in the MSSM parameter phase space up to masses of 200  $\text{GeV}/c^2$  are excluded. No exclusion can be obtained for the  $m_h^{\text{max}}$ ,  $\mu > 0$  scenario, due to the

TABLE I. Cross section limits as a function of Higgs boson mass. Columns two and three show the expected and observed limits on the cross section times branching fraction to  $b\bar{b}$ . The total one-sigma uncertainty on the expected limits is also displayed. The last column shows the value of  $1 - CL_b$ .

Mass ( $\text{GeV}/c^2$ )	$\sigma \times \mathcal{B}$ Expected (pb)	$\sigma \times \mathcal{B}$ Observed (pb)	$1 - CL_b$ (in %)
90	$170^{+72}_{-52}$	184	39
100	$117^{+48}_{-35}$	128	38
110	$71^{+29}_{-20}$	69	52
120	$41^{+18}_{-9}$	34	73
130	$28^{+12}_{-7}$	24	70
140	$25^{+11}_{-6}$	22	60
160	$17^{+8}_{-4}$	26	12
180	$13^{+5}_{-4}$	23	4.4
200	$9^{+4}_{-3}$	17	7.0
220	$7^{+3}_{-2}$	12	12




FIG. 3 (color online). 95% C.L. exclusion limits in the  $(m_A, \tan\beta)$  plane for  $m_h^{\text{max}}$ ,  $\mu = -200$  GeV, and no-mixing,  $\mu = -200$  GeV and  $\mu = +200$  GeV. The exclusions from LEP are also shown [2]. The width of  $\phi$  is larger than 70% of  $m_A$  above  $\tan\beta = 100$  in the  $m_h^{\text{max}}$ ,  $\mu = -200$  GeV scenario.

decrease in  $\sigma \times \mathcal{B}$  for positive values of  $\mu$  [19]. With the restriction of only using the shape of the discriminant variable, this analysis puts stringent limits in the  $(m_A, \tan\beta)$  plane up to Higgs boson masses of 220 GeV/ $c^2$ , particularly in the  $m_h^{\max}$ ,  $\mu < 0$  scenario.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC (United Kingdom); MSM and GACR (Czech Republic); CRC Program, CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); and the Alexander von Humboldt Foundation.

---

\*Visitor from Augustana College, Sioux Falls, SD, USA.

†Visitor from The University of Liverpool, Liverpool, UK.

‡Visitor from ICN-UNAM, Mexico City, Mexico.

§Visitor from II. Physikalisches Institut, Georg-August-University, Göttingen, Germany.

||Visitor from Helsinki Institute of Physics, Helsinki, Finland.

¶Visitor from Universität Zürich, Zürich, Switzerland.

\*\*Deceased.

- [1] H. P. Nilles, Phys. Rep. **110**, 1 (1984); H. E. Haber and G. L. Kane, *ibid.* **117**, 75 (1985).
- [2] S. Schael *et al.* (The ALEPH, DELPHI, L3, and OPAL Collaborations), Eur. Phys. J. C **47**, 547 (2006).
- [3] T. Affolder *et al.* (CDF Collaboration), Phys. Rev. Lett. **86**, 4472 (2001).
- [4] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **96**, 011802 (2006).
- [5] V. M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **95**, 151801 (2005).
- [6] V. M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **101**, 071804 (2008).
- [7] V. M. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **565**, 463 (2006).
- [8] T. Andeen *et al.*, Report No. FERMILAB-TM-2365, 2007.
- [9] T. Sjöstrand *et al.*, arXiv:hep-ph/0308153; We used PYTHIA version 6.323.
- [10] J. Campbell, R. K. Ellis, F. Maltoni, and S. Willenbrock, Phys. Rev. D **67**, 095002 (2003).
- [11] M. L. Mangano *et al.*, J. High Energy Phys. 07 (2003) 001.
- [12] R. Brun and F. Carminati, CERN Program Library Long Writeup W5013 (1993). GEANT3 was used.
- [13] G. Blazey *et al.*, arXiv:hep-ex/0005012.
- [14] V. M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **101**, 062001 (2008).
- [15] T. Scanlon, Report No. FERMILAB-THESIS-2006-43.
- [16] V. Barger, J. Ohnemus, and R. J. Phillips, Phys. Rev. D **48**, R3953 (1993).
- [17] T. Junk, Nucl. Instrum. Methods Phys. Res., Sect. A **434**, 435 (1999); A. Read, *ibid.* **425**, 357 (1999).
- [18] S. Heinemeyer, W. Hollik, and G. Weiglein, Eur. Phys. J. C **9**, 343 (1999); Comput. Phys. Commun. **124**, 76 (2000); G. Degrandi *et al.*, Eur. Phys. J. C **28**, 133 (2003); M. Frank *et al.*, J. High Energy Phys. 02 (2007) 047. We used FEYNHIGGS version 2.6.3.
- [19] M. Carena, S. Heinemeyer, C. E. M. Wagner, and G. Weiglein, Eur. Phys. J. C **45**, 797 (2006).
- [20]  $M_{\text{SUSY}} = 1$  TeV,  $X_t = 2$  TeV,  $M_2 = 0.2$  TeV  $|\mu| = 0.2$  TeV, and  $m_g = 0.8$  TeV.
- [21]  $M_{\text{SUSY}} = 2$  TeV,  $X_t = 0$  TeV,  $M_2 = 0.2$  TeV  $|\mu| = 0.2$  TeV, and  $m_g = 1.6$  TeV.