

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

U.S. Environmental Protection Agency Papers

U.S. Environmental Protection Agency

---

2008

## Fifteen Years after “Wingspread”—Environmental Endocrine Disrupters and Human and Wildlife Health: Where We are Today and Where We Need to Go

Andrew K. Hotchkiss

*USEPA/NCSU Cooperative Training agreement*

Cynthia V. Rider

*USEPA/NCSU Cooperative Training agreement*

Chad R. Blystone

*USEPA/NCSU Cooperative Training agreement*

Vickie S. Wilson

*Reproductive Toxicology Division*

Phillip C. Hartig

*Reproductive Toxicology Division*

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/usepapapers>

---

Hotchkiss, Andrew K.; Rider, Cynthia V.; Blystone, Chad R.; Wilson, Vickie S.; Hartig, Phillip C.; Ankley, Gerald T.; Foster, Paul M.; Gray, Clark L.; and Gray, L. Earl, "Fifteen Years after “Wingspread”—Environmental Endocrine Disrupters and Human and Wildlife Health: Where We are Today and Where We Need to Go" (2008). *U.S. Environmental Protection Agency Papers*. 188.  
<https://digitalcommons.unl.edu/usepapapers/188>

This Article is brought to you for free and open access by the U.S. Environmental Protection Agency at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in U.S. Environmental Protection Agency Papers by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

## Authors

Andrew K. Hotchkiss, Cynthia V. Rider, Chad R. Blystone, Vickie S. Wilson, Phillip C. Hartig, Gerald T. Ankley, Paul M. Foster, Clark L. Gray, and L. Earl Gray

## REVIEW

# Fifteen Years after “Wingspread”—Environmental Endocrine Disruptors and Human and Wildlife Health: Where We are Today and Where We Need to Go

Andrew K. Hotchkiss,<sup>\*,†</sup> Cynthia V. Rider,<sup>\*,†</sup> Chad R. Blystone,<sup>\*,†</sup> Vickie S. Wilson,<sup>†</sup> Phillip C. Hartig,<sup>†</sup> Gerald T. Ankley,<sup>‡</sup> Paul M. Foster,<sup>§</sup> Clark L. Gray,<sup>¶</sup> and L. Earl Gray<sup>†,1</sup>

<sup>\*</sup>USEPA/NCSU Cooperative Training agreement (CT826512010), Raleigh, North Carolina 27695; <sup>†</sup>Reproductive Toxicology Division, Endocrinology Branch, MD 72, National Health and Environmental Effects Research Laboratory, US Environmental Protection Agency, Research Triangle Park, North Carolina 27711; <sup>‡</sup>Mid-continent Division, Toxic Effects Characterization Research Branch, National Health and Environmental Effects Research Laboratory, US Environmental Protection Agency, Duluth, Minnesota 55804; <sup>§</sup>Toxicology Operations Branch, NIH/NIEHS, Research Triangle Park, North Carolina 27709; and <sup>¶</sup>Geography Department, University of North Carolina at Chapel Hill, North Carolina

Received December 22, 2007; accepted February 8, 2008

In 1991, a group of expert scientists at a Wingspread work session on endocrine-disrupting chemicals (EDCs) concluded that “Many compounds introduced into the environment by human activity are capable of disrupting the endocrine system of animals, including fish, wildlife, and humans. Endocrine disruption can be profound because of the crucial role hormones play in controlling development.” Since that time, there have been numerous documented examples of adverse effects of EDCs in invertebrates, fish, wildlife, domestic animals, and humans. Hormonal systems can be disrupted by numerous different anthropogenic chemicals including antiandrogens, androgens, estrogens, AhR agonists, inhibitors of steroid hormone synthesis, antithyroid substances, and retinoid agonists. In addition, pathways and targets for endocrine disruption extend beyond the traditional estrogen/androgen/thyroid receptor-mediated reproductive and developmental systems. For example, scientists have expressed concern about the potential role of EDCs in increasing trends in early puberty in girls, obesity and type II diabetes in the United States and other populations. New concerns include complex endocrine alterations induced by mixtures of chemicals, an issue broadened due to the growing awareness that EDCs present in the environment include a variety of potent human and veterinary pharmaceutical products, personal care products, nutraceuticals and phytosterols. In this review we (1) address what have we learned about the effects of EDCs on fish, wildlife, and human

health, (2) discuss representative animal studies on (anti)androgens, estrogens and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin-like chemicals, and (3) evaluate regulatory proposals being considered for screening and testing these chemicals.

**Key Words:** endocrine disruptors; androgens; estrogens; dioxins; PCBs; Pharmaceuticals; Mixtures; Screening and Multigenerational Testing.

It has been approximately 15 years since the first World Wildlife Federation (WWF) Wingspread Conference focused on endocrine-disrupting chemicals (EDCs), and 10 years since the U.S. Environmental Protection Agency (USEPA) was given a mandate under the Food Quality Protection and Safe Drinking Water Acts to develop test protocols to screen for endocrine effects of chemicals. This review article will briefly highlight some of the major scientific events that have taken place since the inception of the EDCs issue. In this review, we will (1) discuss EDCs effects on invertebrates, fish, wildlife, and human health, (2) highlight some representative studies on (anti)androgens, estrogens and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD)-like chemicals, and (3) evaluate regulatory proposals being considered for screening and testing these chemicals.

An initial impetus for studies of EDCs arose from the Wingspread work session in 1991 on “Chemically Induced Alterations in Sexual Development: The Wildlife /Human Connection” (Colborn and Clement, 1992). A consensus from that meeting, reached by a panel of expert scientists was that “Many compounds introduced into the environment by human activity are capable of disrupting the endocrine system of animals, including fish, wildlife, and humans. Endocrine disruption can be profound because of the crucial role

<sup>1</sup> To whom correspondence should be addressed. Fax: (919) 541-4017. E-mail: gray.earl@epa.gov.

Disclaimer: The research described in the article has been reviewed by the National Health and Environmental Effects Laboratory, US Environmental Protection Agency, and approved for publication. Approval does not signify that the contents necessarily reflect the views and policies of the agency nor does mention of trade names or commercial products constitute endorsement or recommendation for use. This research was supported [in part] by the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences.

hormones play in controlling development” (Colborn and Clement, 1992). These scientists also “estimated with confidence” that developmental impairments in humans have resulted from exposure to EDCs that are present in the environment as a result of human activities. Within a 5-year span several international organizations held meetings to discuss the EDC issue and research objectives. For example, starting in 1996 the USEPA held a series of workshops highlighting the new EDC issue and research needs in collaboration with the WWF and Chemical Manufacturers Association (Ankley and Giesy, 1998; Ankley *et al.*, 1998; DeVito *et al.*, 1999; Gray *et al.*, 1997a). The European Union (Weybridge Report, European Commission, 1996), Australia, and several Asian countries including Korea, Japan held similar meetings.

Subsequent studies have strengthened the documentation of effects of EDCs on animals. Although most of the effects in humans are due to occupational exposures or pharmaceutical usage, hundreds if not thousands of publications showing associations between background EDC exposures and adverse effects in humans are now available in the peer-reviewed literature. Many of the associations between EDCs and human health effects are controversial, but others, like the effects of polychlorinated biphenyls (PCBs) on neurological and immune function are quite widely accepted. Furthermore, animal laboratory studies corroborate many of these adverse effects observed in the field and, in some cases, provide mechanisms to explain the effects.

About 10 years ago the discussion of “endocrine disrupters” broadened from a focus on environmental estrogens to include additional mechanisms of endocrine toxicity; mechanisms that in many cases may be of equal or greater concern than estrogens. Numerous “new” areas of interest and concern have arisen since the original 1991 Wingspread conference and regulatory mandates. These areas include processes potentially disrupted by new classes of anthropogenic chemicals that act as antiandrogens, androgens, inhibitors of steroid hormone synthesis, antithyroid substances, and retinoid agonists. Within the last few years, scientists also have expressed concern about the potential role of EDCs in increasing trends in obesity and type II diabetes in the United States and other populations. In addition, although research has documented the complexity of the multiple mechanisms by which a single chemical can alter the endocrine milieu, recent evidence demonstrates the need to investigate complex endocrine alterations induced by mixtures of chemicals. We are now not only concerned about pesticides and other toxic substances in the environment, but the issue has broadened considerably due to the growing awareness that the list of EDCs present in the environment from human activities includes potent human and veterinary pharmaceutical products, personal care products, nutraceuticals and phytosterols. Among the drugs found as pharmaceuticals in the environment (PiEs) are potent, long lasting estrogens, antibiotics,  $\beta$ -blockers, antiepileptics, androgenic steroids, and lipid regulating agents. Some PiEs have been linked to dramatic effects in wildlife,

including death in bald eagles (<http://www.fws.gov/southeast/news/2002/12-03SecPoisoningFactSheet.pdf>), threatened extinction of species of vultures in Asia (Shultz *et al.*, 2004) and sex reversal and infertility in several species of fish (WHO, 2002.).

The 1996 legislation mandated that the USEPA both establish validated screening and testing procedures for estrogens and other substances as deemed appropriate and consider combinations of chemicals rather than evaluate the potential risk on a chemical by chemical basis. Both of these mandates encompass new areas of investigation in the EDC field. Furthermore, reports of nonmonotonic (e.g., U-shaped) dose–response relationships, ultra-low dose effects and non-threshold effects for EDCs continue to challenge some of the basic assumptions of toxicology and risk assessment. Although the focus of this debate has centered on the low dose effects of Bisphenol A, U-shaped dose–response curves are well known from many *in vitro* and a few *in vivo* studies. It appears that the basic tenet of toxicology from Ames and Gold (2000) that “dose alone determines the poison” is too limited for some EDCs because the timing of exposure can dictate not only the effect, but also whether the effects are adverse versus beneficial, or permanent versus transient. *In vivo*, however, the U-shaped effects of EDCs are generally limited to one or two effects whereas all other effects display “normal” non-monotonic dose responses. Furthermore, EDCs that induce cellular and molecular alterations of endocrine function at low dosage levels produce a cascade of effects increasing in severity resulting in adverse alterations at higher doses. The effects in the high dose groups may be different from those seen at low doses and the low dose effects are often causally linked to the high dose adverse effects of the chemical. This is true for androgens, natural estrogens, xenoestrogens, and antiandrogens.

The present review will focus on some of the new scientific issues in regard to EDCs that have surfaced since the initial Wingspread conference. The review will focus on effects of endocrine-active chemicals on the androgen, estrogen, and AhR-mediated signaling pathways. Reviews of other hormonal pathways potentially affected by EDCs, such as the thyroid system, can be found elsewhere (Guillette, 2006; Howdeshell, 2002; Tabb and Blumberg, 2006; Zoeller and Crofton, 2005).

## EDC EFFECTS ON FISH AND WILDLIFE, DOMESTIC ANIMALS, AND HUMAN HEALTH

### Invertebrates, Fish, and Wildlife

Endocrine-disrupting compounds have had dramatic effects on some invertebrate, fish, and wildlife populations. The diversity in life histories and habitats of animals in the environment introduce the possibility of species or life-stage-specific responses to EDCs that may not be evident in controlled laboratory experiments. Chemicals including

organotins, organochlorine pesticides, PCBs, and steroidal androgens and estrogens have all been associated with endocrine-mediated dysfunction in fish and wildlife populations. Wildlife studies can identify emerging contaminant problems, as well as inform about mechanisms of toxicity from laboratory work.

Great strides have been made in establishing cause and effect linkages between EDC exposures and observed effects in wildlife despite the inherent challenges. The complexity of environmental exposures often precludes identification of a single chemical responsible for observed population level effects. Additionally, it can be difficult to identify the underlying mechanism of the effect. These obstacles can be overcome by using a weight-of-evidence approach to assess the entire body of work surrounding observed wildlife effects.

Criteria have been developed for assessing the strength of the causal linkage between contaminant exposure and population level effect (Ankley *et al.*, 1998; WHO, 2002). These criteria assess the cause–effect connection based on concepts of temporality, strength of association, consistency of observation, biological plausibility, and evidence of recovery (WHO, 2002). There are several cases where strong evidence supports a causal linkage between exposure and effect (Table 1). Two examples discussed in detail below are organotin-mediated imposex in marine snails and feminization of sewage effluent-exposed male fish.

Exposure to the biocide tributyltin (TBT) causes imposex, or pseudohermaphroditism in female prosobranch gastropods (for reviews see, Ankley and Giesy, 1998; WHO, 2002). Imposex is the imposition of male sex organs, including penis and vas deferens, onto female snails and can lead to reproductive failure in some species (Horiguchi, 2006). It has been documented worldwide in around 150 species (Horiguchi, 2006). Due to these effects, use of TBT was restricted during the 1990's leading to subsequent recovery in multiple marine snail populations (Jorundsdottir *et al.*, 2005). High levels of TBT, however, are still found in some aquatic ecosystems because all usages have not been banned. The observation of imposex in these organisms has been associated in some studies with increased titers of testosterone in snail tissue (Spooner *et al.*, 1991). Although there are several competing hypotheses for this effect, including aromatase inhibition, altered metabolism of testosterone, and disruption of neuro-peptide signaling; a definitive mechanism of action has yet to be fully confirmed (Horiguchi, 2006; Sternberg *et al.*, 2008). New studies indicate that TBT causes imposex in invertebrates by acting as a retinoid X receptor (RXR) agonist, and displays an affinity for both invertebrate and vertebrate retinoid receptors (Nishikawa *et al.*, 2004).

Exposure to sewage effluent has been associated with induction of vitellogenin in male fish and the occurrence of intersex in wild fish (Jobling *et al.*, 2002a, b). Vitellogenin is an estrogen-responsive egg yolk protein precursor not normally expressed in male fish. A gradient of effect exists with fish

closest to the sewage outflow demonstrating the most severe responses (Harries *et al.*, 1999). Initial attention on feminized fish focused primarily on studies from the United Kingdom, however, the phenomenon seems to be relatively global in nature (Ankley *et al.*, 2007; WHO, 2002). For example, in the United States, the Potomac River has been identified as a hot spot for feminized smallmouth bass (Chambers and Leiker, 2006).

Caged fish and laboratory-based exposures confirm that sewage effluent is responsible for the observed increases in vitellogenin (WHO, 2002) and reproductive failure (Martinovic *et al.*, 2007). Although no single chemical has been identified as the culprit, chemical fractionation studies of sewage effluent have shown that synthetic and natural estrogens are often present in biologically relevant quantities (Desbrow *et al.*, 1998; WHO, 2002). These estrogenic compounds include synthetic pharmaceuticals and natural hormones in wastewater, such as ethinyl estradiol and 17 $\beta$ -estradiol. Various laboratory studies have shown that exposure to relatively low levels of these steroids will induce vitellogenin and can cause the development of intersex gonads (ovotestis) in fish and amphibians (Hutchinson *et al.*, 2006). Furthermore, a recent study by Kidd *et al.* (2007) showed that levels of ethinyl estradiol sufficient to cause vitellogenin induction in fish also caused a substantial decrease in the sustainability of the wild fish populations.

In some instances, there is moderate evidence for a causal linkage between EDC exposure and effects in wildlife; for example, PCB-induced reproductive and immune dysfunction in Baltic seals (Table 1). Great Lakes trout populations were adversely affected by PCB and dioxin exposures in the 1960–1970's, PCB-exposed cormorants displayed crossed-bills, great blue herons were infertile, and mink fed PCB contaminated fish from the Great Lakes either died after high exposures or were infertile at lower exposure levels. Proposed associations, which remain weak and also require elucidation, include reproductive and developmental anomalies in frogs exposed to atrazine and cryptorchidism in Florida panthers (Table 1).

### Domestic Animals

Naturally occurring compounds with estrogenic and other endocrine activities are widespread in nature. Over 400 species of plants contain known estrogens such as isoflavonoids or coumestans, or are suspected of being estrogenic based on biological grounds. Plants also may contain other compounds, in addition to estrogens, that can affect reproductive performance, such as antiandrogens in oil from saw palmetto (Gray *et al.*, 2001a).

Some plant estrogens occur at sufficient concentrations to cause reproductive alterations in domestic animals. “Clover disease,” which is characterized by dystocia, prolapse of the uterus and infertility, is observed in sheep grazed on clover

TABLE 1

Field and Lab Criteria for Determining a Causal Linkage between Endocrine Disruptors and Wildlife Effects (Ankley and Giesy, 1998; Gray *et al.*, 1998; WHO, 2002)

Animal	Notable effect	Chemical(s)	Field criteria						Lab criteria		
			Exposure–effect correlation	Effect involved in multiple populations	Specific effect(s) observed in exposed population	Exposure correlates with effect onset	Correlation of effect with exposure overtime	Correlation of decrease in effect with exposure remediation	Replication of field effects in lab species with relevant concentrations	Identification of mechanism of action	Replication of effect with like-acting chemicals
Marine snails	Imposex	TBT	+	+	+	+	+	+	+	+/-	+
Predatory birds	Egg shell thinning	<i>p,p'</i> -1,1-Dichloro-2,2-bis (p-chlorophenyl) ethylene (DDE)	+	+	+	+	+	+	+	+/-	+
Fish	Feminization	Sewage effluent	+	+	+		+	+	+	+	+
Fish	Masculinization	Pulp/paper mill effluent	+	+	+		+	+	+	+	+
Lake Trout	Low hatchability of fertilized eggs	TCDD-like compounds	+	+	+	+	+	+	+	+/-	+
Mink/otter	Reproductive failure	PCBs	+	+	+	+	+	+	+	+/-	+
Seals	Reproductive failure	PCBs	+	+	+	+	+		+	+/-	
Seals	Immune dysfunction	PCBs	+	+	+	+	+		+	+/-	
Alligator	Demasculinization	<i>p,p'</i> -DDE and other organochlorine pesticides	+	+	+				+/-		
Frogs	Hermaphroditism/Demasculinization	Atrazine	+/-	+	+				+		+
Polar bears	Demasculinization	PCBs	+		+				+		
Florida panthers	Cryptorchidism	Agricultural chemicals	+		+				+		

Note. + indicates that data support the listed criteria; +/- indicate that some data support the listed criteria, but more work is needed to confirm.



containing potent plant estrogens. Permanent infertility can be produced in ewes by much lower amounts of estrogen over a longer time period than are needed to produce clover disease (Adams, 1995).

In domestic animals, feeds contaminated with the zearalenone-producing fungus (*Fusarium* spp.) induce adverse reproductive effects in a wide variety of domestic animals, including impaired fertility in cows and hyperestrogenism in swine (Meyer *et al.*, 2000) and turkeys (Gray *et al.*, 2001a).

## Humans

Several chemicals are known to directly affect the endocrine system by altering development or reproduction in humans. Although other chemicals affect the reproductive system and indirectly alter endocrine function resulting in endocrine disruption and disruption of other components of the reproductive system. The examples of anthropogenic chemicals known to interact with the human endocrine system discussed herein includes accidental exposures to pesticides, toxic substances and pharmaceuticals with adverse side effects. In addition, we also discuss a few cases where pesticides or their metabolites and plant derived chemicals have been found to possess sufficient endocrine activity to be considered for use as pharmaceuticals, although no adverse effects have been associated with use of these chemicals. Some of the chemicals discussed are hormone agonists or antagonists or directly alter hormone synthesis. Others affect multiple cell types in the reproductive system including those like the Sertoli cell, cells that have both endocrine (secretes several hormones including antimüllerian hormone and inhibin) and nonendocrine functions (support germ cell development).

### Reproductive Effects of Chemicals in Humans Resulting from Occupational or Accidental Exposures or Pharmacological Side Effects

Accidental, occupational and therapeutic exposures to EDCs have produced adverse effects in humans (e.g., effects seen with PCBs (AhR and non-AhR pathways), TCDD, 2-bromopropane (likely affects the endocrine system indirectly), 4,4'-diaminostilbene-2,2'-disulfonic acid (DAS) (possible estrogenic mechanism), diethylstilbestrol (DES, estrogenic mechanism), the androgenic drugs danazol and methyltestosterone, and aminoglutethimide (drug that was taken that inhibits aromatase directly) (reviewed in Gray *et al.*, 2001a, b).

### Developmental Reproductive Toxicants Acting via an Endocrine Mechanism

#### *In Utero*

In a very well-documented example of EDC effects in humans, *in utero* exposure to the estrogenic chemical DES

alters reproductive development. Specifically, DES taken during pregnancy to prevent premature birth, which it failed to do, caused developmental alterations in the daughters including clear cell adenocarcinoma of the vagina, and gross structural abnormalities of the cervix, uterus, and fallopian tube. Women who were exposed to DES *in utero* are more likely to have spontaneous abortions, ectopic pregnancies and premature births (Steinberger and Lloyd, 1985). Developmental DES exposure has been reported to cause women to have less well-established sex-partner relationships; reduced sexual desire, excitability, and coital functioning, and an increased masculine pattern of cerebral lateralization on a verbal task (reviewed by Gray *et al.*, 2001a). Although far less prevalent and certain (Joffe, 2003), there are reports that males are also affected by developmental exposure to DES, displaying underdevelopment or absence of the vas deferens, epididymis, and seminal vesicles and persistence of the Müllerian ducts (Steinberger and Lloyd, 1985). They also reported that DES causes epididymal cysts, hypotrophic testes and infertility, and, in some males reduced ejaculate volume and numbers of motile sperm (Steinberger and Lloyd, 1985).

*In utero* exposure to the androgenic drug, danazol, also is contraindicated during pregnancy. Brunskill (1992) reported that of 94 completed pregnancies, 37 resulted in the birth of normal males, 34 in nonvirilized females, and 23 in virilized females. Virilization occurred in a proportion of female fetuses with a pattern of clitoromegaly, fused labia and urogenital sinus formation. The abnormality has not been reported where danazol therapy had been discontinued before the eighth week of pregnancy. Although more common in the higher dosages, virilization was reported in one case with a 200 mg daily dosage. Similar effects have been reported for other androgenic drugs (Grumbach and Ducharme, 1960; Saunders, 1968). The anticonvulsant drug aminoglutethimide, which inhibits aromatase and the production of estrogens from androgens, has also been associated with pseudohermaphroditism in daughters (LeMaire *et al.*, 1972).

#### *Pubertal*

In another more recent example of unintended effects of EDCs in humans, estrogen-containing shampoos (Tiwary, 1998) and skin oils (Henley *et al.*, 2007) were shown to be causative factors for pseudoprecocious puberty in girls and gynecomastia in boys (Felner and White, 2000). Some of the shampoos, for example, contained up to 4 mg of estradiol per 100 g. Others contained up to 2 g of the much less potent estril per 100 g (Tiwary, 1998).

Virilization in young children as young as 17 months of age is uncommon and is produced by androgens (Kunz *et al.*, 2004), which may be from endogenous or exogenous sources. Recently, the increased availability and use of commercial androgen products for cutaneous application has increased the

risk of virilization in children through skin contact and passive absorption.

### Reproductive Toxicity in Adults Resulting from an Endocrine Mechanism

Clinical studies by The National Institute of Occupation Safety and Health documented sexual impotence in chemical factory workers exposed to DAS, a DES-like stilbene derivative. These studies were carried out in response to complaints of impotence and decreased libido among male workers involved in the manufacture of DAS, a key ingredient in the synthesis of dyes and fluorescent whitening agents. The data from the studies showed that workers manufacturing DAS had lower serum testosterone levels and reduced libido (Grajewski *et al.*, 1996; Whelan *et al.*, 1996) as compared with control workers. They found that “Current and former DAS workers had lower mean total testosterone (TT) levels compared with additives workers (458 and 442, respectively, vs. 556 ng/dl;  $p = 0.05$  and  $0.04$ ). Current and former DAS workers were 3.6 (95% CI, 0.5–24.4) and 2.2 (95% CI, 0.3–18.0) times more likely than additives workers to have lowest quartile TT levels ( $< 386$  ng/dl) after adjustment for age and body mass index. Duration of employment in DAS production was negatively related to the workers’ testosterone levels.” Studies in rodents suggest that DAS has estrogenic activity, possibly providing a possible mechanistic explanation for the complaints of impotency in factory workers exposed to these substances (Smith and Quinn, 1992).

The *o,p'*-dichlorodiphenyltrichloroethane (DDT) metabolite, *o,p'*-Ortho,para,dichlorodiphenyl dichloroethane (DDD) has historically been considered as an environmental contaminant. However, *o,p'*-DDD also has clinical uses due to its ability to alter endocrine function and it has been reported to induce both beneficial and adverse side effects. *o,p'*-DDD (Mitotane), has been used to treat adrenal steroid hypersecretion associated with adrenal tumors in humans and pets (Reine, 2007). It has been reported to act by inhibiting P450 steroidogenic enzymes and also has some estrogenic activity in humans (Nader *et al.*, 2006). In addition, mitotane treatment restored menstruation in women with high androgen levels of spanomenorrhea associated with hypertrichosis with pregnancies occurring in about a third of the treated women (Hayes and Laws, 1991). However, Mitotane treatment has been associated with atrophy of the spermatogenic tubules of the testis in men (Sparagana, 1987). Sparagana studied a patient who developed impotency due to primary testicular failure at the time that he was treated with Mitotane. A testicular biopsy, performed after the drug was discontinued, showed normal appearing Leydig cells and atrophy of the seminiferous tubules with the picture of a maturation arrest. In the four and one half years since he last received mitotane, the patient’s libido has slowly improved and his plasma testosterone, gonadotropins and LH response to gonadotropin-releasing hormone have become essentially

normal. He proposed that mitotane was cytotoxic to the testis as it is to the adrenal cortex.

### Reproductive Toxicity in Adults that Indirectly Produces Endocrine Alterations

Yet another example of a chemical altering reproductive and endocrine function in humans is 2-bromopropane (CERHR, NTP, 2003) (Kim *et al.*, 1996). Korean electric workers exposed to solvents containing this chemical displayed reproductive and endocrine abnormalities. Specifically, women showed secondary amenorrhea, primary ovarian failure with high follicle-stimulating hormone levels, hot flashes and undetectable estradiol levels (below 13.6 pg/ml the LOD). Exposed men displayed azoospermia, some degree of oligospermia, or reduced sperm motility (Kim *et al.*, 1996) but had normal libido and based upon hormonal levels it appears that the germ cells and not the Leydig cells were directly affected by 2-bromopropane (2003).

Dibromochloropropane (DBCP) is another human reproductive toxicant that indirectly disrupts endocrine function in humans (Lag *et al.*, 1989; Potashnik and Porath, 1995; Potashnik and Yanai-Inbar, 1987; Slutsky *et al.*, 1999; Whorton and Foliart, 1983; Whorton and Milby, 1980; Whorton *et al.*, 1979). The pesticide DBCP acts directly on the germ cells as a mutagen. Occupational exposures during manufacturing and application in several countries (Teitelbaum, 1999) resulted in permanent oligospermia, azoospermia, and infertility, and indirectly elevated FSH levels and non-significant reductions in testosterone. Unlike 2-bromopropane, above, few women were exposed to DBCP. In rodents, *in utero* DBCP severely disrupts fetal testis development (Warren *et al.*, 1988). Effects on sperm counts and FSH levels persist in some workers for at least 17 years. Despite the ban on the use of DBCP, this pesticide remains persistent in soil and continues to be detected as a groundwater contaminant in areas of past high use, in particular California’s Central Valley (Clark and Snedeker, 2005; Teitelbaum, 1999). Although case-control, cohort and ecological epidemiology studies have not found significant, positive associations between DBCP exposure and cancer in exposed populations, concerns persist due to the cited limitations in these studies (Clark and Snedeker, 2005).

### EDCs in Humans and Effects at Background Exposure Levels

In contrast to the above examples, background human EDC exposure-effect relationships have only occasionally approached a significant degree of certainty. There are literally thousands of studies on the potential effects of EDCs on human health. Although many of the have shown associations among EDC and reproductive alterations (sperm abnormalities, shortened anogenital distance in boys, altered sex ratio, testis



cancer, cryptorchidism, hypospadias, testes cancer, prostate cancer, advanced onset of puberty, breast cancer, endometriosis, obesity, attention deficit disorder, uterine cancer, and allergies) many others have failed to find such associations and, for this reason, there remains a great deal of uncertainty about the effects of background levels of EDCs on human reproduction. Although it is very important that scientists continue to evaluate the effects of background levels of EDCs on human disease, it is more important to prevent as opposed to detect the effects of EDCs on humans, fish, and wildlife. As stated by Lucier (2007; <http://domesticpolicy.oversight.house.gov/documents/20070925143750.pdf>), "We should keep in mind that a positive finding in an epidemiology or clinical study is, in reality, a failure of preventive medicine policy."

In contrast, the effects of the PCBs on human neurological and immune development, is one case where many scientists have concluded that the data are robust enough to conclude that they approach the level of cause and effect (reviewed in Brouwer *et al.*, 1999; Selgrade, 2007). Even here, there remains considerable uncertainty. In contrast to background levels of PCBs, children accidentally exposed to high levels of PCBs and dibenzofurans display neurobehavioral abnormalities from exposure *in utero* and during early postnatal development (Guo *et al.*, 1995). These changes included low birth weight, malformations, and reduced IQ and cognitive abilities. Furthermore, the exposed males exhibited abnormal sperm morphology, reduced sperm motility, and reduced *in vitro* sperm fertilizing capacity (Guo *et al.*, 1995). Selgrade (2007) recently concluded that "suppression of immune responses in rodents is predictive of suppression of immune responses in humans and that there is a relationship between immune suppression following developmental exposure to the toxicants and enhanced risk of infectious or neoplastic disease in humans" for PCBs as well as cigarette smoke and arsenic.

### Adverse Effects of EDCs in Laboratory Animals

#### *In Vivo Studies on Select Classes of EDCs*

The following section briefly reviews selected examples of the effects of EDCs on, mostly, rodent reproductive development. Although this review does not include a discussion of the *in vitro* and short-term studies examining the mode(s) of action of the selected chemicals, such studies are essential to confirm mechanisms of action hypothesized from *in vivo* studies. Specific EDC classes considered here include antiandrogens, androgens, estrogens and AhR agonists (such as TCDD).

*Environmental antiandrogens.* Several environmental chemicals adversely affect male development by interfering with androgen signaling during the critical periods of sexual differentiation and maturation. Depending on the timing of exposure (e.g., *in utero* or peripubertal), these antiandrogens affect androgen-sensitive organs and processes within the male

rat, leading to histological lesions within the reproductive organs, delayed puberty (Monosson *et al.*, 1999), hypospadias/cryptorchidism, reduced fertility, and testicular tumors. Mechanistic work has identified some antiandrogenic chemicals to be androgen receptor (AR) antagonists, e.g., the DDT metabolite *p,p'*-DDE (Kelce *et al.*, 1995), the fungicides vinclozolin (Gray *et al.*, 1994; Kelce *et al.*, 1997), procymidone (Gray *et al.*, 2006; Hosokawa *et al.*, 1993; Ostby *et al.*, 1999) and prochloraz, the herbicide linuron, and polybrominated diphenyl ether (PBDE) flame retardants (Stoker *et al.*, 2005).

Some chemicals can produce antiandrogenic effects via depression of testicular testosterone production. The fungicide prochloraz induces malformations in male offspring and delays the onset of puberty after peripubertal exposure, with mechanistic work revealing that it is not only an AR antagonist but also acts as a potent steroid synthesis inhibitor (Blystone *et al.*, 2007a, b; Noriega *et al.*, 2005; Wilson *et al.*, 2004c). Similarly, linuron, is both an AR antagonist (Lambright *et al.*, 2000) and it reduces testosterone production (Hotchkiss *et al.*, 2004; Wilson *et al.*, 2004c). The AR antagonism and reduced testosterone production displayed by these chemicals (Hotchkiss *et al.*, 2004; Wilson *et al.*, 2004c) likely has a cumulative effect on the male offspring, but the relative contribution of each mechanism requires further clarification.

Several phthalate esters, used as plasticizers in many commercial products, are antiandrogens, which gives cause for concern because there is widespread exposure to these chemicals (Silva *et al.*, 2004). Malformations of male reproductive organs occur after *in utero* exposure to some phthalates (e.g., dibutyl phthalate, butyl benzyl phthalate, DEHP, di-*iso*-nonyl phthalate) and, through a mechanism that has yet to be established, these chemicals reduce fetal rat testosterone production (Foster, 2006; Gray *et al.*, 2000; Mylchreest *et al.*, 1998; Parks *et al.*, 2000; Sharpe, 2006; Wilson *et al.*, 2004c). In addition to reduced testosterone, phthalate exposure also reduces *insl-3* gene expression (Wilson *et al.*, 2004c); the *insl-3* peptide hormone is necessary for transabdominal testis descent (Klonisch *et al.*, 2004).

The spectrum of malformations that result from disrupting the androgen signaling pathway varies depending on the mechanism of action. For example, hypospadias is prominent at high doses of vinclozolin (whose metabolites are AR antagonists), whereas high levels of phthalates (which reduce fetal testosterone and *insl-3* expression) induce epididymal agenesis. However, linuron and prochloraz, two chemicals that one would expect to display similar effects based upon similar dual mechanisms, differ in the relative rates of hypospadias and epididymal agenesis that they induce (Gray *et al.*, 2006). Together this area of EDC research has characterized the effects of several antiandrogens and identified relevant and diverse mechanisms of action pertinent to human health risk assessment because this pathway is highly conserved among vertebrates. Disruptions of this pathway either via genetic

mutations or steroid-like chemical exposures results in profound examples of pseudohermaphroditism in experimental animals and humans (Quigley *et al.*, 1995).

The reproductive system is also sensitive to phthalate exposure during the peripubertal and neonatal stages of life. Rats treated with diethyl hexyl phthalate (DEHP) orally or by injection as neonates display permanent alterations lesions of the testis and sperm numbers (Cammack *et al.*, 2003; Dostal *et al.*, 1988). When administered during peripubertal life, di-n-phthalate esters with side chains with three to six carbons (Foster *et al.*, 1981) disrupt testis structure and function, reduces testosterone production, delays pubertal onset and reduces androgen-dependent organ weights in male rats (Gray *et al.*, 1999). These effects are not limited to Murid and Cricetid rodents like the rat and hamster, respectively, and have been seen in several other mammalian species including the guinea pig (Gray *et al.*, 1982), ferret (Lake *et al.*, 1976) and rabbit (Higuchi *et al.*, 2003).

Sjoberg *et al.* (1986) found an age sensitivity in response with the adults more resistant to testicular damage than pubertal animals. When he administered DEHP in the diets of 25-, 40-, and 60-day-old male rats for 14 days, testis weight was reduced in males in 25-day-old males (at 1 g/kg/day) and 40 (at 1.7 g/kg/day) and all testis tubules were damaged in 25-day-old males (at 1.7 g/kg/day). No testicular effects were observed in 60-day-old animals (Sjoberg *et al.*, 1986).

**Environmental androgens.** It was noted in the 1970's that female fish of several species living in rivers contaminated with pulp mill effluent (PME) displayed masculinized sexual traits and starting in 2001 two different research teams reported that PME displayed androgenic activity in *in vitro* assays (Jenkins *et al.*, 2003; Parks *et al.*, 2001). Since then, androgenic activity has been detected in PME from other rivers in Florida, the Baltic Sea, the Great Lakes and New Zealand (Ellis *et al.*, 2003; Larsson and Forlin, 2002; Parks *et al.*, 2001). PME effluents from sites on the Fenholloway River in Florida include a chemical mixture that binds AR and induces androgen-dependent gene expression *in vitro*. This mode of action is consistent with the observation of masculinized female mosquitofish (*Gambusia holbrooki*) collected from contaminated sites on the river. Male-biased sex ratios of fish embryos have been reported near a pulp mill in broods of eelpout (*Zoarces viviparus*) in the vicinity of a large kraft pulp mill on the Swedish Baltic coast, suggesting that masculinizing compounds in the effluent were affecting gonadal differentiation and promoting skewed sex ratios. Although androgenic chemicals have been isolated from PME, efforts to date have not conclusively identified the chemicals in PME responsible for masculinization of the female fish (Durhan *et al.*, 2002).

Effluents from cattle concentrated animal feeding operations (CAFO) in Nebraska and Ohio have also been shown to display androgenicity. Orlando *et al.* (2004) found that water associated with a CAFO in Nebraska exhibited androgenic

activity and found that fish (fathead minnow; *Pimephales promelas*) collected at the site displayed small gonads, fewer embryos (Orlando *et al.*, 2007) and morphological differences compared with fish from a reference site. Durhan *et al.* (2006) detected the synthetic androgens 17 $\alpha$ -trenbolone and 17 $\beta$ -trenbolone in water samples associated with a beef CAFO in Ohio, where trenbolone acetate implants were used to stimulate weight gain; samples collected from a direct discharge from the feedlot displayed significant androgenic activity *in vitro*. Complementary laboratory studies revealed both trenbolone isomers were androgenic in the fathead minnow (Ankley *et al.*, 2003; Jensen *et al.*, 2006) and the rat (Hotchkiss *et al.*, 2007; Wilson *et al.*, 2002b). For example, *in utero* administration of 17 $\beta$ -trenbolone to dams resulted in masculinized female rat offspring, displaying increased anogenital distance (AGD), vaginal agenesis and induced male sex accessory tissues in females (Hotchkiss *et al.*, 2007; Wilson *et al.*, 2002b).

**Environmental estrogens.** Much of the interest in environmental estrogens grew out of reports of reproductive cancers, reproductive dysfunction, and potential behavioral alterations of women who had been developmentally exposed to the pharmaceutical estrogen, DES. Estrogens are critical for female reproduction both developmentally and in adulthood. Two different estrogen receptors ( $\alpha$ ,  $\beta$ ) have been characterized in mammals with 3 different receptors ( $\alpha$ ,  $\beta$ 1,  $\beta$ 2) identified in teleost fish (Hawkins *et al.*, 2005). Differential expression of these receptors in tissues as well as different binding affinities for various ligands suggests diversity in physiological function.

Studies with mammals have shown that estrogenic compounds are quite varied in structure and potency, including pharmaceuticals (ethinylestradiol, DES), insecticides (methoxychlor, DDT, Chlordane), surfactants (octylphenol, nonylphenol), phytoestrogens (genistein, coumesterol), sun screens, and plastics (bisphenol A) (Gray *et al.*, 2001a).

High dose oral *in utero* exposure to potent estrogens such as ethinyl estradiol induces reproductive tract malformations in rats with females affected at lower doses than male offspring. In contrast, estrogenic pesticides or toxic substances have not been shown to induce similar structural malformations. Perinatal oral maternal exposure to weaker estrogenic chemicals like methoxychlor defeminizes the central nervous system (CNS) of the female rat such that they are acyclic and infertile and have a shortened reproductive life span due to an early onset of irregular estrous cycles. Male offspring from such studies are fertile but occasionally display some effects on reproductive tissues at higher dosage levels. Although one laboratory has reported that ip injections of pregnant rats with high doses of methoxychlor during gonadal differentiation induces testicular lesions and reduced fertility in offspring, with such effects being transmitted through several generations (Anway *et al.*, 2005), other investigators have not observed similar testicular effects or reduced fertility in F1 males using oral exposures (a more relevant exposure route for risk assessment) during this period of development (Gray *et al.*, 1989).

Exposure of developing female rodents (rats and hamsters) to ethinyl estradiol, methoxychlor, chlordecone, and zearalenone, for example, can result in altered reproductive behaviors, sexually dimorphic nonreproductive behavior, neuroendocrine function (estrous cycles, onset of puberty), fertility, and reproductive tract morphology. Developing males are not immune to the adverse effects of exogenous estrogens. Studies with ethinyl estradiol in males showed altered testicular development among other effects (Howdeshell *et al.*, 2007).

Oral exposure to all potent and may xenoestrogens after weaning of female rats causes pseudoprecocious puberty (early vaginal opening), decreased food intake, and body weight, altered activity, and altered reproductive behavior in both males and females (Gray *et al.*, 1989; Gray and Ostby, 1998). Finally, there are ongoing discussions about the possibility of low dose effects of this class of chemicals, especially Bisphenol A, and this continues to be an active area of research and debate about the reproducibility and relevance of the low dose findings given the lack of adverse effects at any dose level (CERHR, NTP Bisphenol A Expert Panel Final Nov 26, 2007) (Guillette, 2006).

*Environmental AhR agonists.* Exposure to Ah receptor agonists such as TCDD, PCBs, and PCDFs has been causally linked to developmental/reproductive toxicity in humans (Brouwer *et al.*, 1999; Guo *et al.*, 1995), nonhuman primates (Rier *et al.*, 2001), rodents (see below), mink (Hornshaw *et al.*, 1983), fish (Ankley and Giesy, 1998), and other wildlife species (Ankley and Giesy, 1998). Agonists of the AhR (which is a cellular steroid hormone-like receptor) form a complex with the receptor that acts as a transcription factor by binding to specific DREs (dioxin response elements) on specific genes. Some, if not all of the toxicity of TCDD, appears to result from activation of the Ah receptor. TCDD is an “endocrine disrupter” that acts on multiple components of the endocrine axis. TCDD exposure alters the levels of many hormones and growth factors, and their receptors (see Part II, Draft EPA Dioxin Risk Assessment document: <http://www.epa.gov/ncea/pdfs/dioxin/part2/dritoc.pdf>).

In addition to the effects of these chemicals seen in humans from accidental exposures, exposure to very low doses of TCDD produces infertility in rodent progeny (Bjerke and Peterson, 1994; Bjerke *et al.*, 1994a, b; Mably *et al.*, 1992). Exposure to a single low dose of TCDD ranging from 50 ng to 2 µg/kg during sex differentiation of the rat or Syrian hamster results in a number of unusual reproductive alterations in male and female progeny (Gray and Ostby, 1995; Gray *et al.*, 1995, 1997b, c; Wolf *et al.*, 1999). In female rats, gestational exposure to TCDD induced clefting of the phallus with a mild degree of hypospadias in females and a permanent “thread” of tissue across the opening of the vagina of the progeny. Female progeny, treated earlier in gestation displayed reduced fecundity, a high incidence of constant estrus, and cystic endometrial hyperplasia at middle-age. Female hamsters,

exposed *in utero* with TCDD also displayed clitoral clefting, reduced fertility as a result of several functional reproductive problems, but they did not display the vaginal thread. In TCDD-exposed male rat and hamster offspring, puberty was delayed, ejaculated and epididymal sperm numbers were reduced, whereas the reductions in ventral prostate, seminal vesicle, and testis size, displayed during peripubertal life, were attenuated with age. When administered chronically prior to and throughout gestation, TCDD delays puberty in the male rat at doses as low as 2.4 ng/kg/day (Bell *et al.*, 2007).

Mating behavior was normal in male hamsters, whereas male rats had some difficulty achieving intromissions. No malformations were noted in male rat or hamster offspring at these dosage levels. Furthermore, a PCB congener, 169, is an Ah receptor agonist with a toxic equivalency factor of about 0.001, as compared with the potency of TCDD. PCB 169 treatment during pregnancy alters reproductive development of Long Evans hooded male and female rats in a manner almost identical to TCDD (Gray and Ostby, 1998).

TCDD and related compounds also have been linked to endometriosis in nonhuman primates (Rier *et al.*, 2001) and women (Louis *et al.*, 2005). Exposure to TCDD is associated with a dose-dependent increase in the incidence and severity of endometriosis in the rhesus monkey and it was reported that the serum levels of TCDD and specific dioxin-like PHAH congeners were increased in TCDD-treated rhesus monkeys with endometriosis 13 years after the TCDD exposure.

Other classes of vertebrates are adversely affected by low concentrations of TCDD and related compounds. These chemicals are toxic to some embryonic fish (Zabel *et al.*, 1995) and avian (Sanderson *et al.*, 1994) species in the field and in the laboratory.

## NEW AREAS OF RESEARCH

### *Other Pathways Affected by EDCs*

It has become apparent that endocrine disruption includes mechanisms and targets beyond the traditional estrogen/androgen/thyroid (EAT) receptor-mediated systems. Examples of “nontraditional” endocrine disruption include various inhibitors of the steroidogenic pathway including fetal Leydig cell androgen production and aromatase inhibitors (Gray *et al.*, 2006). The herbicide, atrazine, binds to a new class of membrane-bound progesterone receptors, inducing neuronal mast cell degranulation *in vitro* (Mizota and Ueda, 2006) and several EDCs have been reported to interact with membrane-bound estrogen receptors. However, the physiological role of these receptors and the biological significance of these *in vitro* observations have yet to be defined and for these reasons, binding of these receptors can only be hypothetically linked to adverse effects *in vivo*.

Recently, nitrates have come into focus as potential endocrine disruptors. Although historically linked to

methemoglobinemia, nitrates are capable of disrupting gonadal steroidogenesis and thyroid function (Guillette, 2006). In invertebrates, other receptor systems such as the retinoid receptor systems (RXR) and ecdysteroids (EcR) play critical roles in reproduction and growth (Rodriguez *et al.*, 2006).

Recent evidence also suggests that EDCs may disrupt terminal metabolism of endogenous and exogenous materials by altering the activating hepatic enzymes. For example, the pregnane X receptor in the liver is upregulated by numerous EDCs including nonylphenol, DDT, and methoxychlor in a way that alters the metabolism of various xenobiotics and endogenous hormones (Kretschmer and Baldwin, 2005). This altered metabolism may affect androgen and estrogen levels as well as the balance of bound and free thyroid ligands (Kretschmer and Baldwin, 2005).

In addition to novel mechanisms of action, nonreproductive targets of EDCs have been identified. These targets include the cardiovascular system, the digestive system, and adipose tissue (WHO, 2002). Immune function, long known to be sensitive to steroids, has also been identified as an EDC target (Inadera, 2006). Hormone receptors have been identified on various immune cells and cytokine receptors localized to various endocrine tissues and the brain. In addition, increasing incidence of autoimmune diseases, sex differences in immune function, and documented immunomodulatory effects of sex steroids have highlighted the potential for endocrine-active chemicals to affect this dynamic system. Examples of chemicals interfering with immune function via endocrine interactions have been described for numerous compounds including androgens (Hotchkiss and Nelson, 2007), estrogens, organotins, and dioxins (Inadera, 2006). Selgrade (2007) stated that there is now sufficient data, at least for the effects of toxicants on the developing immune system, to conclude that the concern raised about human risk by the rodent data is justified (Selgrade, 2007).

Another target of EDCs is the CNS. Not only are some sexually dimorphic behaviors affected permanently by exposure to chemicals with androgenic or estrogenic activities (Gray *et al.*, 2006) neural development also can be altered by exposure to thyroid-active EDCs (Schantz and Widholm, 2001). The hypothalamic-pituitary-adrenal axis also is a potential target of EDCs. Chemical activation of this system by PCBs or through interactions with the glucocorticoid receptors (Guillette, 2006; Johansson *et al.*, 1998) can have adverse effects on a number of different systems, thereby expanding the number of potential targets for EDCs.

### Mixtures

Mixtures of EDCs in the environment present a major emerging issue for research and risk assessment. Toxicological studies have typically focused on single chemical exposures and their effects; however, exposure to mixtures in the environment is the rule, not the exception (Kolpin *et al.*,

2002). It is essential to develop and validate methods to accurately predict effects of endocrine-disrupting mixtures in order to protect humans and wildlife from the risk associated with potentially cumulative effects of these mixtures.

The evidence from both *in vitro* and *in vivo* studies generally indicates that mixtures of EDCs with the same mechanism of action display additive effects, however, chemical mixtures also can act in an antagonistic or synergistic manner. Initially, studies focused on estrogenic chemicals. For example, *in vitro* studies using the yeast estrogen screen assay (Payne *et al.*, 2000), as well as the breast cancer cell proliferation assay (Payne *et al.*, 2001; Silva *et al.*, 2002) found that mixtures of xenoestrogens elicit additive activation of the estrogen receptor. Furthermore, Silva *et al.* (2002) showed that chemicals present at their no observed effect concentrations contribute to a cumulative effect of the mixture. *In vivo* studies by Brian *et al.* (2005) demonstrated that a mixture of five estrogenic compounds induced vitellogenin in male fish corresponding to predictions based on the concept of concentration addition.

An increasing incidence of male reproductive tract developmental abnormalities highlights the importance of assessing antiandrogenic mixtures for their potential effects (Skakkebaek *et al.*, 2001). Recent studies in our own lab have demonstrated that *in utero* exposure to mixtures of chemicals that target the androgen signaling pathway at multiple sites (e.g., AR antagonists and inhibitors of testosterone) elicit dose-additive effects on the male rat reproductive tract (Gray *et al.*, 2001, 2006; Hotchkiss *et al.*, 2004; Howdeshell *et al.*, 2007; Rider *et al.*, 2008). The fact that chemicals that act on different fetal tissues via diverse cellular mechanism of action produce additive effects indicates that the current framework for conducting cumulative risk assessments should not only consider including chemicals from different classes with the same "mechanism of toxicity", but also should include chemicals that disrupt differentiation of the same fetal tissue at different sites in the androgen signaling pathway.

*In vivo* studies (Birkhoj *et al.*, 2004; Gray *et al.*, 2001b, 2006; Hotchkiss *et al.*, 2004; Nellemann *et al.*, 2003) have repeatedly found that mixtures of AR antagonists elicit additive effects on young male rats exposed to the chemicals *in utero*. Dose-additive effects also have been demonstrated *in vitro* using AR activation assays (Birkhoj *et al.*, 2004; Nellemann *et al.*, 2003). It is likely, for example, that androgen-dependent tissues cannot discriminate in terms of cellular responses, between the blockade of the AR and significantly lowering the concentration of the ligand (T and/or dihydrotestosterone). It would be reasonable to expect, therefore, that both of these actions could produce dose-additive adverse responses in the differentiating tissues.

The finding that mixtures of chemicals targeting endocrine signaling at multiple levels adhere to dose additivity was also observed with mixtures of thyroid disrupting chemicals (Crofton *et al.*, 2005). Overall, current mixtures research



TABLE 2  
Strategy for Ongoing Studies with Vertebrate and Invertebrate Estrogen and ARs

	Invertebrate		Reptiles and amphibians			Fish		Bird	Mammal	
	Mud snail	Water flea	Northern Leopard Frog	Giant Salamander	American Alligator	Rainbow trout	Fathead minnow	Japanese Quail	Chimpanzee	Human
Obtain animal tissues	X	X	X	N/A	N/A	N/A	X	N/A	N/A	N/A
Prepare complimentary DNA library	X	X	X	N/A	N/A	N/A	X	N/A	N/A	N/A
Isolate ER or AR			Ongoing	N/A	N/A	N/A	ER $\alpha$ , AR	N/A	N/A	N/A
Sequence ER or AR				ER $\alpha^1$	ER $\alpha^1$	ER $\alpha^3$ , AR <sup>2</sup>	ER $\alpha$ , AR	ER $\alpha^1$	AR <sup>1</sup>	ER $\alpha^1$ , AR <sup>1</sup>
Express ER or AR				ER $\alpha$	ER $\alpha$	ER $\alpha$	ER $\alpha$ , AR	ER $\alpha$	AR	ER $\alpha$ , AR
ER or AR Function					Ongoing	AR	ER $\alpha$ , AR		AR	ER $\alpha$ , AR
Compare function across species						AR	ER $\alpha^4$ , AR		ARongoing	ER $\alpha^4$ , AR

Note. ER $\alpha^1$  = sequence obtained from GenBank. AR<sup>2</sup> = expression vector generously provided by Takeo and Yamashita (1999); ER $\alpha^3$  = expression vector generously provided by F. Pakdel (Pakdel *et al.*, 2000), ER $\alpha^4$  = manuscript in preparation, N/A = not applicable—sequence obtained from outside sources or GenBank. Mud snail = *Ilyanassa obsoleta*, Water flea = *Daphnia magna*, Northern Leopard Frog = *Rana pipiens*, Japanese Quail = *Coturnix japonica*, Giant salamander = *Andrias davidianus*, Rainbow trout = *Oncorhynchus mykiss*, fathead minnow = *Pimephales promelas*, American alligator = *Alligator mississippiensis*, Chimpanzee = *Pan troglodytes*.

points to two major conclusions: (1) individual chemicals present below their NOAELs contribute to overall mixture effects and (2) true risk associated with exposure to EDCs can only be determined when the cumulative effects of chemicals that target a common signaling pathway are taken into consideration.

### Cross-Species Comparisons

Assessment of the validity of cross-species extrapolation is problematic on several issues. Fish and wildlife are often viewed as “sentinel species.” Hence, estrogenic and androgenic effects in animals in the environment are generally assumed to be of concern to other species, including humans. However, preliminary cross-species studies suggest that surrogate laboratory species may not always be relevant to the species of concern and that data from human health risk assessments may not accurately reflect the risk to fish and wildlife. Current prescreening protocols only include mammalian receptor-based assays, and thus may not provide as accurate an assessment for fish and wildlife as would similar assays utilizing receptors obtained from nonmammalian species.

Assessing the impact of EDCs on fish and wildlife has been facilitated by molecular cloning and gene synthesis (Table 2; Wilson *et al.*, 2004). Generation of cDNA libraries from specific tissues can effectively immortalize and clone the genes of a species, frequently utilizing only one animal or surgical specimen. (Wilson *et al.*, 2004b). These libraries act as repositories for genes associated with endocrine function, as well as providing a source of material for future studies. Additionally, databases of gene sequences derived from

analysis of overlapping gene fragments provide information for the synthesis and expression of an entire gene without utilization of additional animals or tissues. Both approaches are useful in generating an unlimited source of receptor proteins in a controlled environment. These proteins can then be subjected to binding assays using similar protocols, to determine if EDCs affect steroid hormone receptors equally across species (Wilson *et al.*, 2007). These binding assays can utilize nonmammalian recombinant receptors, in various stages of purification, or expressed proteins in living cells. Both approaches provide valuable data with regard to ligand binding, but provide little information about how ligand–receptor–EDC interactions might affect gene regulation. Current mammalian transcriptional activation assays, utilizing mammalian receptor and receptor specific reporters in homologous cells lines, are more fully developed than for lower vertebrates. However, generation of homologous cell lines and species-specific reporter constructs would facilitate similar studies of alteration of gene induction in fish and wildlife species.

The utility of any species in addressing EDC questions is enhanced if species are from taxa not commonly studied, have the potential for laboratory and field studies, have been affected by EDCs in the wild and can help resolve conflicting reports of receptor binding. Studies of the fungicide vinclozolin and its metabolites have reinforced the validity of this approach to species selection. Despite the high degree of homology in the ligand binding domain of human and the fathead minnow AR, subtle differences in the binding affinities for these EDCs have been documented between mammals and fish, and suggest that across species, the impact of a given EDC may vary (Wilson *et al.*, 2004b, 2007). Similar studies comparing the estrogen receptor (ER) binding affinity of plasticizers across animal



classes have also documented differences, and established the need for additional studies assessing the effect of EDCs on other receptors (e.g., the estrogen receptor) of diverse species (Rider *et al.*, in press). This is needed to fully assess the default assumption that the interaction of EDCs with steroid hormone receptors in one species can be extrapolated to another.

## REGULATORY SCREENING AND TESTING

### *Mammalian Screening and Testing*

One response of the USEPA to the Safe Drinking Water and Food Quality and Protection Acts of 1996 was to form an advisory group, the Endocrine Disruptor Screening and Testing Advisory Committee (EDSTAC), to help formulate a tiered screening and testing strategy for EDCs (<http://www.epa.gov/scipoly/oscp/edspoverview/finalrpt.htm>). The screening battery recommended by EDSTAC was designed to detect alterations in developmental and reproductive processes controlled by the hypothalamic-pituitary-gonadal (HPG) and hypothalamic-pituitary-thyroid (HPT) axes. Based on a “weight-of-evidence” analysis, chemicals positive in Tier 1 screening (T1S) would be considered as potential EDCs and subjected to testing (Tier 2). Because T1S would be less expensive and time consuming than Tier 2 testing, equivocal effects in T1S could be replicated or evaluated further in additional short-term assays before more extensive Tier 2 testing was initiated. The goal of T1S is to use assays sensitive enough to detect EDCs, whereas issues of “dose–response, relevance of the route of exposure, sensitive life stages and adversity” would be resolved in the Tier 2 testing phase.

*In vitro* T1S assays that are under development would evaluate chemicals for AR and ER binding with rat AR and ER cytosolic, rat AR recombinant, and human ER $\alpha$  recombinant receptors, ER or AR dependent gene expression (Wilson *et al.*, 2002a, 2004a) or cell proliferation (Soto *et al.*, 2004) assays, aromatase enzyme inhibition, and inhibition of steroidogenesis in the H295R cell line (Heneweer *et al.*, 2004). Due to limitations of *in vitro* assays, it is necessary to include *in vivo* assays in the screening battery. *In vitro* assays alone cannot account for absorption, distribution, metabolism, and excretion (ADME), and they yield false negative and false-positive results. In particular, many false negatives result from the inability of the *in vitro* assays to activate EDCs metabolically. In addition, *in vitro* assays at high concentrations lack specificity and produce many false-positive results. Because *in vivo* assays not only account for ADME, but also can integrate all of the endocrine and nonendocrine toxicities, the combination of both *in vivo* and *in vitro* assays is recommended for screening.

To avoid the limitations described above, the EDSTAC proposed three short-term *in vivo* mammalian assays for the

T1S battery: the uterotrophic, Hershberger, and pubertal female rat assays (Gray, 1998a). Estrogen agonists and antagonists would be detected using the 3-day uterotrophic assay using subcutaneous administration of the test compound. Based on the evaluation of four variations of the uterotrophic assay protocol in organization for economic cooperation and development (OECD) interlaboratory studies, all of the protocols have produced acceptable responses without regard to rat strain, diet, or housing conditions (Owens and Koeter, 2003; Owens *et al.*, 2006). The selected uterotrophic assays for estrogens and antiestrogens use either the intact juvenile or the castrated ovariectomized adult/juvenile female rat that is dosed either by oral or subcutaneous routes.

The second *in vivo* assay in T1S, the Hershberger assay, detects antiandrogenic activity simply by weighing androgen-dependent tissues in the castrated male rat (Gray, 1998a, b; Hershberger *et al.*, 1953; Owens *et al.*, 2006, 2007). In this assay, weights of the ventral prostate, Cowper’s glands, seminal vesicles (with coagulating glands and fluids), glans penis, and levator ani/bulbocavernosus muscles are measured in castrated, testosterone-treated (or untreated) male rats after 10 days of oral treatment with the test compound. This assay is very sensitive for detection of androgens and antiandrogens. Other useful end points that help reveal the mode of action and specificity of the response include weights of the adrenal, liver, and kidney, and measurements of serum (collected by cardiac puncture) levels of testosterone and luteinizing hormone. The Hershberger assay shows high sensitivity and specificity to chemicals with AR-mediated activity. Weak antiandrogenic pesticides such as *p,p'*-DDE and linuron are easily detected in the Hershberger assay (Lambright *et al.*, 2000; Parks *et al.*, 2000; Yamasaki *et al.*, 2003). Chemicals such as finasteride, which inhibit 5 $\alpha$  reductase activity, also are active in this assay. 5 $\alpha$  Reductase inhibitors dramatically reduce male accessory sex gland weights with less effect on the levator ani/bulbocavernosus muscle, which has low levels of this enzyme (Blohm *et al.*, 1986). Chemicals that are positive in the Hershberger assay often produce adverse effects after *in utero* exposure and during puberty.

Both the rat uterotrophic and Hershberger assays have been used for several decades to screen chemicals for estrogenicity and androgenicity (Dorfman and Dorfman, 1962). Because estrogens are required for uterine growth and androgens for sex accessory gland growth in both rats and humans, chemicals that are positive in the rat assays for these endocrine activities can be expected to produce predictable responses in humans. These endocrine pathways are highly conserved; in fact, many drugs with antiestrogenic activity or antiandrogenic activity have been shown to produce the anticipated responses in both species. The OECD-led effort to standardize and validate the Uterotrophic and Hershberger assay has been completed with several publications on the different phases available, and OECD test guidelines for both these assays have been peer-reviewed, public comments addressed and are listed as draft

Test Guidelines on the OECD web site: [http://www.oecd.org/document/62/0,3343,en\\_2649\\_34377\\_2348606\\_1\\_1\\_1\\_1,00.html#Hershberger](http://www.oecd.org/document/62/0,3343,en_2649_34377_2348606_1_1_1_1,00.html#Hershberger).

The third *in vivo* mammalian/rat assay included in the proposed EDSTAC T1S battery, the pubertal female rat assay, has been used in our laboratory for nearly two decades (Gray, 1998a; Gray *et al.*, 1988a, b, 1989). This assay detects alterations in thyroid hormone status, HPG function, inhibition of steroidogenesis, and direct effects of estrogens, and antiestrogens. In this assay, weanling female rats are dosed daily by gavage for 21 days while the age at vaginal opening (puberty) is monitored. The females are necropsied at about 42 days of age and measurements include serum thyroid hormones, uterine and ovarian weight, and histology (reviewed by Goldman *et al.*, 2000) ([http://www.epa.gov/oscpmont/oscpendo/pubs/assayvalidation/pubertal\\_female\\_pr.htm](http://www.epa.gov/oscpmont/oscpendo/pubs/assayvalidation/pubertal_female_pr.htm)).

Alternative *in vivo* assays were also discussed by EDSTAC and are being evaluated by the EPA. If they are of sufficient sensitivity, specificity, and relevance, they might replace or augment current T1S assays.

One promising alternative assay also used extensively in our laboratory is the pubertal male rat assay (Gray *et al.*, 1999; Kelce *et al.*, 1995; Monosson *et al.*, 1999; Stoker *et al.*, 2000), which detects alterations in thyroid function, HPG maturation, steroidogenesis, and altered steroid hormone function. Intact weanling males are exposed to the test substance for approximately 30 days during which the age at puberty is determined by measuring the age at preputial separation, and reproductive tissues are evaluated and serum taken for optional hormonal analyses at necropsy. We suggest measurement of serum testosterone be included because it would assist in determining chemical mode of action (e.g., separating AR antagonist vs. inhibitors of hormone synthesis) and enhance the sensitivity of the assay to chemicals that inhibit steroid hormone synthesis: [http://www.epa.gov/oscpmont/oscpendo/pubs/assayvalidation/pubertal\\_male\\_pr.htm](http://www.epa.gov/oscpmont/oscpendo/pubs/assayvalidation/pubertal_male_pr.htm).

The EDSTAC recommended that the EPA develop and evaluate an *in utero* lactational screening assay due to the unique sensitivity of the fetal reproductive system to disruption by some toxicants. For example, TCDD alters sexual differentiation of male and female rats and hamsters at dosage levels approximately two orders of magnitude below those required to produce adverse effects in pubertal or adult rats (Gray *et al.*, 1997b, c; Wolf *et al.*, 1999). One version of the proposed *in utero* lactational assay now being evaluated by the EPA takes about 80 days and uses approximately 10 litters per group (120–150 pups). In this protocol, androgens and antiandrogens can be detected in approximately 2 to 3 wk, and EDCs with antithyroid activity can be detected in infant or weanling offspring after 4–5 weeks of maternal treatment. However, an EPA Science Advisory Panel recently concluded the above protocol is more suitable for testing than use as a screening assay due to the size and duration of the study but that efforts to streamline such an assay for use in screening

were worthy of consideration, <http://www.epa.gov/scipoly/sap/meetings/index.htm#january>. A scaled-down protocol that might be useful for screening with *in utero* and lactational exposure could include five to six dose groups over a broad dose range, three to four litters per dose group and carefully examine all the pups of both sexes (i.e., Rider *et al.*, 2008). If effects were noted, then this would be followed by a definitive multigenerational study to define NOAELs.

#### *Tier 2 Testing as Defined by EDSTAC and the USEPA*

The purpose of Tier 2 testing is to provide “definitive” information for hazard characterization of endocrine-disrupting agents. This information adds to the body of knowledge essential for risk assessment for an EDC by


1. Either confirming or refuting the observations noted in the T1S screens/assays. The Tier 2 *in vivo* assays data can expand on the information derived from the *in vitro* and *in vivo* screens and use a route of exposure that will be reflective of likely exposure of humans to the agent under investigation.

2. For the first time in the tiered process, studies involving mammals utilize an *in utero* exposure paradigm that includes exposures in critical developmental windows and assessments not evaluated in standard prenatal developmental toxicity studies. For many EDCs, this is the most sensitive life stage for the induction of adverse effects leading to potentially permanent changes in phenotype and function that would not be noted in adult animal toxicity studies.

3. Identifying in the experimental animal species (most likely the rat) the activity of suspected EDCs for end points for which concerns have been raised in humans (e.g., hypospadias, cryptorchidism) as well as identify other endocrine-like activity associated with androgens, estrogens, and antithyroid acting agents.

4. Not only detecting activity, but also providing the appropriate dose–response information for use in any risk assessment on individual chemicals.

When the EPA first proposed a Tier 2 test for mammals it noted that the multigeneration study (Fig. 1) was the only one likely to have an appropriate exposure period covering the major developmental life stages of interest. EPA also had built into the protocol, appropriate times for evaluations of effects, especially those that might show latency (e.g., an exposure *in utero* only producing a readily identifiable change at adulthood). Even though the improvements in protocol study design in the most recent EPA test guidelines ([http://www.epa.gov/opptsfrs/publications/OPPTS\\_Harmonized/870\\_Health\\_Effects\\_Test\\_Guidelines/Series/870-3800.pdf](http://www.epa.gov/opptsfrs/publications/OPPTS_Harmonized/870_Health_Effects_Test_Guidelines/Series/870-3800.pdf)) did incorporate more endocrine-mediated end points (e.g., sperm analyses, indices of puberty in males and females), EDSTAC realized that some improvements would have to be made to the end points incorporated in the multigeneration protocol to make the assay more comprehensive in the detection of effects (e.g., improved pathology and endocrine assessment, sexually



**FIG. 1.** Schematic of a standard multigenerational test.

dimorphic phenotypic assessments). It was also noted that many registrants and producers could bypass T1S screening and go directly to Tier 2 (e.g., for pesticides), but that the current protocols might be insensitive to both detection and providing accurate dose–response information.

There is also a large gap (in terms of comprehensiveness and cost) between the T1S screens and Tier 2 tests. Many investigators had successfully used a transgenerational approach (usually employing approximately 10 litters per dose group but analyzing every pup in the litter on reaching adulthood) which employed fewer animals, and was significantly cheaper than a standard multigeneration study in identifying adverse endocrine-mediated end points or found them at lower dose levels than seen in conventional multigeneration studies (e.g., linuron, Gray *et al.*, 1999; Lambright

*et al.*, 2000; McIntyre *et al.*, 2000; di-*iso*-nonyl phthalate, Gray *et al.*, 1999; Waterman *et al.*, 2000).

The use of a lower litter number but increasing the number of pups per litter examined provides greater statistical power over the conventional multigeneration study with 20 litters per dose group, but evaluation of only one male and one female offspring per litter at adulthood. The degree to which the statistical power increases with examination of multiple pups per litter is dependent upon the degree to which pups within a litter differ from one another (the intralitter correlation coefficient [ICC]). We have calculated ICCs and determined how examining multiple male pups in a litter affected the power calculations for a number of reproductive end points from several of our studies. In these studies, dams were exposed to EDCs *in utero* and the offspring examined later in

TABLE 3  
Calculation of Cox's Ratios and ICC From proc Mixed Analyses

Adult necropsy weight data	Litter variance	Pups-residual variance	Cox's ratio $4 \times$ (pups/litters)	ICC litter/(litter + pups)
Epididymis	5021	5167	4.116	0.493
Liver	4.95	6.22	5.031	0.443
Body	2579	3481	5.399	0.426
Kidney	23,537	32,670	5.552	0.419
Cowper's glands	1129	1694	6.002	0.400
Testis	21,506	32,376	6.022	0.399
LABC	10,417	15,736	6.042	0.398
Glans penis	31.2	56.5	7.244	0.356
Seminal vesicle	25,980	71,543	11.015	0.266
Ventral prostate	5346	23,258	17.402	0.187
Variable pubertal necropsy <sup>a</sup>	Litter variance	Pups-residual variance	Cox's ratio $4 \times$ (pups/litters)	ICC litter/(litter + pups)
Body weight, day 18	14.225	7.131	2.005	0.666
Body weight, day 44	348.2	201.02	2.309	0.634
Body weight final	753.87	492.14	2.611	0.605
AGD, day 2	0.05503	0.06889	5.007	0.444
Glans penis weight	44.4	70.85	6.383	0.385
Epididymis weight	839.64	1708.64	8.140	0.329
Testis weight	23,297	61,044	10.481	0.276
LABC	2666	7417.5	11.129	0.264
Seminal vesicle weight	4925	14,445	11.732	0.254
Adrenal weight	8.214	34	16.719	0.193
Age at preputial separation	1.391	8.0744	23.219	0.147
Weight at preputial separation	107.00	720.00	26.916	0.129
Cowper's glands	14.9954	189.57	50.568	0.073

Note. LABC, levator ani bulbocavernosus weight. Power curves flatten out when the number of pups sampled per litter exceeds Cox's ratio (Bergerud, 1995). End points are ordered based upon ICC values, ranging from high (pups more correlated within the litter) to low (pups less correlated within the litter).

<sup>a</sup>The "pubertal" necropsy included all the males dosed by gavage from 18 to 65 days of age with the same dose level administered to the dam during gestation and lactation. For males necropsied as adults, only the dam was treated and exposure was discontinued on day 18 of lactation. The lower the ICC, the less related are the values of the pups within litters.


life. In the data shown here as an example, DEHP was administered to the dam orally from day 8 of gestational to day 18 of lactation, half the male rat offspring being further exposed from 18 to 65 days of age and all the males were examined thoroughly (Gray *et al.*, unpublished data). The results of the DEHP study indicate (Table 3; Fig. 2) that sampling two to three pups from 15 litters, or all of the pups from ten litters/dose provides the same or greater statistical power as sampling one pup from 20 litters. More information on these calculations is provided in the text below.

Using more pups per litter also addresses the issue of statistical power to detect malformations that occur at low incidences in the lower dosage groups. It is not just the continuous variables but also the malformations that can have improved detection and provide better assessments of dose-response relationships (Fig. 2).

These studies also did not employ triggered end points, but made measurements in all offspring and used other sensitive

measures of endocrine activity (e.g., anogenital distance in F<sub>1</sub> offspring, assessment of areola and nipple retention in preweaning F<sub>1</sub> pups and at adulthood) not currently incorporated into standard multigeneration studies. Such trans-generational studies may offer an intermediate tier between T1S and Tier 2. Although they do not evaluate the reproductive performance of the F1 offspring, such studies could serve as a less expensive intermediate tier to re-enforce the occurrence of endocrine activity that was flagged in T1S, before embarking on any modified multigeneration study, or perhaps could act as a bridging study from an older multigeneration study.

Whatever improvement may be required in the current EPA multigeneration protocols that would serve as the definitive Tier 2 test it would be prudent to think that all the information obtained in T1S should be used in tailoring the study design to move away from a "one size fits all" approach. If no T1S data are obtained, Tier 2 must represent a comprehensive evaluation



**FIG. 2.** Power curves for reproductive organ weights, malformation rates, anogenital distance and age at preputial separation (puberty). The data from our transgenerational study with DEHP were analyzed using PROC MIXED available on SAS to obtain estimates of components of variation (for litters vs. pups within litters) that make up the overall variability of the means in order to assess the implications for statistical power of using data from multiple pups per litter. Following this analysis, we calculated power to detect the treatment effects as described by Raudenbush and Xiao-Feng (2001). The objective of this was to determine the proportion of the overall error variance due to litter-to-litter variability versus the proportion accounted for by the pups within litters. This retrospective analysis can be useful for designing optimal sampling strategies for future studies. It basically allows one to determine how much the statistical power of a study is enhanced by examining several pups from the same litter rather than using only one pup/per sex/litter. Because the pups in the DEHP treated litters do not respond identically, the more variable the pups are within the litter the more power is enhanced, and hence the standard error of the mean is reduced, by examining multiple pups from the same litter.



of activity with sufficient statistical and experimental power to be able to detect events likely to have an endocrine-mediated mode of action (e.g., reproductive tract malformations). The current studies with limited end points and numbers of offspring examined per litter, have much poorer resolving power than, for example, the current prenatal developmental toxicity study to detect birth defects, such that incidences of reproductive tract malformations in excess of 25% with a very low control background incidence cannot be detected.

Newer efforts to move to a more tailored approach to studying reproductive toxicity have been proposed by the Agricultural and Chemical Safety Assessment committee of ILSI (Cooper *et al.*, 2006). This life-stage approach, although never having been tested, contains many end points and approaches that have been used successfully in the past and acknowledges that a flexible approach needs to be taken based on all the available information to adequately test for reproductive toxicity, including those aspects of postnatal development that have been of serious concern in the evaluation of EDCs. However, only three chemicals are being selected with known reproductive, neuro- or immunotoxicity for evaluation in the protocol. This approach is based on many of the transgenerational types of studies that have been previously published. The authors also acknowledge the weakness in having too few pups to examine in every litter and advocate testing more, in different cohorts that would evaluate neuro- and immunotoxicity.

Whilst this represents a new initiative to incorporate the best science into testing, the protocol presented in the publication unfortunately fell short of the ideal, definitive test for EDCs. In particular, the different cohorts of animals are examined at different times so that at sexual maturity (usually deemed to be ~90 days of age in the rat) there is only 1 cohort available with 1 male and 1 female representative of the original litter, which is no better than we have with the current multigeneration protocol. In common with the transgenerational studies, there is no routine evaluation of the reproductive performance of the  $F_1$  offspring. These shortcomings have been discussed and an improved protocol was presented to a USEPA Science Advisory Panel in, 2007 and this effort is being pursued by the OECD.

Cooper *et al.* (2006) advocated the use of triggers to incorporate additional end points into the design. Unfortunately, our experience with triggered end points in the current EPA multigeneration study design has not been good. In many instances, this is not a fault of the experimenters, but more because of lack of time for evaluation of data. So, for example, male pubertal indices are captured on the study around postnatal day 45. With the normal staggering of individuals on a reproduction study (not all are born on the same day) it would be several days, or even longer if an effect is noted, before all the males had complete evaluations of puberty. These data would then need to be statistically analyzed corrected for body weight and any other confounders and then a decision

made on the conduct of a triggered endpoint (which may involve a Contract Research Organization contacting sponsors for approval). This may have to be done within 40 days, which places significant pressure on the scientists conducting these studies and may not be reliably achievable. A much better approach would be the application of "opt out" triggers. That is, all the end points for a study would be collected unless there was sufficient information and justification not to undertake a specific measurement. As the protocol currently exists in its published form (Cooper *et al.*, 2006), this would not suffice as a replacement for the current multigeneration study, but could be adapted to make it meet this need and we are optimistic that this approach will serve as a useful template which can in time be fashioned into a very useful protocol.

#### *Power Calculations, Analysis of Components of Variation, and Intralitter correlation coefficients*

The data for our DEHP transgenerational study were analyzed using PROC MIXED available on SAS to obtain estimates of components of variation (for litters vs. pups within litters) that make up the overall variability of the means in order to calculate power to detect the treatment effects (Raudenbush and Xiao-Feng, 2001) (Fig. 2; Table 3). Because the pups in the DEHP treated litters do not respond identically, the more variable the pups are within the litter the more power is enhanced by examining multiple pups from the same litter. We also employed Cox's Ratio as a general rule to determine how many pups per litter could be sampled to enhance the power of the study (Bergerud, 1995) (<http://www.for.gov.bc.ca/hre/bio-pamph/pamp50.pdf>). ICC also were calculated to describe the degree of similarity of pups within litters for the multiple end points measured in the current study. According to Cox's Rule Of Thumb (Bergerud, 1995) there is little or no increase in power when the number of subsamples (pups in this case) is greater than Cox's Ratio [Cox's ratio =  $4 \times (\text{variation due to pups within litters} / \text{variation due to litters})$ ]. The power curves flatten out when the number of pups/litter sampled is increased, indicating that the most efficient design would not sample any more pups per litter beyond this Ratio. When the components of variance were estimated for reproductive measures it is clear that, even in the most conservative case, sampling all the male pups in each litter would enhance the power of a transgenerational reproduction study. As examples, power curves were calculated for the control and high dose group (300 mg/kg/day) for one to five pups per litter and 10, 15, and 20 (standard sample size) litters per dose.

Power calculations also were calculated for categorical effects based upon the numbers of malformed males versus males without malformations per dose group using SigmaStat 3.1 software (Fig. 2). If 20 animals per dose group are examined for malformations then lesions occurring at an incidence of 25% or greater can be detected whereas an incidence of 10% can be detected if all the pups are examined

from 20 litters. If only ten males per group are examined, as recommended for histopathological analyses in some regulatory agency test guidelines, then effects are only detected statistically if about 50% or more of the tissues/organs are affected; a level of statistical power that many would consider inadequate.

### Nonmammalian Screening and Testing

As discussed previously, there is compelling evidence for adverse effects of EDCs on invertebrates, fish, and wildlife. This has provided one impetus for the development of EDC screening and testing assays with nonmammalian species; however, there also are examples where nonmammalian models may prove superior to rodents in detecting some EDC MOA.

The class of animals that has received the most attention in terms of EDC testing is fish. Many different methodological approaches have been applied in fish EDC tests; an experimental design that has been used quite frequently is a 21-day assay initiated with reproductively mature adults. Different freshwater small fish models, including the Japanese medaka (*Oryzias latipes*) and zebrafish (*Danio rerio*), have been used in 21-day EDC screening assays (Seki *et al.*, 2006); however in the United States, the small fish species of choice for the 21-day design has been the fathead minnow (*Pimephales promelas*; status of this and other assays reviewed at (<http://www.epa.gov/endo/pubs/assayvalidation/status.htm>). The 21-day Tier 1 fathead minnow test features a range of apical and mechanistic end points designed not only to detect the adverse reproductive effects of exposure to the test chemicals, but target specific classes of EDCs (Ankley *et al.*, 2001). For example, an endpoint in the assay that is exquisitely sensitive to (and specific for) estrogenic EDCs is induction of vitellogenin (egg yolk protein) in males, which, under normal physiological conditions, do not express the protein (Ankley *et al.*, 2001; Seki *et al.*, 2006). Decreases in plasma vitellogenin concentrations in females have been shown to be indicative of inhibitors of steroidogenic enzymes, including CYP19 aromatase (Ankley *et al.*, 2005). The 21-day assay also is effective in identifying androgenic EDCs, in that female fish exposed to AR agonists exhibit readily apparent morphological masculinization (Ankley *et al.*, 2001; Seki *et al.*, 2006).

Fish also are being used in longer-term (e.g., Tier 2) tests to detect full life-cycle and multigenerational effects of EDCs. Species that have received the most attention in the United States in this regard have been an estuarine fish, the sheepshead minnow (*Cyprinodon variegates*), and the Japanese medaka, which is attractive for this work due to its relatively rapid life cycle (Ankley and Johnson, 2004). The medaka also is widely used in Japan for long-term EDC testing, as is the zebrafish in some European countries (Peterson *et al.*, 2001). Responses assessed in the longer-term EDC assays with fish are similar to the apical and diagnostic end points used in the 21-day

reproduction tests, but typically also include measurements reflective of chemical effects on the endocrine system during early life stages, such as gonadal development and sexual differentiation in F1 and, perhaps, F2 generations (Ankley and Johnson, 2004).

Amphibian species have received significant attention in the development and implementation of EDC testing programs as recommended by EDSTAC. Amphibian metamorphosis is almost certainly the most visually obvious manifestation of effects on HPT function in a vertebrate; from this, it was initially recommended that a simple tail resorption assay with the model lab species *Xenopus laevis* could serve as the basis of a screen for chemicals that act as agonists or antagonists of thyroid hormone function. Subsequent research suggested that the tail resorption endpoint was for various reasons problematic, but metamorphic frogs nonetheless were a good model for detecting thyroid-active EDCs (Degitz *et al.*, 2005; Opitz *et al.*, 2006). Currently the endpoint that seems best suited for this purpose is thyroid gland histology, collected in conjunction with some indication of developmental stage; this basic approach (with *X. laevis*) is being used for EDC screening programs being implemented both by the USEPA and Organization for Economic Cooperation and Development (Degitz *et al.*, 2005; Opitz *et al.*, 2006).

There has been interest in North America in the development of a full life-cycle test with amphibian species that could serve as a tool (e.g., in Tier 2) for assessing the long-term ecological risk of EDCs. Research in this area has focused more on *Xenopus tropicalis* than *X. laevis*, due in large part to the more rapid life cycle of the former species (Degitz, unpublished data). In addition to considering responses controlled by thyroid function, such a test would feature end points (e.g., vitellogenin production, gonad histopathology) related to disruption of the HPG axis by (anti-)estrogenic/androgenic chemicals (Degitz, unpublished data).

In addition to fish and amphibians, there have been efforts with other nonmammalian vertebrates to develop standardized assays suitable for assessing the effects of EDCs. A notable effort in this regard involves birds. Full life-cycle tests with birds have been used for some time for pesticide registration; with some adaptation, these basic tests would be applicable to Tier 2 testing of EDCs (Touart, 2004). For example, to detect possible transgenerational effects, it has been proposed that the test design encompass two generations of animals, with (dietary) chemical exposures starting with sexually mature adults, proceeding through a complete F<sub>1</sub> generation, and culminating with production of F<sub>2</sub> young. Either of two species would be amenable to this type of experimental design; however, due to their relatively more rapid life cycle, the Japanese quail (*Coturnix japonica*) likely would be a more resource-effective model than the northern bobwhite quail (*Colinus virginianus*) for routine testing. In addition to apical responses like fecundity and fertility, end points specifically indicative of endocrine-disruptive effects, like gonad development and differentiation

and certain changes in behavior, are being incorporated into the two-generation avian test design (Touart, 2004).

There also have been efforts to develop standardized tests for EDCs with different invertebrate species (deFur, 2004). A challenge in this area has been definition of what invertebrate tests are intended to detect because the role, if any, of many of the mammalian steroidal and protein hormones is uncertain in invertebrates. If the assays are to be used to identify chemicals that affect the vertebrate HPG axis, then there is some question as to whether functional aspects of the axis are sufficiently conserved across animal phyla to expect that sensitive, biologically meaningful tests could be developed in invertebrate models. On the other hand, invertebrates possess a variety of endocrine systems/function that could be adversely affected by chemicals, which often would not be “flagged” as EDCs using vertebrate tests (deFur, 2004). Under this scenario there is an easily justifiable need for standardized tests for EDCs using invertebrate models. One test recommended for inclusion in the formal EPA testing program for EDCs is a life-cycle test with *Mysid* shrimp. Other invertebrates that have received significant attention relative to full life-cycle tests for EDCs include daphnids and copepods (deFur, 2004).

## CONCLUSIONS

The field of endocrine disrupters has come a long way since its initial impetus in 1991. Many of the concepts advanced in this area have now been embraced by the general toxicological community resulting in useful enhancements of testing assays, new mechanism-based screening assays and incorporation of toxicant-induced endocrine alterations into risk assessment. Concepts such as the special susceptibility of the developing organism and early induction of latent effects, for example, are now widely held and the mechanism-based approach to toxicology testing and screening has been incorporated into many other branches of toxicology. Parallel to the EDCs there also is an emerging field of “developmental basis of adult health,” or “fetal programming of adult disease” (de Boo and Harding, 2006; Young, 2001). This hypothesis is a variation of the many of the EDC concepts, but applies to potential adverse effects on endpoints such as blood pressure, pancreatic function, insulin sensitivity, and even brain and lung function. The concept is the same: exposure during development may induce profound changes in adult function, and extends that concept past the endocrine system.

Since the Wingspread Conference of 1991, there have been numerous examples of adverse effects of EDCs in fish, wildlife, domestic animals, and humans. Endocrine systems have been disrupted by anthropogenic chemicals including antiandrogens, androgens, estrogens, AhR agonists, inhibitors of steroid hormone synthesis, antithyroidal compounds, and retinoid agonists. In addition, potential pathways and targets for endocrine disruption extend beyond the traditional EAT receptor-mediated reproductive and developmental systems.

TABLE 4  
Where Do We Go from Here?

1. Ecosystem health
  - 1.a. Better characterization of the extent and population impacts on fish of estrogenic municipal effluents, and further research concerning the possible impacts on fish and wildlife of other known EDCs entering the environment (e.g., from pulp and paper mills, CAFOs).
  - 1.b. Clearer delineation of exact chemicals responsible for effluent estrogenicity (i.e., ethinyl estradiol vs. E2 vs. nonylphenol) as a basis for identifying mitigation options.
  - 1.c. Identification of pharmaceuticals that could possibly cause ecological effects through properties similar to EDCs, that is, high-potency chemicals focused on biological pathways (other than the HPG axis) that have a substantial degree of phylogenetic conservation.
2. EDC screening
  - 2.a. Evaluation of the EPA T1S assay suite to determine “holes” and redundancies with respect to identifying HPG-active chemicals (i.e., we aren’t going to be able to apply the full suite of tests to very many chemicals-too expensive); an understanding of endpoint/species extrapolation will be critical here.
  - 2.b. Improve predictiveness of screens and tests for EDC activities.
  - 2.c. Enhance and expand the *in vitro* assays for T1S to replace animal-derived proteins with recombinant receptors, adapt *in vitro* assays for high throughput.
  - 2.d. Determine the need for *in vitro* assays using nonmammalian pathways and develop the assays for screening, as warranted.
  - 2.e. Develop an *in utero* lactational screening assay for possible inclusion in T1S.
  - 2.f. Implement a T1S battery that meets or exceeds the EDSTAC 1998 recommendations.
3. Multigenerational and transgenerational testing including a postnatal examination of offspring
  - 3.a. Enhanced multigenerational assays that include sensitive and reliable endpoints, that use animals more efficiently (fewer litters/dose, examine more pups/litter, increase histopathological examinations to include all F<sub>1</sub> animals) without losing statistical power.
4. Clearer communication and cross-disciplinary training on the following:
  - 4.a. Appropriate experimental designs and statistical analyses of postnatal study data for scientists from different disciplines as well as understanding of basic toxicological issues (endocrinology, toxicogenomics, proteomics, epigenetics, etc.) so that research resources are not wasted. It is evident from the literature that special attention needs to be given to assure that studies that examine more than one pup per litter account for this in the analysis of the data. Furthermore, investigators that cross-foster pups should do so properly, retaining both prenatal and postnatal litter information.
  - 4.b. Use of reproductive and developmental toxicity data in risk assessment. Promote an understanding of the difference between and no observed effect level and a NOAEL and the need to causally link effects not generally recognized as adverse to clear adverse effects.
5. Maintain an integrated biological approach to study endocrine action (beware of systems biology and the drilling down to molecular levels *in vitro* as cells and tissues live in complex environments). Integrate EDC latent effects research with other projects on the fetal basis of adult disease.
6. Risk assessment on EDCs
  - 6.a. Develop a scientifically defensible and rational framework for cumulative risk assessments, incorporating the knowledge that dose additivity applies to chemicals that target the same endocrine pathway via very different cellular mechanisms. This redefines what we mean by mode of action and encompasses multiple mechanisms inducing the same phenotypic changes.
  - 6.b. Expanded use of endocrine data in risk assessments not only to support mode of action but also as critical effects.

Although much has been learned over the past 15 plus years, future research needs to address critical issues that have arisen since the early 1990's. For example, reports of nonmonotonic (e.g., U-shaped), and low dose effects and nonthreshold effects for EDCs continue to be a challenge for risk assessment. Future well-designed studies are required to address the biological plausibility of low dose effects as well as the ramifications for risk assessment. Furthermore, new mechanisms of endocrine action continue to be discovered. Membrane-bound steroid receptors, novel inhibitors of steroidogenesis, and altered metabolism are just a few of the new areas of investigation.

In addition to the remaining questions of the field, there are multiple emerging areas to be investigated. One of these areas involves mixtures toxicology. It is clear that, rather than focusing on single chemical exposures, data are required to determine and predict potential cumulative effects of EDCs. Pharmaceuticals and veterinary drugs in the environment also present a new challenge. Pharmaceuticals, be they from personal, medical, or agricultural uses have been detected in the environment and these chemicals may be disrupting both target and nontarget systems in exposed populations of wildlife and fish (Ankley *et al.*, 2007). The potential role of EDCs in nontraditional biological pathways also warrants further investigation. Studies implicating EDCs in diabetes, obesity, and behavioral and immune deficiencies have underscored the need to elucidate these areas of endocrine disruptor research. These new areas of research may eventually alter the screening and testing batteries currently proposed by regulatory agencies such as the USEPA and OECD. Until that time, there are fundamental questions remaining about the proposed batteries of screens and tests as described above. Finally, from a regulatory standpoint, current interest in using toxicogenomic data for risk assessment requires further evaluation. Inherent problems with variability, reproducibility, and biological significance continue to challenge application of these promising technologies. Risk assessors have traditionally required that changes in gene expression be validated by some other measure such as endocrine or biochemical "biomarkers" such that a causal link can be established between mechanistic data and an effect that is clearly adverse. Furthermore, one then needs to identify a "point of departure" or the degree of change in the mechanistic endpoint that consistently produces an adverse effect.

In summary, research on EDCs has progressed significantly and now provides concrete examples of chemicals in the environment adversely affecting endocrine systems in diverse fauna, ranging from invertebrates to humans. Many of the concepts introduced in the consensus statement from the Wingspread Conference in 1991 by Dr Theo Colborn (Colborn and Clement, 1992) and the other workshop participants are now generally accepted by the scientific community which in turn has improved our understanding of how "upstream," cellular and molecular alterations in toxicity pathways produce adverse effects "downstream" at higher dose levels and how

exposure to EDCs during critical developmental periods can result in adverse effects later in life. Although much has been learned, the field is relatively new, and many relevant and timely questions remain. Some of the important areas of emphasis in the field are presented in Table 4 "*Where do we go from here?*". The continued introduction of novel chemicals (drugs, toxic substances, pesticides, nanomaterials, etc.) into the environment requires ongoing research and monitoring to ensure the safeguarding of the environment for generations to come.

## FUNDING

USEPA/NCSU Cooperative Training agreement (CT826512010) to A.K.H, C.V.R, C.R.B.; Integrated Graduate Education, Research and Training grant from the National Science Foundation to C.L.G.

## ACKNOWLEDGMENTS

We would like to thank Dr. Chris Wiesen, Odum Institute for Research in Social Science, University of North Carolina at Chapel Hill, for statistical assistance with the calculation of power from the proc mixed output from SAS.

## REFERENCES

- Adams, N. R. (1995). Organizational and activational effects of phytoestrogens on the reproductive tract of the ewe. *Proc. Soc. Exp. Biol. Med. Soc. Exp. Biol. Med.* **208**, 87–91.
- Ames, B. N., and Gold, L. S. (2000). Paracelsus to parascience: the environmental cancer distraction. *Mutation research* **447**, 3–13.
- Ankley, G. T., Brooks, B. W., Huggett, D. B., and Sumpter, J. P. (2007). Repeating History: pharmaceuticals in the environment. *Environ. Sci. Technol.* **41**, 8211–8217.
- Ankley, G., Mihaich, E., Stahl, R., Tillitt, D., Colborn, T., McMaster, S., Miller, R., Bantle, J., Campbell, P., Denslow, N., *et al.* (1998). Overview of a workshop on screening methods for detecting potential [anti-] estrogenic/androgenic chemicals in wildlife. *Environ. Toxicol. Chem.* **17**, 68–87.
- Ankley, G. T., and Giesy, J. P., Eds. (1998). *Endocrine Disruptors in Wildlife: A Weight of Evidence Perspective*. SETAC Press, Pensacola.
- Ankley, G. T., Jensen, K. M., Durhan, E. J., Makynen, E. A., Butterworth, B. C., Kahl, M. D., Villeneuve, D. L., Linnum, A., Gray, L. E., Cardon, M., *et al.* (2005). Effects of two fungicides with multiple modes of action on reproductive endocrine function in the fathead minnow (*Pimephales promelas*). *Toxicol. Sci.* **86**, 300–308.
- Ankley, G. T., Jensen, K. M., Kahl, M. D., Korte, J. J., and Makynen, E. A. (2001). Description and evaluation of a short-term reproduction test with the fathead minnow (*Pimephales promelas*). *Environ. Toxicol. Chem. SETAC* **20**, 1276–1290.
- Ankley, G. T., Jensen, K. M., Makynen, E. A., Kahl, M. D., Korte, J. J., Hornung, M. W., Henry, T. R., Denny, J. S., Leino, R. L., Wilson, V. S., *et al.* (2003). Effects of the androgenic growth promoter 17-beta-trenbolone on fecundity and reproductive endocrinology of the fathead minnow. *Environ. Toxicol. Chem. SETAC* **22**, 1350–1360.



- Ankley, G. T., and Johnson, R. D. (2004). Small fish models for identifying and assessing the effects of endocrine-disrupting chemicals. *ILAR J.* **45**, 469–483.
- Anway, M. D., Cupp, A. S., Uzumcu, M., and Skinner, M. K. (2005). Epigenetic transgenerational actions of endocrine disruptors and male fertility. *Science* **308**, 1466–1469.
- Bell, D. R., Clode, S., Fan, M. Q., Fernandes, A., Foster, P. M., Jiang, T., Loizou, G., MacNicol, A., Miller, B. G., Rose, M., *et al.* (2007). Toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in the developing male Wistar(Han) rat. II: Chronic dosing causes developmental delay. *Toxicol. Sci.* **99**, 224–233.
- Bergerud, W. (1995). Power analysis and sample sizes for randomized block designs with subsampling. *Biom. Inf. B. C. Min. For. Range Res. Pamphlet* **50**, 1–4.
- Birkhoj, M., Nellemann, C., Jarfelt, K., Jacobsen, H., Andersen, H. R., Dalgaard, M., and Vinggaard, A. M. (2004). The combined antiandrogenic effects of five commonly used pesticides. *Toxicol. Appl. Pharmacol.* **201**, 10–20.
- Bjerke, D. L., Brown, T. J., MacLusky, N. J., Hochberg, R. B., and Peterson, R. E. (1994a). Partial demasculinization and feminization of sex behavior in male rats by in utero and lactational exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin is not associated with alterations in estrogen receptor binding or volumes of sexually differentiated brain nuclei. *Toxicol. Appl. Pharmacol.* **127**, 258–267.
- Bjerke, D. L., and Peterson, R. E. (1994). Reproductive toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in male rats: Different effects of in utero versus lactational exposure. *Toxicol. Appl. Pharmacol.* **127**, 241–249.
- Bjerke, D. L., Sommer, R. J., Moore, R. W., and Peterson, R. E. (1994b). Effects of in utero and lactational 2,3,7,8-tetrachlorodibenzo-p-dioxin exposure on responsiveness of the male rat reproductive system to testosterone stimulation in adulthood. *Toxicol. Appl. Pharmacol.* **127**, 250–257.
- Blohm, T. R., Laughlin, M. E., Benson, H. D., Johnston, J. O., Wright, C. L., Schatzman, G. L., and Weintraub, P. M. (1986). Pharmacological induction of 5 alpha-reductase deficiency in the rat: Separation of testosterone-mediated and 5 alpha-dihydrotestosterone-mediated effects. *Endocrinology* **119**, 959–966.
- Blystone, C. R., Furr, J., Lambright, C. S., Howdeshell, K. L., Ryan, B. C., Wilson, V. S., Leblanc, G. A., and Gray, L. E., Jr. (2007a). Prochloraz inhibits testosterone production at dosage below those that affect androgen-dependent organ weights or the onset of puberty in the male Sprague Dawley rat. *Toxicol. Sci.* **97**, 65–74.
- Blystone, C. R., Lambright, C. S., Howdeshell, K. L., Furr, J., Sternberg, R. M., Butterworth, B. C., Durhan, E. J., Makynen, E. A., Ankley, G. T., Wilson, V. S., *et al.* (2007b). Sensitivity of fetal rat testicular steroidogenesis to maternal prochloraz exposure and the underlying mechanism of inhibition: Prochloraz reduced fetal testis testosterone. *Toxicol. Sci.* **97**, 512–519.
- Brian, J. V., Harris, C. A., Scholze, M., Backhaus, T., Booy, P., Lamoree, M., Pojana, G., Jonkers, N., Runnalls, T., Bonfa, A., *et al.* (2005). Accurate prediction of the response of freshwater fish to a mixture of estrogenic chemicals. *Environ. Health Perspect.* **113**, 721–728.
- Brouwer, A., Longnecker, M. P., Birnbaum, L. S., Coglian, J., Kostyniak, P., Moore, J., Schantz, S., and Winneke, G. (1999). Characterization of potential endocrine-related health effects at low-dose levels of exposure to PCBs. *Environ. Health Perspect.* **107**(Suppl. 4), 639–649.
- Brunskill, P. J. (1992). The effects of fetal exposure to danazol. *Br. J. Obstet. Gynaecol.* **99**, 212–215.
- Cammack, J. N., White, R. D., Gordon, D., Gass, J., Hecker, L., Conine, D., Bruen, U. S., Friedman, M., Echols, C., Yeh, T. Y., *et al.* (2003). Evaluation of reproductive development following intravenous and oral exposure to DEHP in male neonatal rats. *Int. J. Toxicol.* **22**, 159–174.
- CERHR, NTP. (2003). NTP-CERHR monograph on the potential human reproductive and developmental effects of 2-bromopropane (2-BP). *Ntp. Cerhr. Mon.* i–III11.
- Chambers, D. B., and Leiker, T. J. (2006). A Reconnaissance for Emerging Contaminants in the South Branch Potomac River, Cacapon River, and Williams River Basins, West Virginia. USGS.
- Clark, H. A., and Snedeker, S. M. (2005). Critical evaluation of the cancer risk of dibromochloropropane (DBCP). *J. Environ. Sci. Health* **23**, 215–260.
- Colborn, T., and Clement, C. (1992). Chemically-Induced Alterations in Sexual and Functional Development: The Wildlife/Human Connection. Princeton Scientific Publishing Co., Inc., Princeton, NJ.
- Cooper, R. L., Lamb, J. C., Barlow, S. M., Bentley, K., Brady, A. M., Doerr, N. G., Eisenbrandt, D. L., Fenner-Crisp, P. A., Hines, R. N., Irvine, L. F., *et al.* (2006). A tiered approach to life stages testing for agricultural chemical safety assessment. *Crit. Rev. Toxicol.* **36**, 69–98.
- Crofton, K. M., Craft, E. S., Hedge, J. M., Gennings, C., Simmons, J. E., Carchman, R. A., Carter, W. H., Jr., and DeVito, M. J. (2005). Thyroid-hormone-disrupting chemicals: Evidence for dose-dependent additivity or synergism. *Environ. Health Perspect.* **113**, 1549–1554.
- de Boo, H. A., and Harding, J. E. (2006). The developmental origins of adult disease (Barker) hypothesis. *Aust. N. Z. J. Obstet. Gynaecol.* **46**, 4–14.
- deFur, P. L. (2004). Use and role of invertebrate models in endocrine disruptor research and testing. *ILAR J.* **45**, 484–493.
- Degitz, S. J., Holcombe, G. W., Flynn, K. M., Kosian, P. A., Korte, J. J., and Tietge, J. E. (2005). Progress towards development of an amphibian-based thyroid screening assay using *Xenopus laevis*. Organismal and thyroidal responses to the model compounds 6-propylthiouracil, methimazole, and thyroxine. *Toxicol. Sci.* **87**, 353–364.
- Desbrow, C., Routledge, E., Brighty, G., Sumpter, J., and Waldock, M. (1998). Identification of estrogenic chemicals in STW effluent. 1. Chemical fractionation and in vitro biological screening. *Environ. Sci. Technol.* **32**, 1549–1558.
- DeVito, M., Biegel, L., Brouwer, A., Brown, S., Brucker-Davis, F., Cheek, A. O., Christensen, R., Colborn, T., Cooke, P., Crissman, J., *et al.* (1999). Screening methods for thyroid hormone disruptors. *Environ. Health Perspect.* **107**, 407–415.
- Dorfman, R. I., and Dorfman, A. S. (1962). Assay of subcutaneously administered androgens on the chick's comb. *Acta. Endocrinol.* **41**, 101–106.
- Dostal, L. A., Chapin, R. E., Stefanski, S. A., Harris, M. W., and Schwetz, B. A. (1988). Testicular toxicity and reduced Sertoli cell numbers in neonatal rats by di(2-ethylhexyl)phthalate and the recovery of fertility as adults. *Toxicol. Appl. Pharmacol.* **95**, 104–121.
- Durhan, E. J., Lambright, C., Wilson, V., Butterworth, B. C., Kuehl, O. W., Orlando, E. F., Guillelte, L. J., Jr., Gray, L. E., and Ankley, G. T. (2002). Evaluation of androstenedione as an androgenic component of river water downstream of a pulp and paper mill effluent. *Environ. Toxicol. Chem. SETAC* **21**, 1973–1976.
- Durhan, E. J., Lambright, C. S., Makynen, E. A., Lazorchak, J., Hartig, P. C., Wilson, V. S., Gray, L. E., and Ankley, G. T. (2006). Identification of metabolites of trenbolone acetate in androgenic runoff from a beef feedlot. *Environ. Health Perspect.* **114**(Suppl. 1), 65–68.
- Ellis, R. J., van den Heuvel, M. R., Bandelj, E., Smith, M. A., McCarthy, L. H., Stuthridge, T. R., and Dietrich, D. R. (2003). In vivo and in vitro assessment of the androgenic potential of a pulp and paper mill effluent. *Environ. Toxicol. Chem. SETAC* **22**, 1448–1456.
- European Commission. (1996). *European Workshop on the Impact of Endocrine Disrupters on Human Health and Wildlife*. Proceedings from a workshop, December 2–4, 1996, Weybridge, UK. Report reference EUR 17549. Brussels, Belgium: European Commission.
- Felner, E. I., and White, P. C. (2000). Prepubertal gynecomastia: Indirect exposure to estrogen cream. *Pediatrics* **105**, E55.
- Foster, P. M. (2006). Disruption of reproductive development in male rat offspring following in utero exposure to phthalate esters. *Int. J. Androl.* **29**, 140–147.



- Foster, P. M., Lake, B. G., Cook, M. W., Thomas, L. V., and Gangolli, S. D. (1981). Structure-activity requirements for the induction of testicular atrophy by butyl phthalates in immature rats: Effect on testicular zinc content. *Adv. Exp. Med. Biol.* **136**(Pt A), 445–452.
- Goldman, J. M., Laws, S. C., Balchak, S. K., Cooper, R. L., and Kavlock, R. J. (2000). Endocrine-disrupting chemicals: Prepubertal exposures and effects on sexual maturation and thyroid activity in the female rat. A focus on the EDSTAC recommendations. *Crit. Rev. Toxicol.* **30**, 135–196.
- Grajewski, B., Whelan, E. A., Schnorr, T. M., Mouradian, R., Alderfer, R., and Wild, D. K. (1996). Evaluation of reproductive function among men occupationally exposed to a stilbene derivative: I. Hormonal and physical status. *Am. J. Ind. Med.* **29**, 49–57.
- Gray, L. E., Jr. (1998a). Tiered screening and testing strategy for xenoestrogens and antiandrogens. *Toxicol. Lett.* **102–103**, 677–680.
- Gray, L. E., Jr. (1998b). Xenoendocrine disrupters: Laboratory studies on male reproductive effects. *Toxicol. Lett.* **102–103**, 331–335.
- Gray, L. E., Jr., Kelce, W. R., Monosson, E., Ostby, J. S., and Birnbaum, L. S. (1995). Exposure to TCDD during development permanently alters reproductive function in male Long Evans rats and hamsters: Reduced ejaculated and epididymal sperm numbers and sex accessory gland weights in offspring with normal androgenic status. *Toxicol. Appl. Pharmacol.* **131**, 108–118.
- Gray, L. E., Jr., Kelce, W. R., Wiese, T., Tyl, R., Gaido, K., Cook, J., Klinefelter, G., Desaulniers, D., Wilson, E., Zacharewski, T., et al. (1997a). Endocrine Screening Methods Workshop report: Detection of estrogenic and androgenic hormonal and antihormonal activity for chemicals that act via receptor or steroidogenic enzyme mechanisms. *Reprod. Toxicol.* **11**, 719–750.
- Gray, L. E., Lambright, C., Parks, L., Tyl, R. W., Orlando, E. F., Guillette, L. J., Wolf, C. J., Seeley, J. C., Chang, T.-Y., Wilson, V., et al., Eds. (2001a). *Antiandrogenic Effects of Environmental Endocrine Disruptors*. Springer-Verlag, New York.
- Gray, L. E., Jr., and Ostby, J. S. (1995). In utero 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) alters reproductive morphology and function in female rat offspring. *Toxicol. Appl. Pharmacol.* **133**, 285–294.
- Gray, L. E., Jr., and Ostby, J. (1998). Effects of pesticides and toxic substances on behavioral and morphological reproductive development: Endocrine versus nonendocrine mechanisms. *Toxicol. Ind. Health* **14**, 159–184.
- Gray, L. E., Jr., Ostby, J., Ferrell, J., Rehnberg, G., Linder, R., Cooper, R., Goldman, J., Slott, V., and Laskey, J. (1989). A dose-response analysis of methoxychlor-induced alterations of reproductive development and function in the rat. *Fundam. Appl. Toxicol.* **12**, 92–108.
- Gray, L. E., Jr., Ostby, J. S., Ferrell, J. M., Sigmon, E. R., and Goldman, J. M. (1988b). Methoxychlor induces estrogen-like alterations of behavior and the reproductive tract in the female rat and hamster: Effects on sex behavior, running wheel activity, and uterine morphology. *Toxicol. Appl. Pharmacol.* **96**, 525–540.
- Gray, L. E., Jr., Ostby, J., Furr, J., Price, M., Veeramachaneni, D. N., and Parks, L. (2000). Perinatal exposure to the phthalates DEHP, BBP, and DINP, but not DEP, DMP, or DOTP, alters sexual differentiation of the male rat. *Toxicol. Sci.* **58**, 350–365.
- Gray, L. E., Ostby, J., Furr, J., Price, M., Wolf, C. J., Lambright, C., Parks, L., Veeramachaneni, D. N. R., Wilson, V., Hotchkiss, A., Orlando, E., and Guillette, L. (2001a). Effects of environmental antiandrogens in experimental animals. *Hum. Reprod. Update* **7**, 248–264.
- Gray, L. E., Ostby, J., Furr, J., Price, M., Wolf, C. J., Lambright, C., Parks, L., Veeramachaneni, D. N. R., Wilson, V., Hotchkiss, A., et al. (2001b). Effects of environmental antiandrogens in experimental animals. *Hum. Reprod. Update* **7**, 248–264.
- Gray, L. E., Jr., Ostby, J. S., and Kelce, W. R. (1994). Developmental effects of an environmental antiandrogen: The fungicide vinclozolin alters sex differentiation of the male rat. *Toxicol. Appl. Pharmacol.* **129**, 46–52.
- Gray, L. E., Ostby, J. S., and Kelce, W. R. (1997b). A dose-response analysis of the reproductive effects of a single gestational dose of 2,3,7,8-tetrachlorodibenzo-p-dioxin in male Long Evans Hooded rat offspring. *Toxicol. Appl. Pharmacol.* **146**, 11–20.
- Gray, L. E., Jr., Ostby, J., Monosson, E., and Kelce, W. R. (1999). Environmental antiandrogens: Low doses of the fungicide vinclozolin alter sexual differentiation of the male rat. *Toxicol. Ind. Health* **15**, 48–64.
- Gray, L. E., Jr., Ostby, J., Sigmon, R., Ferrell, J., Rehnberg, G., Linder, R., Cooper, R., Goldman, J., and Laskey, J. (1988a). The development of a protocol to assess reproductive effects of toxicants in the rat. *Reprod. Toxicol.* **2**, 281–287.
- Gray, L. E., Jr., Wilson, V. S., Stoker, T., Lambright, C., Furr, J., Noriega, N., Howdeshell, K., Ankley, G. T., and Guillette, L. (2006). Adverse effects of environmental antiandrogens and androgens on reproductive development in mammals. *Int. J. Androl.* **29**, 96–104; discussion 105–108.
- Gray, L. E., Wolf, C., Mann, P., and Ostby, J. S. (1997c). In utero exposure to low doses of 2,3,7,8-tetrachlorodibenzo-p-dioxin alters reproductive development of female Long Evans hooded rat offspring. *Toxicol. Appl. Pharmacol.* **146**, 237–244.
- Gray, T. J., Rowland, I. R., Foster, P. M., and Gangolli, S. D. (1982). Species differences in the testicular toxicity of phthalate esters. *Toxicol. Lett.* **11**, 141–147.
- Grumbach, M. M., and Ducharme, J. R. (1960). The effects of androgens on fetal sexual development: Androgen-induced female pseudohermaphroditism. *Fertil. Steril.* **11**, 157–180.
- Guillette, L. J., Jr. (2006). Endocrine disrupting contaminants—beyond the dogma. *Environ. Health Perspect.* **114**(Suppl. 1), 9–12.
- Guo, Y. L., Lambert, G. H., and Hsu, C. C. (1995). Growth abnormalities in the population exposed in utero and early postnatally to polychlorinated biphenyls and dibenzofurans. *Environ. Health Perspect.* **103**(Suppl. 6), 117–122.
- Harries, J. E., Janbakhsh, A., Jobling, S., Mattiessen, P., Sumpter, J. P., and Tyler, C. R. (1999). Estrogenic potency of effluent from two sewage treatment works in the United Kingdom. *Environ. Toxicol. Chem.* **18**, 932–937.
- Hawkins, M. B., Godwin, J., Crews, D., and Thomas, P. (2005). The distributions of the duplicate oestrogen receptors ER-beta a and ER-beta b in the forebrain of the Atlantic croaker (*Micropogonias undulatus*): Evidence for subfunctionalization after gene duplication. *Proceedings* **272**, 633–641.
- Hayes, W. J., and Laws, E. R. (1991). In *Handbook of Pesticide Toxicology*. Academic Press, Inc, San Diego.
- Heneweer, M., van den Berg, M., and Sanderson, J. T. (2004). A comparison of human H295R and rat R2C cell lines as in vitro screening tools for effects on aromatase. *Toxicol. Lett.* **146**, 183–194.
- Henley, D. V., Lipson, N., Korach, K. S., and Bloch, C. A. (2007). Prepubertal gynecomastia linked to lavender and tea tree oils. *N. Engl. J. Med.* **356**, 479–485.
- Hershberger, L. G., Shipley, E. G., and Meyer, R. K. (1953). Myotrophic activity of 19-nortestosterone and other steroids determined by modified levator ani muscle method. *Proc. Soc. Exp. Biol. Med. Soc. Exp. Biol. Med.* **83**, 175–180.
- Higuchi, T. T., Palmer, J. S., Gray, L. E., Jr., and Veeramachaneni, D. N. (2003). Effects of dibutyl phthalate in male rabbits following in utero, adolescent, or postpubertal exposure. *Toxicol. Sci.* **72**, 301–313.
- Horiguchi, T. (2006). Masculinization of female gastropod mollusks induced by organotin compounds, focusing on mechanism of actions of tributyltin and triphenyltin for development of imposex. *Environ. Sci.* **13**, 77–87.
- Hornshaw, T. C., Aulerich, R. J., and Johnson, H. E. (1983). Feeding Great Lakes fish to mink: Effects on mink and accumulation and elimination of PCBS by mink. *J. Toxicol. Environ. Health* **11**, 933–946.
- Hosokawa, S., Murakami, M., Ineyama, M., Yamada, T., Yoshitake, A., Yamada, H., and Miyamoto, J. (1993). The affinity of procymidone to androgen receptor in rats and mice. *J. Toxicol. Sci.* **18**, 83–93.

- Hotchkiss, A. K., Furr, J., Makynen, E. A., Ankley, G. T., and Gray, L. E., Jr. (2007). In utero exposure to the environmental androgen trenbolone masculinizes female Sprague-Dawley rats. *Toxicol. Lett.* **174**, 31–41.
- Hotchkiss, A. K., and Nelson, R. J. (2007). An environmental androgen, trenbolone, affects delayed-type hypersensitivity and reproductive tissues in male mice. *J. Toxicol. Environ. Health* **70**, 138–140.
- Hotchkiss, A. K., Parks-Saldutti, L. G., Ostby, J. S., Lambright, C., Furr, J., Vandenberg, J. G., and Gray, L. E., Jr. (2004). A mixture of the “antiandrogens” linuron and butyl benzyl phthalate alters sexual differentiation of the male rat in a cumulative fashion. *Biol. Reprod.* **71**, 1852–1861.
- Howdeshell, K. L. (2002). A model of the development of the brain as a construct of the thyroid system. *Environ. Health Perspect.* **110**(Suppl. 3), 337–348.
- Howdeshell, K. L., Furr, J., Lambright, C. R., Wilson, V. S., Ryan, B. C., and Gray, L. E., Jr. (2007). Gestational and lactational exposure to ethinyl estradiol, but not bisphenol a, decreases androgen-dependent reproductive organ weights and epididymal sperm abundance in the male Long Evans hooded rat. *Toxicol. Sci.* 10.1093/toxsci/kfm306.
- Hutchinson, T. H., Ankley, G. T., Segner, H., and Tyler, C. R. (2006). Screening and testing for endocrine disruption in fish-biomarkers as “signposts,” not “traffic lights,” in risk assessment. *Environ. Health Perspect.* **114**(Suppl. 1), 106–114.
- Inadera, H. (2006). The immune system as a target for environmental chemicals: Xenoestrogens and other compounds. *Toxicol. Lett.* **164**, 191–206.
- Jenkins, R. L., Wilson, E. M., Angus, R. A., Howell, W. M., and Kirk, M. (2003). Androstenedione and progesterone in the sediment of a river receiving paper mill effluent. *Toxicol. Sci.* **73**, 53–59.
- Jensen, K. M., Makynen, E. A., Kahl, M. D., and Ankley, G. T. (2006). Effects of the feedlot contaminant 17 $\alpha$ -trenbolone on reproductive endocrinology of the fathead minnow. *Environ. Sci. Technol.* **40**, 3112–3117.
- Jobling, S., Beresford, N., Nolan, M., Rodgers-Gray, T., Brighty, G. C., Sumpter, J. P., and Tyler, C. R. (2002a). Altered sexual maturation and gamete production in wild roach (*Rutilus rutilus*) living in rivers that receive treated sewage effluents. *Biol. Reprod.* **66**, 272–281.
- Jobling, S., Coey, S., Whitmore, J. G., Kime, D. E., Van Look, K. J. W., McAllister, B. G., Beresford, N., Henshaw, A. C., Brighty, G., Tyler, C. R., et al. (2002b). Wild intersex roach (*Rutilus rutilus*) have reduced fertility. *Biol. Reprod.* **67**, 515–524.
- Joffe, M. (2003). Infertility and environmental pollutants. *Br. Med. Bull.* **68**, 47–70.
- Johansson, M., Nilsson, S., and Lund, B. O. (1998). Interactions between methylsulfonyl PCBs and the glucocorticoid receptor. *Environ. Health Perspect.* **106**, 769–772.
- Jorundsdottir, K., Svavarsson, J., and Leung, K. M. (2005). Imposed levels in the dogwhelk *Nucella lapillus* (L.)—Continuing improvement at high latitudes. *Mar. Pollut. Bull.* **51**, 744–749.
- Kelce, W. R., Lambright, C. R., Gray, L. E., Jr., and Roberts, K. P. (1997). Vinclozolin and p,p'-DDE alter androgen-dependent gene expression: In vivo confirmation of an androgen receptor-mediated mechanism. *Toxicol. Appl. Pharmacol.* **142**, 192–200.
- Kelce, W. R., Stone, C. R., Laws, S. C., Gray, L. E., Kemppainen, J. A., and Wilson, E. M. (1995). Persistent DDT metabolite p,p'-DDE is a potent androgen receptor antagonist. *Nature* **375**, 581–585.
- Kidd, K. A., Blanchfield, P. J., Mills, K. H., Palace, V. P., Evans, R. E., Lazorchak, J. M., and Flick, R. W. (2007). Collapse of a fish population after exposure to a synthetic estrogen. *Proc Natl Acad Sci U S A* **104**, 8897–8901.
- Kim, Y., Jung, K., Hwang, T., Jung, G., Kim, H., Park, J., Kim, J., Park, J., Park, D., Park, S., et al. (1996). Hematopoietic and reproductive hazards of Korean electronic workers exposed to solvents containing 2-bromopropane. *Scand. J. Work Environ. Health* **22**, 387–391.
- Klonisch, T., Fowler, P. A., and Hombach-Klonisch, S. (2004). Molecular and genetic regulation of testis descent and external genitalia development. *Dev. Biol.* **270**, 1–18.
- Kolpin, D. W., Furlong, E. T., Meyer, M. T., Thurman, E. M., Zaugg, S. D., Barber, L. B., and Buxton, H. T. (2002). Pharmaceuticals, hormones, and other organic wastewater contaminants in US streams, 1999–2000: A national reconnaissance. *Environ. Sci. Technol.* **36**, 1202–1211.
- Kretschmer, X. C., and Baldwin, W. S. (2005). CAR and PXR: Xenosensors of endocrine disrupters? *Chem. Biol. Interact.* **155**, 111–128.
- Kunz, G. J., Klein, K. O., Clemons, R. D., Gottschalk, M. E., and Jones, K. L. (2004). Virilization of young children after topical androgen use by their parents. *Pediatrics* **114**, 282–284.
- Lag, M., Soderlund, E. J., Brunborg, G., Dahl, J. E., Holme, J. A., Omichinski, J. G., Nelson, S. D., and Dybing, E. (1989). Species differences in testicular necrosis and DNA damage, distribution and metabolism of 1,2-dibromo-3-chloropropane (DBCP). *Toxicology* **58**, 133–144.
- Lake, B. G., Brantom, P. G., Gangolli, S. D., Butterworth, K. R., and Grasso, P. (1976). Studies on the effects of orally administered di-(2-ethylhexyl) phthalate in the ferret. *Toxicology* **6**, 341–356.
- Lambright, C., Ostby, J., Bobseine, K., Wilson, V. S., Hotchkiss, A. K., Mann, P. C., and Gray, L. E. J. (2000). Cellular and molecular mechanisms of action of linuron: An antiandrogenic herbicide that produces reproductive malformations in male rats. *Toxicol. Sci.* **56**, 389–399.
- Larsson, D. G., and Forlin, L. (2002). Male-biased sex ratios of fish embryos near a pulp mill: Temporary recovery after a short-term shutdown. *Environ. Health Perspect.* **110**, 739–742.
- LeMaire, W. J., Cleveland, W. W., Bejar, R. L., Marsh, J. M., and Fishman, L. (1972). Aminoglutethimide: A possible cause of pseudohermaphroditism in females. *Am. J. Dis. Child.* **124**, 421–423.
- Louis, G. M., Weiner, J. M., Whitcomb, B. W., Sperrazza, R., Schisterman, E. F., Lobdell, D. T., Crickard, K., Greizerstein, H., and Kostyniak, P. J. (2005). Environmental PCB exposure and risk of endometriosis. *Hum. Reprod.* **20**, 279–285.
- Mably, T. A., Bjerke, D. L., Moore, R. W., Gendron-Fitzpatrick, A., and Peterson, R. E. (1992). In utero and lactational exposure of male rats to 2,3,7,8-tetrachlorodibenzo-p-dioxin. 3. Effects on spermatogenesis and reproductive capability. *Toxicol. Appl. Pharmacol.* **114**, 118–126.
- Martinovic, D., Hogarth, W. T., Jones, R. E., and Sorensen, P. W. (2007). Environmental estrogens suppress hormones, behavior, and reproductive fitness in male fathead minnows. *Environ. Toxicol. Chem. SETAC* **26**, 271–278.
- McIntyre, B. S., Barlow, N. J., Wallace, D. G., Maness, S. C., Gaido, K. W., and Foster, P. M. (2000). Effects of in utero exposure to linuron on androgen-dependent reproductive development in the male Crl:CD(SD)BR rat. *Toxicol. Appl. Pharmacol.* **167**, 87–99.
- Meyer, K., Usleber, E., Martlbauer, E., and Bauer, J. (2000). [Occurrence of zearalenone, alpha- and beta-zearalenol in bile of breeding sows in relation to reproductive performance]. *Berl. Munch. Tierarztl. Wochenschr.* **113**, 374–379.
- Mizota, K., and Ueda, H. (2006). Endocrine disrupting chemical atrazine causes degeneration through Gq/11 protein-coupled neurosteroid receptor in mast cells. *Toxicol. Sci.* **90**, 362–368.
- Monosson, E., Kelce, W. R., Lambright, C., Ostby, J., and Gray, L. E., Jr. (1999). Peripubertal exposure to the antiandrogenic fungicide, vinclozolin, delays puberty, inhibits the development of androgen-dependent tissues, and alters androgen receptor function in the male rat. *Toxicol. Ind. Health* **15**, 65–79.
- Mylchreest, E., Cattley, R. C., and Foster, P. M. (1998). Male reproductive tract malformations in rats following gestational and lactational exposure to di(n-butyl) phthalate: An antiandrogenic mechanism? *Toxicol. Sci.* **43**, 47–60.

- Nader, N., Raverot, G., Emptoz-Bonneton, A., Dechaud, H., Bonnay, M., Baudin, E., and Pugeat, M. (2006). Mitotane has an estrogenic effect on sex hormone-binding globulin and corticosteroid-binding globulin in humans. *J. Clin. Endocrinol. Metab.* **91**, 2165–2170.
- Nellemann, C., Dalgaard, M., Lam, H. R., and Vinggaard, A. M. (2003). The combined effects of vinclozolin and procymidone do not deviate from expected additivity in vitro and in vivo. *Toxicol. Sci.* **71**, 251–262.
- Nishikawa, J., Mamiya, S., Kanayama, T., Nishikawa, T., Shiraishi, F., and Horiguchi, T. (2004). Involvement of the retinoid X receptor in the development of imposex caused by organotins in gastropods. *Environ. Sci. Technol.* **38**, 6271–6276.
- Noriega, N. C., Ostby, J., Lambright, C., Wilson, V. S., and Gray, L. E., Jr. (2005). Late gestational exposure to the fungicide prochloraz delays the onset of parturition and causes reproductive malformations in male but not female rat offspring. *Biol. Reprod.* **72**, 1324–1335.
- Opitz, R., Hartmann, S., Blank, T., Braunbeck, T., Lutz, I., and Kloas, W. (2006). Evaluation of histological and molecular endpoints for enhanced detection of thyroid system disruption in *Xenopus laevis* tadpoles. *Toxicol. Sci.* **90**, 337–348.
- Orlando, E. F., Bass, D. E., Caltabiano, L. M., Davis, W. P., Gray, L. E., Jr., and Guillelte, L. J., Jr. (2007). Altered development and reproduction in mosquitofish exposed to pulp and paper mill effluent in the Fenholloway River, Florida USA. *Aquat. Toxicol.* **84**, 399–405.
- Orlando, E. F., Kolok, A. S., Binzick, G. A., Gates, J. L., Horton, M. K., Lambright, C. S., Gray, L. E., Jr., Soto, A. M., and Guillelte, L. J., Jr. (2004). Endocrine-disrupting effects of cattle feedlot effluent on an aquatic sentinel species, the fathead minnow. *Environ. Health Perspect.* **112**, 353–358.
- Ostby, J., Kelce, W. R., Lambright, C., Wolf, C. J., Mann, P., and Gray, L. E., Jr. (1999). The fungicide procymidone alters sexual differentiation in the male rat by acting as an androgen-receptor antagonist in vivo and in vitro. *Toxicol. Ind. Health* **15**, 80–93.
- Owens, W., Gray, L. E., Zeiger, E., Walker, M., Yamasaki, K., Ashby, J., and Jacob, E. (2007). The OECD program to validate the rat Hershberger bioassay to screen compounds for in vivo androgen and antiandrogen responses: Phase 2 dose-response studies. *Environ. Health Perspect.* **115**, 671–678.
- Owens, W., and Koeter, H. B. (2003). The OECD program to validate the rat uterotrophic bioassay: An overview. *Environ. Health Perspect.* **111**, 1527–1529.
- Owens, W., Zeiger, E., Walker, M., Ashby, J., Onyon, L., and Gray, L. E., Jr. (2006). The OECD program to validate the rat Hershberger bioassay to screen compounds for in vivo androgen and antiandrogen responses. Phase 1: Use of a potent agonist and a potent antagonist to test the standardized protocol. *Environ. Health Perspect.* **114**, 1259–1265.
- Pakdel, F., Metivier, R., Flouriot, G., and Valotaire, Y. (2000). Two estrogen receptor (ER) isoforms with different estrogen dependencies are generated from the trout ER gene. *Endocrinology*. **141**, 571–580.
- Parks, L. G., Lambright, C. S., Orlando, E. F., Guillelte, L. J., Jr., Ankley, G. T., and Gray, L. E., Jr. (2001). Masculinization of female mosquitofish in Kraft mill effluent-contaminated Fenholloway River water is associated with androgen receptor agonist activity. *Toxicol. Sci.* **62**, 257–267.
- Parks, L. G., Ostby, J. S., Lambright, C. R., Abbott, B. D., Klinefelter, G. R., Barlow, N. J., and Gray, L. E., Jr. (2000). The plasticizer diethylhexyl phthalate induces malformations by decreasing fetal testosterone synthesis during sexual differentiation in the male rat. *Toxicol. Sci.* **58**, 339–349.
- Payne, J., Rajapakse, N., Wilkins, M., and Kortenkamp, A. (2000). Prediction and assessment of the effects of mixtures of four xenoestrogens. *Environ. Health Perspect.* **108**, 983–987.
- Payne, J., Scholze, M., and Kortenkamp, A. (2001). Mixtures of four organochlorines enhance human breast cancer cell proliferation. *Environ. Health Perspect.* **109**, 391–397.
- Peterson, G. I., Norrgren, L., Holbech, H., Lundgren, A., and Kovisto, S. (2001). Suitability of Zebrafish as a Test Organism for Detection of Endocrine-Disrupting Chemicals. Vol. 2001. Nordic Council of Ministers, Uppsala.
- Potashnik, G., and Porath, A. (1995). Dibromochloropropane (DBCP): A 17-year reassessment of testicular function and reproductive performance. *J. Occup. Environ. Med.* **37**, 1287–1292.
- Potashnik, G., and Yanai-Inbar, I. (1987). Dibromochloropropane (DBCP): An 8-year reevaluation of testicular function and reproductive performance. *Fertil. Steril.* **47**, 317–323.
- Quigley, C. A., De Bellis, A., Marschke, K. B., el-Awady, M. K., Wilson, E. M., and French, F. S. (1995). Androgen receptor defects: Historical, clinical, and molecular perspectives. *Endocr. Rev.* **16**, 271–321.
- Raudenbush, S. W., and Xiao-Feng, L. (2001). Effects of study duration, frequency of observation, and sample size on power in studies of group differences in polynomial change. *Psychol. Methods* **6**, 387–401.
- Reine, N. J. (2007). Medical management of pituitary-dependent hyperadrenocorticism: Mitotane versus trilostane. *Clin. Tech. Small Anim. Pract.* **22**, 18–25.
- Rider, C. V., Furr, J., Wilson, V. S., and Gray, L. E., Jr. (2008). A mixture of seven antiandrogens induces reproductive malformations in rats. *Int. J. Androl.* In press.
- Rier, S. E., Turner, W. E., Martin, D. C., Morris, R., Lucier, G. W., and Clark, G. C. (2001). Serum levels of TCDD and dioxin-like chemicals in Rhesus monkeys chronically exposed to dioxin: Correlation of increased serum PCB levels with endometriosis. *Toxicol. Sci.* **59**, 147–159.
- Rodriguez, E. M., Medesani, D. A., and Fingerman, M. (2007). Endocrine disruption in crustaceans due to pollutants: A review. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **146**, 661–671.
- Sanderson, J. T., Elliott, J. E., Norstrom, R. J., Whitehead, P. E., Hart, L. E., Cheng, K. M., and Bellward, G. D. (1994). Monitoring biological effects of polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls in great blue heron chicks (*Ardea herodias*) in British Columbia. *J. Toxicol. Environ. Health* **41**, 435–450.
- Saunders, F. J. (1968). Effects of sex steroids and related compounds on pregnancy and on development of the young. *Physiol. Rev.* **48**, 601–643.
- Schantz, S. L., and Widholm, J. J. (2001). Cognitive effects of endocrine-disrupting chemicals in animals. *Environ. Health Perspect.* **109**, 1197–1206.
- Seki, M., Fujishima, S., Nozaka, T., Maeda, M., and Kobayashi, K. (2006). Comparison of response to 17 beta-estradiol and 17 beta-trenbolone among three small fish species. *Environ. Toxicol. Chem. SETAC* **25**, 2742–2752.
- Selgrade, M. K. (2007). Immunotoxicity—The risk is real. *Toxicol. Sci.*
- Sharpe, R. M. (2006). Pathways of endocrine disruption during male sexual differentiation and masculinization. *Best Pract. Res. Clin. Endocrinol. Metab.* **20**, 91–110.
- Shultz, S., Baral, H. S., Charman, S., Cunningham, A. A., Das, D., Ghalsasi, G. R., Goudar, M. S., Green, R. E., Jones, A., Nighot, P., et al. (2004). Diclofenac poisoning is widespread in declining vulture populations across the Indian subcontinent. *Proceedings* **271**(Suppl. 6), S458–S460.
- Silva, E., Rajapakse, N., and Kortenkamp, A. (2002). Something from “nothing”—Eight weak estrogenic chemicals combined at concentrations below NOECs produce significant mixture effects. *Environ. Sci. Technol.* **36**, 1751–1756.
- Silva, M. J., Barr, D. B., Reidy, J. A., Malek, N. A., Hodge, C. C., Caudill, S. P., Brock, J. W., Needham, L. L., and Calafat, A. M. (2004). Urinary levels of seven phthalate metabolites in the U.S. population from the National Health and Nutrition Examination Survey (NHANES) 1999–2000. *Environ. Health Perspect.* **112**, 331–338.



- Sjoberg, P., Lindqvist, N. G., and Ploen, L. (1986). Age-dependent response of the rat testes to di(2-ethylhexyl) phthalate. *Environ. Health Perspect.* **65**, 237–242.
- Skakkebaek, N. E., Rajpert-De Meyts, E., and Main, K. M. (2001). Testicular dysgenesis syndrome: An increasingly common developmental disorder with environmental aspects. *Hum. Reprod.* **16**, 972–978.
- Slutsky, M., Levin, J. L., and Levy, B. S. (1999). Azoospermia and oligospermia among a large cohort of DBCP applicators in 12 countries. *Int. J. Occup. Environ. Health* **5**, 116–122.
- Smith, E. R., and Quinn, M. M. (1992). Uterotropic action in rats of amsonic acid and three of its synthetic precursors. *J. Toxicol. Environ. Health* **36**, 13–25.
- Soto, A. M., Calabro, J. M., Precht, N. V., Yau, A. Y., Orlando, E. F., Daxenberger, A., Kolok, A. S., Guille, L. J., Jr., le Bizet, B., Lange, I. G., et al. (2004). Androgenic and estrogenic activity in water bodies receiving cattle feedlot effluent in Eastern Nebraska, USA. *Environ. Health Perspect.* **112**, 346–352.
- Sparagana, M. (1987). Primary hypogonadism associated with o,p' DDD (mitotane) therapy. *J. Toxicol.* **25**, 463–472.
- Spooner, N., Gibbs, P. E., Bryan, G. W., and Goad, L. J. (1991). The effect of tributyltin upon steroid titres in female dogwhelk, *nucella lapillus*, and the development of imposex. *Mar. Environ. Res.* **32**, 37–49.
- Steinberger, E., and Lloyd, J. A. (1985). In *Chemicals Affecting the Development of Reproductive Capacity*. Raven Press, New York.
- Sternberg, R. M., Hotchkiss, A. K., and Leblanc, G. A. (2008). The contribution of steroidal androgens and estrogens to reproductive maturation of the eastern mud snail *Ilyanassa obsoleta*. *Gen. Comp. Endocrinol.* 10.1016/j.ygcen.2007.12.002
- Stoker, T. E., Cooper, R. L., Lambright, C. S., Wilson, V. S., Furr, J., and Gray, L. E. (2005). In vivo and in vitro anti-androgenic effects of DE-71, a commercial polybrominated diphenyl ether (PBDE) mixture. *Toxicol. Appl. Pharmacol.* **207**, 78–88.
- Stoker, T. E., Parks, L. G., Gray, L. E., and Cooper, R. L. (2000). Endocrine-disrupting chemicals: Prepubertal exposures and effects on sexual maturation and thyroid function in the male rat. A focus on the EDSTAC recommendations. *Crit. Rev. Toxicol.* **30**, 197–202.
- Takeo, J., and Yamashita, S. (1999). Two distinct isoforms of cDNA encoding rainbow trout androgen receptors. *J. Biol. Chem.* **274**, 5674–5680.
- Tabb, M. M., and Blumberg, B. (2006). New modes of action for endocrine-disrupting chemicals. *Mol. Endocrinol.* **20**, 475–482.
- Teitelbaum, D. T. (1999). The toxicology of 1,2-dibromo-3-chloropropane (DBCP): A brief review. *Int. J. Occup. Environ. Health* **5**, 122–126.
- Tiwary, C. M. (1998). Premature sexual development in children following the use of estrogen- or placenta-containing hair products. *Clin. Pediatrics* **37**, 733–739.
- Touart, L. W. (2004). Factors considered in using birds for evaluating endocrine-disrupting chemicals. *ILAR J.* **45**, 462–468.
- Warren, D. W., Ahmad, N., and Rudeen, P. K. (1988). The effects of fetal exposure to 1,2-dibromo-3-chloropropane on adult male reproductive function. *Biol. Reprod.* **39**, 707–716.
- Waterman, S. J., Keller, L. H., Trimmer, G. W., Freeman, J. J., Nikiforov, A. I., Harris, S. B., Nicolich, M. J., and McKee, R. H. (2000). Two-generation reproduction study in rats given di-isooctyl phthalate in the diet. *Reprod. Toxicol.* **14**, 21–36.
- Whelan, E. A., Grajewski, B., Wild, D. K., Schnorr, T. M., and Alderfer, R. (1996). Evaluation of reproductive function among men occupationally exposed to a stilbene derivative: II. Perceived libido and potency. *Am. J. Ind. Med.* **29**, 59–65.
- WHO (2002). ICPS global assessment of the state-of-the-science of endocrine disruptors. In *International Programme on Chemical Safety, WHO/PCS/EDC/02.2* (T. Damstra, S. Barlow, A. Bergman, R. Kavlock, and G. v d Kraak, Eds.). World Health Organization, Geneva, Switzerland.
- Whorton, D., Milby, T. H., Krauss, R. M., and Stubbs, H. A. (1979). Testicular function in DBCP exposed pesticide workers. *J. Occup. Med.* **21**, 161–166.
- Whorton, M. D., and Foliat, D. E. (1983). Mutagenicity, carcinogenicity and reproductive effects of dibromochloropropane (DBCP). *Mutat. Res.* **123**, 13–30.
- Whorton, M. D., and Milby, T. H. (1980). Recovery of testicular function among DBCP workers. *J. Occup. Med.* **22**, 177–179.
- Wilson, V. S., Cardon, M. C., Thornton, J., Korte, J. J., Ankley, G. T., Welch, J., Gray, L. E., Jr., and Hartig, P. C. (2004). Cloning and in vitro expression and characterization of the androgen receptor and isolation of estrogen receptor alpha from the fathead Minnow (*Pimephales promelas*). *Environ. Sci. Technol.* **38**, 6314–6321.
- Wilson, V. S., Bobseine, K., and Gray, L. E. J. (2004a). Development and characterization of a cell line that stably expresses an estrogen-responsive luciferase reporter for the detection of estrogen receptor agonist and antagonists. *Toxicol. Sci.* **81**, 69–77.
- Wilson, V. S., Bobseine, K., Lambright, C. R., and Gray, L. E., Jr. (2002a). A novel cell line, MDA-kb2, that stably expresses an androgen- and glucocorticoid-responsive reporter for the detection of hormone receptor agonists and antagonists. *Toxicol. Sci.* **66**, 69–81.
- Wilson, V. S., Cardon, M. C., Gray, L. E., Jr., and Hartig, P. C. (2007). Competitive binding comparison of endocrine-disrupting compounds to recombinant androgen receptor from fathead minnow, rainbow trout, and human. *Environ. Toxicol. Chem. SETAC* **26**, 1793–1802.
- Wilson, V. S., Cardon, M. C., Thornton, J., Korte, J. J., Ankley, G. T., Welch, J., Gray, L. E., Jr., and Hartig, P. C. (2004b). Cloning and in vitro expression and characterization of the androgen receptor and isolation of estrogen receptor alpha from the fathead Minnow (*Pimephales promelas*). *Environ. Sci. Technol.* **38**, 6314–6321.
- Wilson, V. S., Lambright, C., Furr, J., Ostby, J., Wood, C., Held, G., and Gray, L. E., Jr. (2004c). Phthalate ester-induced gubernacular lesions are associated with reduced *insl3* gene expression in the fetal rat testis. *Toxicol. Lett.* **146**, 207–215.
- Wilson, V. S., Lambright, C., Ostby, J., and Gray, L. E., Jr. (2002b). In vitro and in vivo effects of 17beta-trenbolone: A feedlot effluent contaminant. *Toxicol. Sci.* **70**, 202–211.
- Wolf, C. J., Ostby, J. S., and Gray, L. E., Jr. (1999). Gestational exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) severely alters reproductive function of female hamster offspring. *Toxicol. Sci.* **51**, 259–264.
- Yamasaki, K., Sawaki, M., Ohta, R., Okuda, H., Katayama, S., Yamada, T., Ohta, T., Kosaka, T., and Owens, W. (2003). OECD validation of the Hershberger assay in Japan: Phase 2 dose response of methyltestosterone, vinclozolin, and p,p'-DDE. *Environ. Health Perspect.* **111**, 1912–1919.
- Young, L. E. (2001). Imprinting of genes and the Barker hypothesis. *Twin Res.* **4**, 307–317.
- Zabel, E. W., Walker, M. K., Hornung, M. W., Clayton, M. K., and Peterson, R. E. (1995). Interactions of polychlorinated dibenzo-p-dioxin, dibenzofuran, and biphenyl congeners for producing rainbow trout early life stage mortality. *Toxicol. Appl. Pharmacol.* **134**, 204–213.
- Zoeller, R. T., and Crofton, K. M. (2005). Mode of action: Developmental thyroid hormone insufficiency—Neurological abnormalities resulting from exposure to propylthiouracil. *Crit. Rev. Toxicol.* **35**, 771–781.