

2011

Moving micronutrients from the soil to the seeds: Genes and physiological processes from a biofortification perspective


Brian M. Waters

University of Nebraska-Lincoln, bwaters2@unl.edu

Renuka P. Sankaran

Lehman College, Bronx, NY

Follow this and additional works at: <https://digitalcommons.unl.edu/agronomyfacpub>

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), [Botany Commons](#), [Horticulture Commons](#), [Other Plant Sciences Commons](#), and the [Plant Biology Commons](#)

Waters, Brian M. and Sankaran, Renuka P., "Moving micronutrients from the soil to the seeds: Genes and physiological processes from a biofortification perspective" (2011). *Agronomy & Horticulture -- Faculty Publications*. 850.
<https://digitalcommons.unl.edu/agronomyfacpub/850>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Moving micronutrients from the soil to the seeds: Genes and physiological processes from a biofortification perspective

Brian M. Waters¹ and Renuka P. Sankaran²

1. Department of Agronomy and Horticulture, University of Nebraska, Lincoln, NE 68583-0915, USA

2. Department of Biological Sciences, Lehman College, Bronx, NY 10468, USA

Corresponding author – Brian M. Waters, email bwaters2@unl.edu

Abstract

The micronutrients iron (Fe), zinc (Zn), and copper (Cu) are essential for plants and the humans and animals that consume plants. Increasing the micronutrient density of staple crops, or biofortification, will greatly improve human nutrition on a global scale. This review discusses the processes and genes needed to translocate micronutrients through the plant to the developing seeds, and potential strategies for developing biofortified crops.

Keywords: iron, zinc, copper, remobilization, micronutrient transporters, biofortification

Contents

1. Introduction	563
2. What are the sources of seed micronutrients?	563
2.1. Which tissues supply micronutrients to seeds, and at which developmental stages?	563
2.2. Remobilization or continuous uptake?	563
3. Defining the pathway of mineral micronutrients through the plant body	564
3.1. Uptake from rhizosphere	564
3.2. Xylem loading and transfer from root to shoot	565
3.3. Translocation to leaf or seed covering tissues	566
3.4. Phloem loading for translocation to seeds	566
3.5. Gradient to the seed and co-transport with N	567
3.6. Transport into seeds and storage forms	567
4. Future discovery	567
4.1. Quantitative genetics	568
4.2. Candidate genes by expression pattern and predicted function	568
5. Strategies for biofortification of mineral micronutrients	569
6. Conclusion	569
Acknowledgments	569
References	569

1. Introduction

Seeds are major sources of human and animal foods, and also the input material for cultivating most agricultural crops. Seed quality determines suitability of seeds for both of these purposes. One component of seed quality is chemical composition, such as the concentrations of mineral elements, including micronutrients such as Fe, Zn, and Cu. The maternal plant supplies the metabolic building blocks that ultimately make up the seed. These inputs are influenced by environmental conditions and by the genetic makeup of the maternal plant. An increased understanding of control over supply of nutrients to developing seeds will allow development of improved varieties that can respond favorably to maintain seed production and quality in adverse environments. This understanding will also allow production of biofortified seeds with increased concentrations of micronutrients.

Plant derived foods provide an important source of proteins and dietary minerals. This is especially true in developing countries where plant foods are a predominant portion of the diet. The concentrations of some minerals, especially iron, zinc, iodine, and selenium, are inherently low in plants as opposed to animal derived foods. As a result, more than 3 billion people worldwide suffer from micronutrient malnutrition [1,2]. Hence, there is a need to improve the mineral concentrations of important seed crops such as rice (*Oryza sativa*), wheat (*Triticum aestivum* or *T. durum*), maize (*Zea mays*), as well as common bean (*Phaseolus vulgaris*) and other legumes. Biofortification is a recent approach aimed at increasing the bioavailable nutrients, such as Fe and Zn, in these staple crops [3] rather than using fortificants or supplements. However, breeding or developing transgenic varieties for increased micronutrient density will require the enhancement of several processes involving both membrane localized transport proteins and long distance systems. The quantities of minerals in seeds depends on uptake from the rhizosphere into the roots, translocation to the transpiring shoots in the xylem, transfer into leaves or other tissues, and finally, translocation into the seeds in the phloem. A major challenge of biofortification is an incomplete understanding of the pathways and the rate limiting steps involved in translocating minerals to the seeds.

This review will focus on the genes (summarized in fig. 1) and processes in the plant that act to supply nutrients, specifically the mineral micronutrients Fe and Zn, to developing seeds. While Cu in human diets is not limiting on a widespread basis like Fe and Zn, Cu interacts with Fe and Zn, and shares common transport proteins and mechanisms. As such, a discussion of Cu transport is also included in this review.

2. What are the sources of seed micronutrients?

Specific strategies for developing crop varieties with increased seed micronutrient concentration can be developed by knowing which maternal tissues have the greatest influence on final seed composition. With this knowledge, efforts for gene discovery and targeted overexpression strategies can be directed to the most appropriate tissues.

2.1. Which tissues supply micronutrients to seeds, and at which developmental stages?

A series of studies using radioactive Fe and Zn were carried out by nutritionists to determine the best ways to produce labeled food materials for animal feeding studies (table 1) [4–10]. These studies showed that micronutrients can be translocated to seeds from various tissues, hinting at a pathway that includes root,

stem, leaf, and reproductive tissues. Later, more focused studies by plant biologists corroborated these results. In pea (*Pisum sativum*), ^{59}Fe applied to leaflets, stipules, and pod walls was translocated into seeds [11]. In wheat and rice, ^{65}Zn applied to leaf tips was translocated to grain [12,13]. Foliar Zn fertilization was highly effective at increasing Zn concentration in wheat grain, especially when applied at various times during the grain filling period [1,14]. Taken together, these results suggest that the roles of vegetative tissues in supplying micronutrients to grain, and the specific mechanisms involved, should be a continued focus of future research efforts.

2.2. Remobilization or continuous uptake?

During the senescence stage of development, remobilization (i.e., net export of stored or recycled nutrients) of some nutrients occurs from vegetative tissues, such as leaves and stems. At the end of growing seasons, mass senescence typically occurs in leaves and other tissues. These remobilized nutrients are most likely moved to developing seeds in annual crop species, providing that the senescence and seed import are synchronized to provide source-sink relationships. While it is well established that remobilized N is a major source of seed protein components for wheat and barley [15–17], data for other mineral elements is less abundant, although older studies demonstrated remobilization of Cu, Fe, and Zn in legumes [18,19] and wheat [20–22], and some newer studies have found similar results [23,24]. Remobilization of Fe and Zn increased when these micronutrients were withheld from the hydroponic solution post-anthesis, indicating that remobilization mechanisms might be upregulated under nutrient limitation [24]. In this wheat study, flag leaf and lower leaves were the major sources of remobilized micronutrients, in contrast to results from rice, where stems accounted for the major source of Zn remobilized to grain [13]. In another rice study, the authors concluded that remobilization accounted for very little grain Zn, and that uptake after anthesis supplied the majority of Zn to grain, with most Zn passing through stems, and only a small amount passing through leaves [25]. It is possible that differences in plant anatomy between rice and wheat [26] account for these contrasting observations, and highlights the importance of conducting similar experiments in a variety of plant species.

Mineral element concentrations decrease over developmental time in *Arabidopsis thaliana* rosette leaves [27,28]. This clearly suggests remobilization of minerals from these leaves, which may be used to supply seeds. Quantitative analysis of *Arabidopsis* plants over developmental time indicated that remobilization accounted for up to 30% of the total Cu, Fe, and Zn in *Arabidopsis* seeds [28]. Future crop breeding or variety development strategies may include modifications that result in increased efficiency of remobilization to generate biofortified seeds. However, in all of the studies mentioned above, remobilized micronutrient sources did not account for the total quantities of seed micronutrients, suggesting that plants continue to take up and translocate minerals into seeds over the seed filling period. The pattern of Zn and Cd distribution in the plant were similar when *Brassica juncea* plants were simultaneously exposed to both Cd and Zn to examine the transport of Cd to seeds (most likely by Zn transporters, see below). Plants treated with Cd during seed set accumulated the highest concentrations of Cd in seeds, suggesting that uptake during seed fill is an important contributor for seed mineral content [29]. Mineral micronutrients that are supplied to seeds by uptake during seed development must pass through the plant to arrive at the seeds. Thus, understanding the route of these minerals, and the genes responsible for their translocation may lead to strategies to increase flux to seeds and result in greater seed Fe, Zn, and Cu density.




Figure 1. Model of location of Fe, Zn, and Cu uptake or transport genes discussed in this review. (A) Model of wheat plant showing the following translocation steps to the seed: 1, uptake from the rhizosphere; 2, xylem loading; 3, root-to-shoot transfer; 4, distribution to the leaves or seed-covering tissues; 5, phloem loading for movement to seed; 6, loading into the seed. (B) Detail of seed loading. Gene families potentially involved in seed mineral micronutrient transport are pictured in hypothetical or known localizations. Maternal tissues are shown in green, filial tissues in gold. Efflux transporters are shown in light blue, plasma-membrane localized uptake transporters in dark blue, vacuolar uptake transporters in gray, vacuolar efflux transporters in red. Pd: plasmodesmata.

3. Defining the pathway of mineral micronutrients through the plant body

3.1. Uptake from rhizosphere

Uptake of micronutrients from the rhizosphere is the first step for accumulating micronutrients into the plant prior to translocation to seeds. Several recent reviews [30–32] have provided excellent coverage of the literature on Fe uptake processes and regulation of Fe uptake genes, such as *FIT/FER* transcription factors, *FRO* ferric-chelate reductase genes, and *IRT* ferrous-iron uptake transporters in dicots, and *IREG* and *IRO* transcription factors, phyto-

siderophore synthesis genes, and *YS/YSL* uptake genes in grasses [33–57] (table 2). Primary uptake of Zn is less well defined, but in *Arabidopsis* is likely carried out by Zn transporters of the *ZIP* family, some of which are Zn regulated in roots and other tissues [58]. Two transcription factors of the *bZIP* family are involved in up-regulation of several *ZIP* genes in response to Zn deficiency [59]. Uptake of Cu from the rhizosphere is also not well defined, and may be carried out by transporters of the *COPT* family [60,61]. Toxic heavy metals are generally transported in plants by transporters of essential micronutrients. Cadmium and zinc have similar chemical properties and are taken up and translocated within

Table 1. Quantitative isotope labeling of seeds of crop plants

Label	Crop species	Application method	Stage of application	% of label in seed	Reference
A. Radioisotopes					
⁵⁹ FeCl ₃	Wheat	Hydroponic	Prior to anthesis	8	[5]
⁶⁵ ZnCl ₂	Maize	Hydroponic	Prior to anthesis	27.6	[4]
⁶⁵ ZnCl ₂	Soybean	Hydroponic	5 weeks to maturity	21.3	[6]
⁶⁵ ZnCl ₂	Soybean	Hydroponic	Anthesis to maturity	27.6	[6]
⁶⁵ ZnCl ₂	Soybean	Hydroponic	Germination to maturity	21.6	[7]
⁶⁵ ZnCl ₂	Soybean	Hydroponic	4 weeks prior to anthesis	28.3	[7]
⁶⁵ ZnCl ₂	Soybean	Hydroponic	Anthesis to maturity	16.9	[7]
⁶⁵ ZnCl ₂	Soybean	Hydroponic	At anthesis for one week	23.4	[10]
⁶⁵ ZnCl ₂	Soybean	Foliar application	At anthesis	37.5	[10]
⁶⁵ ZnCl ₂	Soybean	Stem injection	At anthesis	64.5	[10]
⁶⁵ ZnCl ₂	Wheat	Hydroponic	At anthesis	2.3	[8]
⁶⁵ ZnCl ₂	Wheat	Foliar application	At anthesis	57.5	[8]
⁶⁵ ZnCl ₂	Wheat	Stem injection	At anthesis	62.6	[8]
Label	Crop species	Application method	Stage of application	Concentration in seed (μg g ⁻¹)	Reference
B. Stable isotope enrichment					
⁶⁵ CuCl ₂	Wheat	None (control)	Germination to maturity	8.1	[9]
⁶⁵ CuCl ₂	Wheat	Foliar application	At anthesis	12.9	[9]
⁶⁵ CuCl ₂	Wheat	Stem injection	At anthesis	22.1	[9]

Table 2. Genes implicated in uptake of Fe

Species	Gene	Description	References
A. Dicots			
<i>Solanaceae lycopersicum</i>	<i>FER</i>	Transcription factor – regulates Fe uptake genes	[37]
<i>Arabidopsis thaliana</i>	<i>FIT</i>	Transcription factor – regulates Fe uptake genes	[42,43]
<i>Arabidopsis thaliana</i>	<i>bHLH38</i>	Transcription factor – interacts with FIT	[51,54]
<i>Arabidopsis thaliana</i>	<i>bHLH39</i>	Transcription factor – interacts with FIT	[51,54]
<i>Arabidopsis thaliana</i>	<i>bHLH100</i>	Transcription factor – unknown targets	[51]
<i>Arabidopsis thaliana</i>	<i>bHLH101</i>	Transcription factor – unknown targets	[51]
<i>Arabidopsis thaliana</i>	<i>FRO2</i>	Ferric-chelate reductase	[33]
<i>Pisum sativum</i>	<i>FRO1</i>	Ferric-chelate reductase	[40]
<i>Cucumis sativus</i>	<i>FRO1</i>	Ferric-chelate reductase	[52]
<i>Solanaceae lycopersicum</i>	<i>FRO1</i>	Ferric-chelate reductase	[44]
<i>Arabidopsis thaliana</i>	<i>IRT1</i>	Iron transporter	[36,38,39]
<i>Pisum sativum</i>	<i>RIT1</i>	Iron transporter	[41]
<i>Cucumis sativus</i>	<i>IRT1</i>	Iron transporter	[52]
<i>Solanaceae lycopersicum</i>	<i>IRT1</i>	Iron transporter	[35]
<i>Arabidopsis thaliana</i>	<i>AHA2</i>	Proton ATPase	[57]
<i>Cucumis sativus</i>	<i>HA1</i>	Proton ATPase	[45]
B. Grass			
<i>Oryza sativa</i>	<i>IDEF1</i>	Transcription factor – regulates <i>IRO2</i>	[49]
<i>Oryza sativa</i>	<i>IDEF2</i>	Transcription factor – regulates <i>IRO2</i>	[53]
<i>Oryza sativa</i>	<i>IRO2</i>	Transcription factor – regulates Fe uptake genes	[48,50]
<i>Oryza sativa</i>	<i>IRT1</i>	Iron transporter	[46]
<i>Oryza sativa</i>	<i>YSL15</i>	Iron-phytosiderophore transporter	[55,56]
<i>Zea mays</i>	<i>YS1</i>	Iron-phytosiderophore transporter	[34]
<i>Hordeum vulgare</i>	<i>YS1</i>	Iron-phytosiderophore transporter	[47]

plants by similar pathways [62,63]. Therefore, tracing the movement of Cd could be helpful to understand the physiology and the transport of Zn. Several physiological studies with Cd and Zn in plants have demonstrated that Zn competitively inhibits Cd uptake in roots, suggesting a common transport mechanism [64–66]. It should be noted that many ZIP family proteins can transport several micronutrients, including Mn, Zn, Cu and Cd [35,41,46,52,58,59,67–87] (table 3). Since mutations in transporters can modify metal specificity [82], selecting for or engineering alleles that encode for Fe or Zn transporters that do not take up Cd presents a possible means to exclude Cd uptake into crop plants. However, some Cd can enter the xylem at the root tips where the Casparian strip is not yet formed or where it is disrupted by emerging lateral roots [88].

3.2. Xylem loading and transfer from root to shoot

Transfer of micronutrients from the root system into the shoot system has been considered a rate-limiting step for translocation to seeds [89]. Since Cd and Zn are loaded into the xylem by similar transport systems and the transfer of Cd from root to shoots (as opposed to Cd uptake at the roots) is the major physiological difference between high grain Cd and low grain Cd wheat and rice [90,91], this potentially offers another opportunity for allele selection or engineering to prevent Cd from entering the food supply. Several genes have been implicated in translocation of metal micronutrients into xylem or across the root-shoot junction (table 4), including *FRD3*, *FPN1*, *HMA2*, *HMA4*, *HMA5*, and *MTP3* [92–103]. *HMA* genes are members of the P_{1B} ATPase family [104], and some of these genes have been implicated in Zn and Cd xylem loading by pumping the ions from the pericycle to the xylem vessels [103]. Multiple copies of *HMA4* are found in the Zn hyperaccumulator *Arabidopsis halleri*, leading to higher *HMA4* expression and a mechanism for high Zn translocation rates to shoots [97]. Although much is to be learned, modification or overexpression of these genes offers potential for increasing movement of micronutrients into the shoot systems. Increased flux into shoots could possibly provide additional micronutrients for seed biofortification, and may activate native homeostasis mechanisms that increase root uptake capacity to keep pace.

Table 3. ZIP protein uptake specificities

Species	Protein	Uptake				Sensitivity	References
		Zn	Fe	Mn	Cu	Cd	
<i>Arabidopsis halleri</i>	IRT3	Y	Y	N ^a		N	[86]
<i>Arabidopsis thaliana</i>	IRT1	Y	Y	Y	N	Y	[68,70,82]
<i>Arabidopsis thaliana</i>	IRT2	Y	Y	N		N	[83]
<i>Arabidopsis thaliana</i>	IRT3	Y	Y	N		N	[86]
<i>Arabidopsis thaliana</i>	ZIP1	Y					[58]
<i>Arabidopsis thaliana</i>	ZIP2	Y			Y		[58,84]
<i>Arabidopsis thaliana</i>	ZIP3	Y					[58]
<i>Arabidopsis thaliana</i>	ZIP4	Y			Y		[59,84]
<i>Cucumis sativus</i>	IRT1		Y				[52]
<i>Glycine max</i>	ZIP1	Y	N	N	N	Y	[76]
<i>Hordeum vulgare</i>	IRT1	Y	Y	Y	Y		[78]
<i>Hordeum vulgare</i>	ZIP3	Y	N	N			[77]
<i>Hordeum vulgare</i>	ZIP5	Y	N	N			[77]
<i>Hordeum vulgare</i>	ZIP8	Y	N	N			[77]
<i>Solanaceae lycopersicum</i> ^b	IRT1	Y	Y	Y	Y		[35]
<i>Solanaceae lycopersicum</i>	IRT2	Y	Y	Y	Y		[35]
<i>Malus xiaojinensis</i>	IRT1		Y				[85]
<i>Medicago truncatula</i>	ZIP1	Y	N	N	N		[74]
<i>Medicago truncatula</i>	ZIP3	N	Y	N	N		[74]
<i>Medicago truncatula</i>	ZIP4	N	N	Y	N		[74]
<i>Medicago truncatula</i>	ZIP5	Y	Y	N	N		[74]
<i>Medicago truncatula</i>	ZIP6	Y	Y	N	N		[74]
<i>Medicago truncatula</i>	ZIP7	N	N	Y	N		[74]
<i>Oryza sativa</i>	IRT1		Y				[46,67]
<i>Oryza sativa</i>	IRT2		Y				[46]
<i>Oryza sativa</i>	ZIP1	Y				Y	[81]
<i>Oryza sativa</i>	ZIP3	Y					[81]
<i>Oryza sativa</i>	ZIP4	Y	Y				[69,72]
<i>Oryza sativa</i>	ZIP5	Y					[72]
<i>Oryza sativa</i>	ZIP7 ^a	Y					[87]
<i>Oryza sativa</i>	ZIP8	Y	Y				[73]
<i>Pisum sativum</i>	RIT1	Y	Y			Y	[41]
<i>Noccaea caerulescens</i> ^c	IRT1		Y			N	[80]
<i>Noccaea caerulescens</i>	Znt1	Y				Y	[71,79]
<i>Thlaspi japonicum</i>	Znt1	Y		Y		Y	[75]
<i>Thlaspi japonicum</i>	Znt2			Y		Y	[75]

a. N = tested with negative results b. Formerly *Lycopersicon esculentum*
c. Formerly *Thlaspi caerulescens*

Table 4. Genes implicated in the transfer of metal micronutrients (or Cd) from roots to shoots

Gene	Mineral	References
FRD3	Fe	[95,96]
FPN1	Fe	[100]
HMA2	Zn	[98]
HMA4	Zn, Cd	[94,97,98,101–103]
HMA5	Cu	[92,99]
MTP3	Zn	[93]

3.3. Translocation to leaf or seed covering tissues

Once in the shoot xylem, transpirational tension can carry micronutrients to leaves, where they may be taken into the symplast once again. Siliques have stomata [105] and carry out photosynthesis [106], and apparently transpire [107], as do glumes of wheat [108,109]. Thus, the potential for xylem transport of micronutrients directly to these seed covering tissues exists. Xylem unloading, or uptake from leaf or other xylem parenchyma apoplastic spaces into live cells, is not well characterized. For Fe, the process may be similar to primary uptake from the rhizosphere at the root surface, utilizing leaf expressed ferric chelate reductase [40,44,110,111] and Fe(II) transporters [52]. Several Zn regulated ZIP genes are expressed in leaves [84], as are COPT genes [60,61]. Studies in hyperaccumulator species have highlighted the involvement of ZIP genes in shoot Zn distribution. In *Thlaspi caerulescens*, the ZIP genes *Znt1* and *Znt5* are highly expressed compared to the nonaccumulator, *T. arvense* [79,112,113]. *T. caerulescens* accumulates Zn in epidermal storage cells of leaves to several-fold higher than in mesophyll cells [114]. Quantitative *in situ* hybridization indicated that *Znt5*, rather than *Znt1* was expressed in the Zn accumulating cells [115,116]. *MTP1* is a member of the cation diffusion facilitator (CDF) family, and is has been associated with tolerance to heavy metals in shoots. *Arabidopsis MTP1* plays a role in Zn homeostasis [117,118] by Zn sequestration in the vacuole. In *T. caerulescens* and *A. halleri*, *MTP1* is highly expressed in the shoot tissues [112,116,119].

In grasses, node localized transporters can act as a sort of molecular switching station, and facilitate the xylem-to-phloem transfer of nutrients to direct translocation preferentially to flag leaves or to the spike. The transporter *Lsi6* unloads silicon from xylem [120] and after panicle emergence directs silicon to the panicle [121]. Boron deficiency affected distribution of boron to the spike or flag leaf [122]. It is very possible that similar transporters for metal micronutrients exist in wheat and other species, and manipulation of these switches could direct more micronutrients toward spikes and the developing seeds, potentially improving both mineral concentrations in food and nutrient use efficiency of crops.

3.4. Phloem loading for translocation to seeds

Loading of micronutrients into the phloem is a requisite step for translocation to seeds. Even if minerals have arrived at the seed covering tissues, physical barriers between the maternal plant and developing seed and lack of transpirational tension will necessitate phloem transport to the seed tissues [126,127]. Experimental results support the idea that micronutrients pass through pods [18] or glumes [128] prior to transport to seeds. Movement of radioactive Zn through phloem from older to younger leaves, or from leaves to roots has been confirmed [129,130]. The pattern of transport and remobilization of Cd from vegetative tissues

to the seeds/grains is similar to Zn [29,131]. Cadmium introduced in the cut stem of wheat was shown to be removed from the transpiration stream, loaded into the phloem in the stem, and transported to the maturing grain similarly to zinc [131,132].

Mineral micronutrients are thought to be in chelated or ligand-bound forms when not incorporated into proteins [133,134], including during phloem transport. A ligand called ITP (iron transport protein) was identified from *Ricinus communis* phloem [135], but orthologs in other species have not been described. One potential phloem chelator is nicotianamine (NA). Chemical properties of NA have been reviewed recently [136], and they include the capacity to bind Cu, Co, Fe(II) and Fe(III), Mn, Ni, and Zn. NA is a precursor to phytosiderophores in grasses [137], but NA is present in dicots as well [138]. The *chln* mutant of tomato lacks the capacity to synthesize NA [139], and has disrupted micronutrient homeostasis and accumulates abnormal micronutrient concentrations in leaves [140–142]. Transgenic tobacco that lacks NA recapitulated this phenotype and had defective floral development [143]. Conversely, when NA synthase was overexpressed, seeds had higher than normal concentrations of Fe and Zn. In *Arabidopsis*, quadruple NA synthase mutants were constructed with two sets of alleles, resulting in a sterile *chln*-like phenotype, or a less severe phenotype that could reproduce, but had seeds with low Fe concentration, was chlorotic, and accumulated excess Fe in leaves at reproductive stage [144]. This body of evidence strongly suggests that NA is a vital chelator of micronutrients for homeostasis during growth, and for translocation within vegetative parts of the plant and also in phloem transport of micronutrients to seeds.

Proteins that transport micronutrient-NA complexes have been identified in recent years. These transporters are the yellow stripe-like (YSL) proteins, which are members of the oligopeptide transport (OPT) family. YellowStripe1 (YS1) is the primary Fe(III)- mugineic acid (phytosiderophore) uptake protein in maize [34] and is also involved in internal translocation of Fe [145,146]. In nongrasses, YSL proteins seem to function in interorgan translocation rather than for primary uptake of iron at the root surface. Expression of YSL genes typically is localized around vascular bundles, and thus may be involved with moving micronutrients to or from the vascular tissue. *Arabidopsis* YSLs 1, 2, and 3 have been shown by yeast Fe uptake mutant complementation to transport Fe(II)-NA [147,148]. A single *ysl1* mutant had increased NA in shoots, and lower NA and Fe concentrations in seeds [149]. A double mutant of *ysl1* and *ysl3* has an even more severe phenotype, with chlorosis resembling the *chln* mutant, low pollen viability, and low seed Fe, Zn, and Cu concentrations [150], implicating nicotianamine and YSL proteins in transport of multiple micronutrients. In a study of mineral partitioning throughout the *Arabidopsis* life cycle, rosette and cauline leaves of the *ysl1ysl3* mutant plants accumulated excess Zn and Cu, while stems had lower levels, indicating that xylem transport was normal, but phloem loading was inhibited [28]. In addition to low seed Fe, Zn, and Cu, silique hulls retained these minerals, suggesting that phloem loading in this seed covering tissue is part of the pathway to import micronutrients into seeds. Grafting of inflorescence stems of *ysl1ysl3* mutants onto wild type rosettes, or the reciprocal experiment, showed that functional YSL1 and YSL3 proteins are needed in tissues of the aerial parts (i.e., flowers, cauline leaves, and silique hulls) for normal floral development and seed loading of Cu, Fe, and Mn [147]. A rice RNAi line with lowered YSL2 expression had decreased Fe translocation to the shoots and seeds when compared to wild type, suggesting that OsYSL2 is involved in long distance transport of Fe to the seeds in rice [151].

Other members of the oligopeptide transporter family are important for transport of mineral micronutrients to seeds. The most characterized gene of the oligopeptide transporter family is *Arabidopsis* *OPT3*. Expression of *OPT3* was upregulated in roots and leaves of Fe-deficient *Arabidopsis* plants, and expression of this gene in yeast complemented growth defects of Cu and Fe uptake mutants [84]. Complete knockout mutants of *AtOPT3* were embryo lethal [152], but a T-DNA insertion in the promoter region resulted in an expression knockdown phenotype [153]. These mutants accumulated high Fe, Cu, Mn, and Zn concentrations in vegetative tissues, but had low Fe concentrations in seeds. Individual *OPT* family members are expressed differently from each other in various tissues [154,155], suggesting specialized functions for each gene in micronutrient transport within the plant. The substrates for plant OPTs are not well defined, but could be micronutrients chelated to NA [155], various tetra- and penta-peptides, and reduced glutathione [156].

3.5. Gradient to the seed and co-transport with N

Movement of phloem sap occurs by bulk flow, resulting from a pressure gradient from the source (leaf, stem, etc.) tissues to the sink (seed) tissues [126]. The pressure gradient is generated by loading nutrients such as carbohydrates and potassium into phloem at source tissues to provide osmotic potential that draws in water. The unloading of these nutrients at the seed decreases the pressure and maintains a gradient. Micronutrients will not be present in high enough concentrations to create a pressure gradient on their own, and thus will move with the major nutrients—that is, K^+ , Cl^- , and sugars [126,157] in the phloem. The alkaline phloem sap, at pH 7.2–8.2 [157], necessitates chelation of Fe, which decreases in solubility as pH rises [158]. As mentioned above, NA is a potential chelator of Fe in the phloem. NA can form a stable complex at alkaline pH [159], and can also chelate Cu and Zn. Other potential phloem chelators are unspecified oligopeptides and certain amino acids [134]. These compounds all contain N, suggesting that phloem transport of N and mineral micronutrients may be directly related. Additional evidence for this relationship between N and micronutrient transport to seeds comes from high correlations between N and Zn in seeds of several grass and dicot species [160]. Additionally, increasing N availability to wheat, while Zn availability remained constant, resulted in increased translocation to and concentration of Zn in wheat grain [12,161]. Thus, improved fertilization practices with N and Zn fertilizers could be a large step toward biofortification. Much of the seed N is supplied as amino acids from protein breakdown during senescence [17]. Wheat lines with delayed senescence had decreased amino acid export from leaves [162] and lower grain N, Fe, and Zn concentrations [163], while an RNAi line with delayed leaf senescence had lower remobilization of N, Fe, and Zn and lower grain concentrations of these nutrients [24], suggesting possible co-transport mechanisms.

3.6. Transport into seeds and storage forms

Seeds contain filial tissues (embryo and endosperm, aleurone) surrounded by maternal tissues (seed coat). The developing seed is connected to the maternal plant by a single vascular trace [126,164]. This vascular bundle ends at the seed coat and is not symplastically connected to the endosperm or embryo. Nutrients moving to the seed are unloaded from the phloem, while excess water is thought to move in the opposite direction in this vascular strand, through the xylem or suberized cells [165]. Phloem delivered nutrients are distributed in the maternal tissue surrounding the seed, and are eventually effluxed into an apoplastic space

that separates the maternal and filial tissues. Thus, specific micronutrient efflux transporters are likely to be required for this action, as well as for uptake into cells of the endosperm and/or embryo. The endosperm tissue is present in mature cereals, but is consumed by the embryo in legumes and *Arabidopsis* during seed maturation [164]. For mineral micronutrients, the transporters that are involved in phloem unloading and uptake by the filial tissues are not well defined, but likely belong to the families discussed as transporters in other parts of the plant. Indeed, laser capture microdissection of barley grain into transfer cells of the maternal vascular bundle, aleurone, endosperm, and embryo fractions, followed by microarray analysis, revealed expression of genes from the following families: HMA, ZIP, MTP, Nramp, NAS, and YSL [166]. Typically the expression was not evenly distributed among the different cell types, suggesting that, as in other tissues, certain transporters are important in specific cell types to move micronutrients through the plant.

Within the *Arabidopsis* seed, Fe is stored in vacuoles of the embryo endodermis [167]. X-ray fluorescence microtomography imaging of developing embryos revealed that Zn is not confined to the same cells, but is distributed throughout the embryo [168]. Fe is loaded into the vacuoles by the VIT1 protein during embryo development [168], while upon germination Fe is remobilized from this organelle by the Nramp3 and Nramp4 transporters [169]. In legumes, Fe is stored in ferritin protein in the embryo [170,171] but ferritin is not found in the seed coat [172]. Ferritin is degraded during seed germination [171] thus releasing stored Fe for the growing seedling. Ferritin is estimated to store 20–40% of legume seed Fe [173], and in addition to amyloplast localized ferritin, Fe is stored in epidermal cells and cells near the provascular tissue of the embryo [174]. In some varieties of soybean and common bean, a substantial percentage of total seed Fe is localized to the seed coat [175,176]. In grains, the highest concentrations of Fe and Zn co-localize with protein and free amino acids in the embryo and bran, with lower concentrations in endosperm [14,160,177]. This is problematic from a human nutrition perspective, since processing of rice, wheat, or other cereals in ways that remove these parts results in greatly reduced nutritional value. The high concentration of the phosphorus storage molecule phytate in cereal grains [178] is also problematic, as phytate can also bind Fe and Zn, and decrease absorption of these nutrients in the animal gut. Increasing wheat grain Zn levels by agronomic practices did not result in increased phytate [160], and lines of cereals with mutations that result in low phytate concentration did not consistently have altered levels of Fe or Zn in rice, maize, wheat, barley, or soybean seeds [179–183]. These results suggest that phytate can be decreased in future variety development while Fe and Zn concentrations remain at current levels or higher. Feeding trials have shown that low phytate maize had substantially increased Zn bioavailability [184,185]. Decreasing antinutrients such as phytate while simultaneously increasing micronutrients could have a synergistic effect on Fe and Zn bioavailability. Strategies for altering seed phytate levels have been reviewed recently [178].

4. Future discovery

Although some of the genes that are important for translocating Fe, Zn, and Cu to seeds have been identified, there are surely many additional genes to be discovered. Knowing the pathways taken by minerals through the plant as they are translocated to seeds will give important clues on where to focus efforts when looking for new genes or validating candidates. Major focus has been on *Arabidopsis* in recent years, as it has been the most resource rich plant species, in terms of genomic information and

mutants. Information for other plant species is quickly catching up as additional genomes are sequenced (e.g., crop species rice [186], sorghum [187], and soybean [188]) and scientific communities develop databases and technical resources. With second generation sequencing technologies, transcriptomic data can be generated from any species, even those with no previous sequence information [189,190]. Although *Arabidopsis* can be a useful species for studying many aspects of understanding mineral transport to seeds, other plants species with different characteristics may be better suited for other aspects. Knowledge may be transferred more quickly to biofortified crops if additional model species are adopted, or if the staple crop species themselves are studied directly.

4.1. Quantitative genetics

Potential methods for discovering new genes that affect translocation of micronutrients to seeds include quantitative trait locus (QTL) mapping. Two methods are commonly used for mapping QTL: linkage mapping and association mapping [191–194]. Linkage mapping requires construction of biparental populations consisting of F_2 , recombinant inbred lines (RILs), or backcross progeny. Association mapping is similar to linkage mapping, except that instead of using populations specifically created for QTL mapping, any sample of individuals can be used, such as existing elite lines from breeding programs or exotic accessions from germplasm collections. In many species, high-throughput genotyping is beginning to make genome-wide association mapping feasible. If genomic information allows, candidate genes for QTLs can be proposed and testing can be initiated. If there are no obvious candidate genes in the QTL region, or if genomic information is lacking, the QT genes can be pinpointed by fine mapping and positional cloning.

One strength of using QTL or association mapping is that no prior knowledge about the new genes is required for discovery. One case in point is the discovery of *NAM-B1*, a transcription factor in bread wheat that influences timing of leaf senescence, grain protein concentration, and grain Fe and Zn concentrations [24,163]. QTL mapping of seed mineral concentration has been carried out in *Arabidopsis* [195–197], *Medicago truncatula* [198], *Lotus japonicus* [199], common bean [200–202], *Brassica napus* [203], *Brassica rapa* [204], rice [205–207], and wheat [208,209]. These studies have identified several QTL for seed mineral concentrations localized in different regions and several of these mineral QTL co-localized with each other, pointing to common transport mechanisms. Several candidate genes for the different QTLs have been identified in many of these species allowing future fine mapping of these quantitative genes aimed at developing biofortified crops.

4.2. Candidate genes by expression pattern and predicted function

Combining knowledge of when and where micronutrients are translocated to seeds with gene expression data will allow identification of candidate genes. Development of genomic resources has allowed tools such as microarrays to become commonplace. These tools allow large-scale transcriptomic profiling for various experimental treatments (i.e., Fe or Zn deficient or replete), between mutants, ecotypes or species with different micronutrient accumulation (i.e., low seed micronutrient mutants) or during development (i.e., several time points over seed filling). For example, differential expression of genes in metal micronutrient (Zn, Ni) and non-nutrient (Cd) hyperaccumulator species compared to non-hyperaccumulator species may offer clues to function of genes in micronutrient translocation to

Table 5. Metal transporter genes with higher expression in hyperaccumulator species.

Non-accumulator species			Hyperaccumulator species		References
<i>Arabidopsis thaliana</i>	<i>Arabidopsis lyrata</i> ssp. <i>petraea</i>	<i>Thlaspi arvense</i>	<i>Arabidopsis halleri</i>	<i>Noccaeae caerulescens</i>	
IRT2				IRT2	[212]
IRT3		IRT3/Znt2	IRT3	IRT3/Znt2	[79,86,112,113,213,216,217]
ZIP1			ZIP1		[119,213]
ZIP3			ZIP3		[213,216]
ZIP4		Znt1	ZIP4	Znt1	[71,79,113,119,211–213]
ZIP5		Znt5	ZIP5	Znt5	[113,210,212]
ZIP6	ZIP6	ZIP6	ZIP6	Znt6	[113,119,210,211,213,214,216]
		ZIP7		ZIP7	[113]
ZIP9			ZIP9		[211,213,216]
ZIP10			ZIP10	ZIP10	[212,213]
ZIP12			ZIP12		[216]
HMA1			HMA1		[213]
HMA3		HMA3	HMA3	HMA3	[113,119,212,213,216,217]
HMA4		HMA4	HMA4	HMA4	[113,212,213,216]
HMA6			HMA6		[213]
Nramp3	Nramp3		Nramp3	Nramp3	[211–214,216]
Nramp5			Nramp5		[216]
NAS3					[119,211,213,217]
NAS2			NAS2	NAS2	[211–213]
NAS4			NAS4		[211,213]
		NAS1		NAS1	[113]
MTP1/ZAT/CDF1	MTP1		MTP1	MTP1	[119,212,213,215–217]
MTP2			MTP2		[216]
MTP3			MTP3		[216]
		MTP5		MTP5	[113]
MTP8			MTP8	MTP8	[212,213]
		MTP11		MTP11	[113,213,217]
		MTP12		MTP12	[113]
FRD3			FRD3	FRD3	[212,213]
FPN1/IREG			FPN1/IREG		[213]
YSL6			YSL6		[213]
YSL7				YSL7	[212]

a Formerly *Thlaspi caerulescens*.

seeds. Hyperaccumulator species have increased expression or duplication of genes involved in metal micronutrient transport across plasma membranes (ZIPs), in root-to-shoot translocation (*HMA*, *FRD3*, *FPN1*), xylem or phloem transfer and chelation (*YSL*, *NAS*) and in vacuole loading/unloading (*MTP*, *Nramp*) (table 5) [71,79,86,112,113,119,210–217]. Since micronutrients pass through leaves and seed covering tissues to reach seeds, genes with increased expression in these tissues during seed fill may be important for translocation to seeds. For example, 240 transporter genes exhibited increased expression during *Arabidopsis* leaf senescence [218], including some members of *YSL*, *ZIP*, *OPT*, and *HMA* families. Knowledge of gene family transport specificity, together with expression profiling to determine which specific family members are differentially expressed in certain stages of development or more highly in hyperaccumulators may allow researchers to focus on specific members of gene families for basic research or biofortification strategies.

5. Strategies for biofortification of mineral micronutrients

To this point, this review has focused on when and where, and by which genes, micronutrients are translocated to seeds. The goal for acquiring this knowledge is the application to crop development to improve human food or animal feed supplies for micronutrient density and bioavailability. The HarvestPlus program (www.harvestplus.org) has set goals to increase Fe concentration in common bean from 50 to 94 µg/g and pearl millet (*Pennisetum glaucum*) from 47 to 77 µg/g, and to increase Zn concentration in wheat from 25 to 33 µg/g, and rice Zn from 16 to 24 µg/g. Although traditional breeding or improved crop management has great potential to improve crop quality, this section will focus on transgenic strategies for biofortification of seeds.

For successful increase of seed micronutrient concentrations, genes to be expressed differently should be targeted to the appropriate localization, at the appropriate developmental stage, and the substrates for transporters (micronutrients themselves or co-transported chelators) should be present. Constitutive expression of transporters may not adequately direct when or where the increased expression of micronutrients takes place or have other unintended consequences. Constitutively overexpressing a *ZIP* transporter in barley resulted in higher short-term Zn uptake, and plants with higher concentrations of Fe and Zn [219], however the seeds were smaller and thus yield was likely reduced. Constitutively overexpressing the rice *ZIP4* gene resulted in aberrant distribution of Zn in the plant and lower seed Zn concentrations [220]. A targeted overexpression strategy may solve these types of problems.

For wheat, increasing Zn supply seems to be adequate to increase Zn grain concentrations [1,24], implying that grain sink strength or translocation to grains are not limiting factors, and that simply increasing Zn uptake may adequately biofortify seeds of some species. However, for rice, large increases in Zn supply and Zn accumulation in the plant body translated to much smaller increases in seed Zn [221]. Similar observations have been made for Fe supply in wild-type and Fe-overaccumulating mutant pea [11], and transgenic soybean that overexpressed root ferric reductase activity [222]. These results suggest that in some species, the rate limiting steps are downstream of uptake into the plant, and that a strategy to “push” more micronutrient into the plant may not be adequate by itself.

A strategy to “pull” micronutrients into the plant by increasing sink strength could be used. In this scenario, homeostatic mechanisms would allow the plant to adjust uptake to meet the additional demand. The main “pull” strategies that have been described utilize overexpression of the soybean Fe storage pro-

tein ferritin in rice grain. This strategy resulted in variable results between transgenic events and segregating lines, with no increase in some, or others with increases of up to threefold Fe concentration [223], including in the polished grain [224]. However, the increase in Fe concentration was not proportional to increases in ferritin protein levels [225], suggesting that an upstream step in micronutrient translocation to seeds was rate limiting. Transgenic maize with soybean ferritin expressed on an endosperm specific promoter had grain Fe increases of 20–70% over the control [226].

Another strategy would be to increase the translocation capacity of micronutrients to seeds by increasing transporters or substrates of transporters. Transgenic tobacco that constitutively overexpressed *NAS* had higher seed micronutrient concentrations [143,227], and rice with higher expression of *NAS3* accumulated higher Fe, Zn, and Cu [228]. Rice *YSL2* overexpression, driven by the constitutive 35S promoter, led to lower rather than higher seed and shoot Fe concentrations, however, overexpression driven by the phloem-localized sucrose transporter *SUT1* promoter resulted in a fourfold increase in seed Fe concentration [151].

The most successful transgenic biofortification of rice Fe concentration to date used a combination of increasing translocation capacity and seed sink strength. The ferritin gene from common bean was expressed on the endosperm-specific globulin promoter, while the *Arabidopsis NAS1* gene was constitutively expressed on the 35S promoter [229]. This resulted in a five- to six-fold increase in grain Fe concentration in polished rice grains.

6. Conclusion

Seed mineral improvement or biofortification will likely require simultaneous enhancement of several physiological processes, such as uptake from the rhizosphere, translocation from roots to shoots, phloem loading, and remobilization. Although progress has been made in terms of understanding micronutrient uptake and translocation and the genes involved, there remains much to be learned. Information about phloem delivery of micronutrients to seeds in different crops is particularly lacking. Multiple genes at different steps of micronutrient movement through the plant may need to be targeted simultaneously for the development of biofortified crops. Driving expression of this set of genes by common developmentally inducible promoters may be a potential strategy, as may tissue and development specific promoters to increase uptake and translocation capacity at the same time. Perhaps QTL mapping of seed mineral concentrations will reveal a “master regulator” that transmits seed micronutrient demand to the maternal plant and homeostatically adjusts uptake and translocation processes. Ideally, manipulation of such a regulator could accomplish all of these steps at once.

Acknowledgments – The authors thank Aaron Lorenz for helpful discussions regarding quantitative genetics. The authors also thank the reviewers for helpful suggestions.

References

- [1] I. Cakmak, Enrichment of cereal grains with zinc: agronomic or genetic biofortification? *Plant Soil* 302 (2008) 1–17.
- [2] P.J. White, M.R. Broadley, Biofortification of crops with seven mineral elements often lacking in human diets: iron, zinc, copper, calcium, magnesium, selenium and iodine, *New Phytol.* 182 (2009) 49–84.
- [3] W.H. Pfeiffer, B. McClafferty, HarvestPlus: breeding crops for better nutrition, *Crop Sci.* 47 (2007) S88–S105.
- [4] W. Garcia, R. Hodgson, C. Blessin, G. Inglett, Preparation of corn

- products endogenously labeled with zinc-65 for use in bioavailability studies, *J. Agric. Food Chem.* 25 (1977) 1290–1295.
- [5] A. Knight, Long term culture methods for the production of isotopically labelled plant material, *New Phytol.* 79 (1977) 573–582.
 - [6] M. Janghorbani, C. Weaver, B. Ting, V. Young, Labeling of soybeans with the stable isotope ^{70}Zn for use in human metabolic studies, *J. Nutr.* 113 (1983) 973–978.
 - [7] H. Schmitt, C. Weaver, Level of application and period of exposure affecting accumulation and distribution of chromium-51 and zinc-65 in hydroponically grown kale, bush beans, and soybeans, *J. Agric. Food Chem.* 32 (1984) 498–503.
 - [8] T.L. Starks, P.E. Johnson, Techniques for intrinsically labeling wheat with Zn-65, *J. Agric. Food Chem.* 33 (1985) 691–698.
 - [9] T.L. Starks, P.E. Johnson, Evaluation of foliar application and stem injection as techniques for intrinsically labeling wheat with copper-65, *J. Agric. Food Chem.* 34 (1986) 23–26.
 - [10] A. Khan, C. Weaver, Pattern of zinc-65 incorporation into soybean seeds by root absorption, stem injection, and foliar application, *J. Agric. Food Chem.* 37 (1989) 855–860.
 - [11] M.A. Grusak, Iron transport to developing ovules of *Pisum sativum*. I. Seed import characteristics and phloem iron-loading capacity of source regions, *Plant Physiol.* 104 (1994) 649–655.
 - [12] E.B. Erenoglu, U.B. Kutman, Y. Ceylan, B. Yildiz, I. Cakmak, Improved nitrogen nutrition enhances root uptake, root-to-shoot translocation and remobilization of zinc (^{65}Zn) in wheat, *New Phytol.* (2010), doi:10.1111/j.1469-8137.2010.03488.x.
 - [13] C.Y. Wu, L.L. Lu, X.E. Yang, Y. Feng, Y.Y. Wei, H.L. Hao, P.J. Stoffella, Z.L. He, Uptake, translocation, and remobilization of zinc absorbed at different growth stages by rice genotypes of different Zn densities, *J. Agric. Food Chem.* 58 (2010) 6767–6773.
 - [14] I. Cakmak, M. Kalayci, Y. Kaya, A.A. Torun, N. Aydin, Y. Wang, Z. Arisoy, H. Erdem, A. Yazici, O. Gokmen, L. Ozturk, W.J. Horst, Biofortification and localization of zinc in wheat grain, *J. Agric. Food Chem.* 58 (2010) 9092–9102.
 - [15] A.J. Barneix, Physiology and biochemistry of source-regulated protein accumulation in the wheat grain, *J. Plant Physiol.* 164 (2007) 581–590.
 - [16] N.M. Heidlebaugh, B.R. Trethewey, A.K. Jukanti, D.L. Parrott, J.M. Martin, A.M. Fischer, Effects of a barley (*Hordeum vulgare*) chromosome 6 grain protein content locus on whole-plant nitrogen reallocation under two different fertilisation regimes, *Funct. Plant Biol.* 35 (2008) 619–632.
 - [17] C. Masclaux-Daubresse, M. Reisdorf-Cren, M. Orsel, Leaf nitrogen remobilisation for plant development and grain filling, *Plant Biol.* 10 (2008) 23–36.
 - [18] P.J. Hocking, J.S. Pate, Mobilization of minerals to developing seeds of legumes, *Ann. Bot.* 41 (1977) 1259–1278.
 - [19] J. Pate, P. Hocking, Phloem and xylem transport in the supply of minerals to a developing legume (*Lupinus albus* L.) fruit, *Ann. Bot.* 42 (1978) 911–921.
 - [20] J. Hill, The remobilization of nutrients from leaves, *J. Plant Nutr.* 2 (1980) 407–444.
 - [21] P.J. Hocking, Dry-matter production, mineral nutrient concentrations, and nutrient distribution and redistribution in irrigated spring wheat, *J. Plant Nutr.* 17 (1994) 1289–1308.
 - [22] R.O. Miller, J.S. Jacobsen, E.O. Skogley, Aerial accumulation and partitioning of nutrients by hard red spring wheat, *Commun. Soil Sci. Plant Anal.* 25 (1994) 1891–1911.
 - [23] T.P. Garnett, R.D. Graham, Distribution and remobilization of iron and copper in wheat, *Ann. Bot.* 95 (2005) 817–826.
 - [24] B.M. Waters, C. Uauy, J. Dubcovsky, M.A. Grusak, Wheat (*Triticum aestivum*) NAM proteins regulate the translocation of iron, zinc, and nitrogen compounds from vegetative tissues to grain, *J. Exp. Bot.* 60 (2009) 4263–4274.
 - [25] W. Jiang, P.C. Struik, J. Lingna, H. van Keulen, Z. Ming, T.J. Stomph, Uptake and distribution of root-applied or foliar-applied Zn-65 after flowering in aerobic rice, *Ann. Appl. Biol.* 150 (2007) 383–391.
 - [26] T.J. Stomph, W. Jiang, P.C. Struik, Zinc biofortification of cereals: rice differs from wheat and barley, *Trends Plant Sci.* 14 (2009) 123–124.
 - [27] E. Himelblau, R.M. Amasino, Nutrients mobilized from leaves of *Arabidopsis thaliana* during leaf senescence, *J. Plant Physiol.* 158 (2001) 1317–1323.
 - [28] B.M. Waters, M.A. Grusak, Whole-plant mineral partitioning throughout the life cycle in *Arabidopsis thaliana* ecotypes Columbia, Landsberg erecta, Cape Verde Islands, and the mutant line *ysl3*, *New Phytol.* 177 (2008) 389–405.
 - [29] R.P. Sankaran, S.D. Ebbs, Transport of Cd and Zn to seeds of Indian mustard (*Brassica juncea*) during specific stages of plant growth and development, *Physiol. Plant* 132 (2008) 69–78.
 - [30] R.F.H. Giehl, A.R. Meda, N. von Wiren, Moving up, down, and everywhere: signaling of micronutrients in plants, *Curr. Opin. Plant Biol.* 12 (2009) 320–327.
 - [31] J. Morrissey, M.L. Guerinot, Iron uptake and transport in plants: the good, the bad, and the ionome, *Chem. Rev.* 109 (2009) 4553–4567.
 - [32] E.L. Walker, E.L. Connolly, Time to pump iron: iron-deficiency-signaling mechanisms of higher plants, *Curr. Opin. Plant Biol.* 11 (2008) 530–535.
 - [33] N.J. Robinson, C.M. Procter, E.L. Connolly, M.L. Guerinot, A ferric-chelate reductase for iron uptake from soils, *Nature* 397 (1999) 694–697.
 - [34] C. Curie, Z. Panaviene, C. Loulergue, S.L. Dellaporta, J.F. Briat, E.L. Walker, Maize *yellow stripe1* encodes a membrane protein directly involved in Fe(III) uptake, *Nature* 409 (2001) 346–349.
 - [35] U. Eckhardt, A.M. Marques, T.J. Buckhout, Two iron-regulated cation transporters from tomato complement metal uptake-deficient yeast mutants, *Plant Mol. Biol.* 45 (2001) 437–448.
 - [36] R. Henriques, J. Jasik, M. Klein, E. Martinoia, U. Feller, J. Schell, M.S. Pais, C. Koncz, Knock-out of *Arabidopsis* metal transporter gene *IRT1* results in iron deficiency accompanied by cell differentiation defects, *Plant Mol. Biol.* 50 (2002) 587–597.
 - [37] H.Q. Ling, P. Bauer, Z. Berczky, B. Keller, M. Ganal, The tomato *fer* gene encoding a bHLH protein controls iron-uptake responses in roots, *Proc. Natl. Acad. Sci. U.S.A.* 99 (2002) 13938–13943.
 - [38] C. Varotto, D. Maiwald, P. Pesaresi, P. Jahns, F. Salamini, D. Leister, The metal ion transporter *IRT1* is necessary for iron homeostasis and efficient photosynthesis in *Arabidopsis thaliana*, *Plant J.* 31 (2002) 589–599.
 - [39] G. Vert, N. Grotz, F. Dedaldechamp, F. Gaymard, M.L. Guerinot, J.F. Briat, C. Curie, *IRT1*, an *Arabidopsis* transporter essential for iron uptake from the soil and for plant growth, *Plant Cell* 14 (2002) 1223–1233.
 - [40] B.M. Waters, D.G. Blevins, D.J. Eide, Characterization of *FRO1*, a pea ferric-chelate reductase involved in root iron acquisition, *Plant Physiol.* 129 (2002) 85–94.
 - [41] C.K. Cohen, D.F. Garvin, L.V. Kochian, Kinetic properties of a micronutrient transporter from *Pisum sativum* indicate a primary function in Fe uptake from the soil, *Planta* 218 (2004) 784–792.
 - [42] E.P. Colangelo, M.L. Guerinot, The essential basic helix-loop-helix protein *FIT1* is required for the iron deficiency response, *Plant Cell* 16 (2004) 3400–3412.
 - [43] M. Jakoby, H.Y. Wang, W. Reidt, B. Weisshaar, P. Bauer, *FRU1* (*BHLH029*) is required for induction of iron mobilization genes in *Arabidopsis thaliana*, *FEBS Lett.* 577 (2004) 528–534.
 - [44] L.H. Li, X.D. Cheng, H.Q. Ling, Isolation and characterization of Fe(III)-chelate reductase gene *LeFRO1* in tomato, *Plant Mol. Biol.* 54 (2004) 125–136.
 - [45] S. Santi, S. Cesco, Z. Varanini, R. Pinton, Two plasma membrane H^{+} -ATPase genes are differentially expressed in iron-deficient cucumber plants, *Plant Physiol. Biochem.* 43 (2005) 287–292.
 - [46] Y. Ishimaru, M. Suzuki, T. Tsukamoto, K. Suzuki, M. Nakazono, T. Kobayashi, Y. Wada, S. Watanabe, S. Matsushashi, M. Takahashi, H. Nakanishi, S. Mori, N.K. Nishizawa, Rice plants take up iron as an Fe^{3+} -phytosiderophore and as Fe^{2+} , *Plant J.* 45 (2006) 335–346.
 - [47] Y. Murata, J.F. Ma, N. Yamaji, D. Ueno, K. Nomoto, T. Iwashita, A specific transporter for iron(III)-phytosiderophore in barley roots, *Plant J.* 46 (2006) 563–572.
 - [48] Y. Ogo, R.N. Itai, H. Inoue, T. Kobayashi, M. Suzuki, M. Takahashi, S. Mori, N.K. Nishizawa, N.K. Nishizawa, Isolation and characterization of *IRO2*, a novel iron-regulated bHLH transcription factor in graminaceous plants, *J. Exp. Bot.* 57 (2006) 2867–2878.

- [49] T. Kobayashi, Y. Ogo, R.N. Itai, H. Nakanishi, M. Takahashi, S. Mori, N.K. Nishizawa, The transcription factor IDEF1 regulates the response to and tolerance of iron deficiency in plants, *Proc. Natl. Acad. Sci. U.S.A.* 104 (2007) 19150–19155.
- [50] Y. Ogo, R.N. Itai, H. Nakanishi, T. Kobayashi, M. Takahashi, S. Mori, N.K. Nishizawa, The rice bHLH protein OsIRO2 is an essential regulator of the genes involved in Fe uptake under Fe-deficient conditions, *Plant J.* 51 (2007) 366–377.
- [51] H.Y. Wang, M. Klatte, M. Jakoby, H. Baumlein, B. Weisshaar, P. Bauer, Iron deficiency-mediated stress regulation of four subgroup Ib BHLH genes in *Arabidopsis thaliana*, *Planta* 226 (2007) 897–908.
- [52] B.M. Waters, C. Lucena, F.J. Romera, G.G. Jester, A.N. Wynn, C.L. Rojas, E. Alcantara, R. Perez-Vicente, Ethylene involvement in the regulation of the H⁺-ATPase CsHA1 gene and of the new isolated ferric reductase CsFRO1 and iron transporter CsIRT1 genes in cucumber plants, *Plant Physiol. Biochem.* 45 (2007) 293–301.
- [53] Y. Ogo, T. Kobayashi, R.N. Itai, H. Nakanishi, Y. Kakei, M. Takahashi, S. Toki, S. Mori, N.K. Nishizawa, A novel NAC transcription factor, IDEF2, that recognizes the iron deficiency-responsive element 2 regulates the genes involved in iron homeostasis in plants, *J. Biol. Chem.* 283 (2008) 13407–13417.
- [54] Y.X. Yuan, H.L. Wu, N. Wang, J. Li, W.N. Zhao, J. Du, D.W. Wang, H.Q. Ling, FIT interacts with AtbHLH38 and AtbHLH39 in regulating iron uptake gene expression for iron homeostasis in *Arabidopsis*, *Cell Res.* 18 (2008) 385–397.
- [55] H. Inoue, T. Kobayashi, T. Nozoye, M. Takahashi, Y. Kakei, K. Suzuki, M. Nakazono, H. Nakanishi, S. Mori, N.K. Nishizawa, Rice OsYSL15 is an iron-regulated iron(III)-deoxymugineic acid transporter expressed in the roots and is essential for iron uptake in early growth of the seedlings, *J. Biol. Chem.* 284 (2009) 3470–3479.
- [56] S. Lee, J.C. Chiecko, S.A. Kim, E.L. Walker, Y. Lee, M.L. Guerinot, G. An, Disruption of OsYSL15 leads to iron inefficiency in rice plants, *Plant Physiol.* 150 (2009) 786–800.
- [57] S. Santi, W. Schmidt, Dissecting iron deficiency-induced proton extrusion in *Arabidopsis* roots, *New Phytol.* 183 (2009) 1072–1084.
- [58] N. Grotz, T. Fox, E. Connolly, W. Park, M.L. Guerinot, D. Eide, Identification of a family of zinc transporter genes from *Arabidopsis* that respond to zinc deficiency, *Proc. Natl. Acad. Sci. U.S.A.* 95 (1998) 7220–7224.
- [59] A.G.L. Assunção, E. Herrero, Y.F. Lin, B. Huettel, S. Talukdar, C. Smaczniak, R.G.H. Immink, M. van Eldik, M. Fiers, H. Schat, M.G.M. Aarts, *Arabidopsis thaliana* transcription factors bZIP19 and bZIP23 regulate the adaptation to zinc deficiency, *Proc. Natl. Acad. Sci. U.S.A.* 107 (2010) 10296–10301.
- [60] T. del Pozo, V. Cambiazio, M. González, Gene expression profiling analysis of copper homeostasis in *Arabidopsis thaliana*, *Biochem. Biophys. Res. Commun.* 393 (2010) 248–252.
- [61] V. Sancenon, S. Puig, H. Mira, D.J. Thiele, L. Penarrubia, Identification of a copper transporter family in *Arabidopsis thaliana*, *Plant Mol. Biol.* 51 (2003) 577–587.
- [62] C.A. Grant, W.T. Buckley, L.D. Bailey, F. Selles, Cadmium accumulation in crops, *Can. J. Plant Sci.* 78 (1998) 1–17.
- [63] N.S. Harris, G.J. Taylor, Remobilization of cadmium in maturing shoots of near isogenic lines of durum wheat that differ in grain cadmium accumulation, *J. Exp. Bot.* 52 (2001) 1473–1481.
- [64] E. Lombi, K.L. Tearall, J.R. Howarth, F.J. Zhao, M.J. Hawkesford, S.P. McGrath, Influence of iron status on cadmium and zinc uptake by different ecotypes of the hyperaccumulator *Thlaspi caerulescens*, *Plant Physiol.* 128 (2002) 1359–1367.
- [65] D.E. Salt, I.J. Pickering, R.C. Prince, D. Gleba, S. Dushenkov, R.D. Smith, I. Raskin, Metal accumulation by aquacultured seedlings of Indian mustard, *Environ. Sci. Technol.* 31 (1997) 1636–1644.
- [66] S. Thomine, R.C. Wang, J.M. Ward, N.M. Crawford, J.I. Schroeder, Cadmium and iron transport by members of a plant metal transporter family in *Arabidopsis* with homology to Nramp genes, *Proc. Natl. Acad. Sci. U.S.A.* 97 (2000) 4991–4996.
- [67] N. Bughio, H. Yamaguchi, N.K. Nishizawa, H. Nakanishi, S. Mori, Cloning an iron-regulated metal transporter from rice, *J. Exp. Bot.* 53 (2002) 1677–1682.
- [68] D. Eide, M. Broderius, J. Fett, M.L. Guerinot, A novel iron-regulated metal transporter from plants identified by functional expression in yeast, *Proc. Natl. Acad. Sci. U.S.A.* 93 (1996) 5624–5628.
- [69] Y. Ishimaru, M. Suzuki, T. Kobayashi, M. Takahashi, H. Nakanishi, S. Mori, N.K. Nishizawa, OsZIP4, a novel zinc-regulated zinc transporter in rice, *J. Exp. Bot.* 56 (2005) 3207–3214.
- [70] Y.O. Korshunova, D. Eide, W.G. Clark, M.L. Guerinot, H.B. Pakrasi, The IRT1 protein from *Arabidopsis thaliana* is a metal transporter with a broad substrate range, *Plant Mol. Biol.* 40 (1999) 37–44.
- [71] M.M. Lasat, N.S. Pence, D.F. Garvin, S.D. Ebbs, L.V. Kochian, Molecular physiology of zinc transport in the Zn hyperaccumulator *Thlaspi caerulescens*, *J. Exp. Bot.* 51 (2000) 71–79.
- [72] S. Lee, H.J. Jeong, S.A. Kim, J. Lee, M.L. Guerinot, G. An, OsZIP5 is a plasma membrane zinc transporter in rice, *Plant Mol. Biol.* 73 (2010) 507–517.
- [73] S. Lee, S.A. Kim, J. Lee, M.L. Guerinot, G. An, Zinc deficiency-inducible OsZIP8 encodes a plasma membrane-localized zinc transporter in rice, *Mol. Cells* 29 (2010) 551–558.
- [74] A.F. Lopez-Millan, D.R. Ellis, M.A. Grusak, Identification and characterization of several new members of the ZIP family of metal ion transporters in *Medicago truncatula*, *Plant Mol. Biol.* 54 (2004) 583–596.
- [75] T. Mizuno, K. Usui, K. Horie, S. Nosaka, N. Mizuno, H. Obata, Cloning of three ZIP/Nramp transporter genes from a Ni hyperaccumulator plant *Thlaspi japonicum* and their Ni²⁺-transport abilities, *Plant Physiol. Biochem.* 43 (2005) 793–801.
- [76] S. Moreau, R.M. Thomson, B.N. Kaiser, B. Trevaskis, M.L. Guerinot, M.K. Udvardi, A. Puppo, D.A. Day, GmZIP1 encodes a symbiosis-specific zinc transporter in soybean, *J. Biol. Chem.* 277 (2002) 4738–4746.
- [77] P. Pedas, J.K. Schjoerring, S. Husted, Identification and characterization of zinc starvation-induced ZIP transporters from barley roots, *Plant Physiol. Biochem.* 47 (2009) 377–383.
- [78] P. Pedas, C.K. Ytting, A.T. Fuglsang, T.P. Jahn, J.K. Schjoerring, S. Husted, Manganese efficiency in barley: identification and characterization of the metal ion transporter HvIRT1, *Plant Physiol.* 148 (2008) 455–466.
- [79] N.S. Pence, P.B. Larsen, S.D. Ebbs, D.L. Letham, M.M. Lasat, D.F. Garvin, D. Eide, L.V. Kochian, The molecular physiology of heavy metal transport in the Zn/Cd hyperaccumulator *Thlaspi caerulescens*, *Proc. Natl. Acad. Sci. U.S.A.* 97 (2000) 4956–4960.
- [80] S. Plaza, K.L. Tearall, F.J. Zhao, P. Buchner, S.P. McGrath, M.J. Hawkesford, Expression and functional analysis of metal transporter genes in two contrasting ecotypes of the hyperaccumulator *Thlaspi caerulescens*, *J. Exp. Bot.* 58 (2007) 1717–1728.
- [81] S.A. Ramesh, R. Shin, D.J. Eide, D.P. Schachtman, Differential metal selectivity and gene expression of two zinc transporters from rice, *Plant Physiol.* 133 (2003) 126–134.
- [82] E.E. Rogers, D.J. Eide, M.L. Guerinot, Altered selectivity in an *Arabidopsis* metal transporter, *Proc. Natl. Acad. Sci. U.S.A.* 97 (2000) 12356–12360.
- [83] G. Vert, J.F. Briat, C. Curie, *Arabidopsis* IRT2 gene encodes a root-periphery iron transporter, *Plant J.* 26 (2001) 181–189.
- [84] H. Wintz, T. Fox, Y.Y. Wu, V. Feng, W.Q. Chen, H.S. Chang, T. Zhu, C. Vulpe, Expression profiles of *Arabidopsis thaliana* in mineral deficiencies reveal novel transporters involved in metal homeostasis, *J. Biol. Chem.* 278 (2003) 47644–47653.
- [85] P. Li, J.L. Qi, L. Wang, Q.N. Huang, Z.H. Han, L.P. Yin, Functional expression of MxIRT1, from *Malus xiaojinensis*, complements an iron uptake deficient mutant for plasma membrane targeting via membrane vesicles trafficking process, *Plant Sci.* 171 (2006) 52–59.
- [86] Y.F. Lin, H.M. Liang, S.Y. Yang, A. Boch, S. Clemens, C.C. Chen, J.F. Wu, J.L. Huang, K.C. Yeh, *Arabidopsis* IRT3 is a zinc-regulated and plasma membrane localized zinc/iron transporter, *New Phytol.* 182 (2009) 392–404.
- [87] X. Yang, J. Huang, Y. Jiang, H.S. Zhang, Cloning and functional identification of two members of the ZIP (Zrt, Irt-like protein) gene family in rice (*Oryza sativa* L.), *Mol. Biol. Rep.* 36 (2009) 281–287.

- [88] A. Lux, M. Martinka, M. Vaculík, P.J. White, Root responses to cadmium in the rhizosphere: a review, *J. Exp. Bot.* (2010).
- [89] M.G. Palmgren, S. Clemens, L.E. Williams, U. Kraemer, S. Borg, J.K. Schjorring, D. Sanders, Zinc biofortification of cereals: problems and solutions, *Trends Plant Sci.* 13 (2008) 464–473.
- [90] J.J. Hart, R.M. Welch, W.A. Norvell, L.V. Kochian, Characterization of cadmium uptake, translocation and storage in near-isogenic lines of durum wheat that differ in grain cadmium concentration, *New Phytol.* 172 (2006) 261–271.
- [91] S. Uruguchi, S. Mori, M. Kuramata, A. Kawasaki, T. Arai, S. Ishikawa, Root-to-shoot Cd translocation via the xylem is the major process determining shoot and grain cadmium accumulation in rice, *J. Exp. Bot.* 60 (2009) 2677–2688.
- [92] N. Andres-Colas, V. Sancenon, S. Rodriguez-Navarro, S. Mayo, D.J. Thiele, J.R. Ecker, S. Puig, L. Penarrubia, The *Arabidopsis* heavy metal P-type ATPase HMA5 interacts with metallochaperones and functions in copper detoxification of roots, *Plant J.* 45 (2006) 225–236.
- [93] S. Arrivault, T. Senger, U. Krämer, The *Arabidopsis* metal tolerance protein AtMTP3 maintains metal homeostasis by mediating Zn exclusion from the shoot under Fe deficiency and Zn oversupply, *Plant J.* 46 (2006) 861–879.
- [94] M. Courbot, G. Willems, P. Motte, S. Arvidsson, N. Roosens, P. Saumitou-Laprade, N. Verbruggen, A major quantitative trait locus for cadmium tolerance in *Arabidopsis halleri* colocalizes with HMA4, a gene encoding a heavy metal ATPase, *Plant Physiol.* 144 (2007) 1052–1065.
- [95] T.P. Durrett, W. Gassmann, E.E. Rogers, The FRD3-mediated efflux of citrate into the root vasculature is necessary for efficient iron translocation, *Plant Physiol.* 144 (2007) 197–205.
- [96] L.S. Green, E.E. Rogers, FRD3 controls iron localization in *Arabidopsis*, *Plant Physiol.* 136 (2004) 2523–2531.
- [97] M. Hanikenne, I.N. Talke, M.J. Haydon, C. Lanz, A. Nolte, P. Motte, J. Kroymann, D. Weigel, U. Krämer, Evolution of metal hyperaccumulation required *cis* regulatory changes and triplication of HMA4, *Nature* 453 (2008), 391–U344.
- [98] D. Hussain, M.J. Haydon, Y. Wang, E. Wong, S.M. Sherson, J. Young, J. Camakaris, J.F. Harper, C.S. Cobbett, P-type ATPase heavy metal transporters with roles in essential zinc homeostasis in *Arabidopsis*, *Plant Cell* 16 (2004) 1327–1339.
- [99] Y. Kobayashi, K. Kuroda, K. Kimura, J.L. Southon-Francis, A. Furuzawa, K. Kimura, S. Iuchi, M. Kobayashi, G.J. Taylor, H. Koyama, Amino acid polymorphisms in strictly conserved domains of a P-type ATPase HMA5 are involved in the mechanism of copper tolerance variation in *Arabidopsis*, *Plant Physiol.* 148 (2008) 969–980.
- [100] J. Morrissey, I.R. Baxter, J. Lee, L.T. Li, B. Lahner, N. Grotz, J. Kaplan, D.E. Salt, M.L. Gueriot, The ferroportin metal efflux proteins function in iron and cobalt homeostasis in *Arabidopsis*, *Plant Cell* 21 (2009) 3326–3338.
- [101] A. Papayan, L.V. Kochian, Identification of *Thlaspi caerulescens* genes that may be involved in heavy metal hyperaccumulation and tolerance. Characterization of a novel heavy metal transporting ATPase, *Plant Physiol.* 136 (2004) 3814–3823.
- [102] F. Verret, A. Gravot, P. Auroy, N. Leonhardt, P. David, L. Nussaume, A. Vavasseur, P. Richaud, Overexpression of AtHMA4 enhances root-to-shoot translocation of zinc and cadmium and plant metal tolerance, *FEBS Lett.* 576 (2004) 306–312.
- [103] C.K.E. Wong, C.S. Cobbett, HMA P-type ATPases are the major mechanism for root-to-shoot Cd translocation in *Arabidopsis thaliana*, *New Phytol.* 181 (2009) 71–78.
- [104] L.E. Williams, R.F. Mills, P-1B-ATPases: an ancient family of transition metal pumps with diverse functions in plants, *Trends Plant Sci.* 10 (2005) 491–502.
- [105] D. Major, Stomatal frequency and distribution in rape, *Can. J. Plant Sci.* 55 (1975) 1077–1078.
- [106] L.H. Gammelvind, J.K. Schjoerring, V.O. Mogensen, C.R. Jensen, J.G.H. Bock, Photosynthesis in leaves and siliques of winter oilseed rape (*Brassica napus* L.), *Plant Soil* 186 (1996) 227–236.
- [107] Q. Gu, C. Ferrandiz, M.F. Yanofsky, R. Martienssen, The FRUIT-FULL MADS-box gene mediates cell differentiation during *Arabidopsis* fruit development, *Development* 125 (1998) 1509–1517.
- [108] J.L. Araus, H.R. Brown, A. Febrero, J. Bort, M.D. Serret, Ear photosynthesis, carbon isotope discrimination and the contribution of respiratory CO₂ to differences in grain mass in durum wheat, *Plant Cell Environ.* 16 (1993) 383–392.
- [109] E.A. Tambussi, S. Nogues, J.L. Araus, Ear of durum wheat under water stress: water relations and photosynthetic metabolism, *Planta* 221 (2005) 446–458.
- [110] I. Mukherjee, N.H. Campbell, J.S. Ash, E.L. Connolly, Expression profiling of the *Arabidopsis* ferric chelate reductase (FRO) gene family reveals differential regulation by iron and copper, *Planta* 223 (2006) 1178–1190.
- [111] H.L. Wu, L.H. Li, J. Du, Y.X. Yuan, X.D. Cheng, H.Q. Ling, Molecular and biochemical characterization of the Fe(III) chelate reductase gene family in *Arabidopsis thaliana*, *Plant Cell Physiol.* 46 (2005) 1505–1514.
- [112] A.G.L. Assunção, P.D. Martins, S. De Folter, R. Vooijs, H. Schat, M.G.M. Aarts, Elevated expression of metal transporter genes in three accessions of the metal hyperaccumulator *Thlaspi caerulescens*, *Plant Cell Environ.* 24 (2001) 217–226.
- [113] J.P. Hammond, H.C. Bowen, P.J. White, V. Mills, K.A. Pyke, A.J.M. Baker, S.N. Whiting, S.T. May, M.R. Broadley, A comparison of the *Thlaspi caerulescens* and *Thlaspi arvense* shoot transcriptomes, *New Phytol.* 170 (2006) 239–260.
- [114] H. Küpper, F.J. Zhao, S.P. McGrath, Cellular compartmentation of zinc in leaves of the hyperaccumulator *Thlaspi caerulescens*, *Plant Physiol.* 119 (1999) 305–311.
- [115] H. Küpper, L.O. Seib, M. Sivaguru, O.A. Hoekenga, L.V. Kochian, A method for cellular localization of gene expression via quantitative *in situ* hybridization in plants, *Plant J.* 50 (2007) 159–175.
- [116] H. Küpper, L.V. Kochian, Transcriptional regulation of metal transport genes and mineral nutrition during acclimatization to cadmium and zinc in the Cd/Zn hyperaccumulator, *Thlaspi caerulescens* (Ganges population), *New Phytol.* 185 (2010) 114–129.
- [117] A.G. Desbrosses-Fonrouge, K. Voigt, A. Schröder, S. Arrivault, S. Thomine, U. Krämer, *Arabidopsis thaliana* MTP1 is a Zn transporter in the vacuolar membrane which mediates Zn detoxification and drives leaf Zn accumulation, *FEBS Lett.* 579 (2005) 4165–4174.
- [118] Y. Kobae, T. Uemura, M.H. Sato, M. Ohnishi, T. Mimura, T. Nakagawa, M. Maeshima, Zinc transporter of *Arabidopsis thaliana* AtMTP1 is localized to vacuolar membranes and implicated in zinc homeostasis, *Plant Cell Physiol.* 45 (2004) 1749–1758.
- [119] M. Becher, I.N. Talke, L. Krall, U. Krämer, Cross-species microarray transcript profiling reveals high constitutive expression of metal homeostasis genes in shoots of the zinc hyperaccumulator *Arabidopsis halleri*, *Plant J.* 37 (2004) 251–268.
- [120] N. Yamaji, N. Mitatni, J.F. Ma, A transporter regulating silicon distribution in rice shoots, *Plant Cell* 20 (2008) 1381–1389.
- [121] N. Yamaji, J.F. Ma, A Transporter at the node responsible for intervascular transfer of silicon in rice, *Plant Cell* 21 (2009) 2878–2883.
- [122] D. Nachiangmai, B. Dell, R. Bell, L.B. Huang, B. Rerkasem, Enhanced boron transport into the ear of wheat as a mechanism for boron efficiency, *Plant Soil* 264 (2004) 141–147.
- [123] T. Tsukamoto, H. Nakanishi, H. Uchida, S. Watanabe, S. Matsushashi, S. Mori, N.K. Nishizawa, Fe-52 translocation in barley as monitored by a Positron-Emitting Tracer Imaging System (PETIS): Evidence for the direct translocation of Fe from roots to young leaves via phloem, *Plant Cell Physiol.* 50 (2009) 48–57.
- [124] M. Suzuki, M. Takahashi, T. Tsukamoto, S. Watanabe, S. Matsushashi, J. Yazaki, N. Kishimoto, S. Kikuchi, H. Nakanishi, S. Mori, N.K. Nishizawa, Biosynthesis and secretion of mugineic acid family phytosiderophores in zinc-deficient barley, *Plant J.* 48 (2006) 85–97.
- [125] M. Suzuki, T. Tsukamoto, H. Inoue, S. Watanabe, S. Matsushashi, M. Takahashi, H. Nakanishi, S. Mori, N.K. Nishizawa, Deoxymugineic acid increases Zn translocation in Zn-deficient rice plants, *Plant Mol. Biol.* 66 (2008) 609–617.
- [126] W.H. Zhang, Y.C. Zhou, K.E. Dibley, S.D. Tyerman, R.T. Furbank, J.W. Patrick, Nutrient loading of developing seeds, *Funct. Plant Biol.* 34 (2007) 314–331.

- [127] Y.X. Wang, A. Specht, W.J. Horst, Stable isotope labelling and zinc distribution in grains studied by laser ablation ICP-MS in an ear culture system reveals zinc transport barriers during grain filling in wheat, *New Phytol.* (2010), doi:10.1111/j.1469-8137.2010.03489.x.
- [128] J.N. Pearson, Z. Rengel, Distribution and remobilization of Zn and Mn during grain development in wheat, *J. Exp. Bot.* 45 (1994) 1829–1835.
- [129] B.S. Haslett, R.J. Reid, Z. Rengel, Zinc mobility in wheat: uptake and distribution of zinc applied to leaves or roots, *Ann. Bot.* 87 (2001) 379–386.
- [130] V. Page, L. Weisskopf, U. Feller, Heavy metals in white lupin: uptake, root-to-shoot transfer and redistribution within the plant, *New Phytol.* 171 (2006) 329–341.
- [131] T. Herren, U. Feller, Effect of locally increased zinc contents on zinc transport from the flag leaf lamina to the maturing grains of wheat, *J. Plant Nutr.* 19 (1996) 379–387.
- [132] T. Herren, U. Feller, Transfer of zinc from xylem to phloem in the peduncle of wheat, *J. Plant Nutr.* 17 (1994) 1587–1598.
- [133] C.A. Blindauer, R. Schmid, Cytosolic metal handling in plants: determinants for zinc specificity in metal transporters and metallo-thioneins, *Metallomics* 2 (2010) 510–529.
- [134] M.J. Haydon, C.S. Cobbett, A novel major facilitator superfamily protein at the tonoplast influences zinc tolerance and accumulation in *Arabidopsis*, *Plant Physiol.* 143 (2007) 1705–1719.
- [135] C. Kruger, O. Berkowitz, U.W. Stephan, R. Hell, A metal-binding member of the late embryogenesis abundant protein family transports iron in the phloem of *Ricinus communis* L., *J. Biol. Chem.* 277 (2002) 25062–25069.
- [136] C. Curie, G. Cassin, D. Couch, F. Divol, K. Higuchi, M. Jean, J. Misson, A. Schikora, P. Czernic, S. Mari, Metal movement within the plant: contribution of nicotianamine and yellow stripe 1-like transporters, *Ann. Bot.* 103 (2009) 1–11.
- [137] K. Higuchi, K. Suzuki, H. Nakanishi, H. Yamaguchi, N.K. Nishizawa, S. Mori, Cloning of nicotianamine synthase genes, novel genes involved in the biosynthesis of phytosiderophores, *Plant Physiol.* 119 (1999) 471–479.
- [138] K. Suzuki, K. Higuchi, H. Nakanishi, N.K. Nishizawa, S. Mori, Cloning of nicotianamine synthase genes from *Arabidopsis thaliana*, *Soil Sci. Plant Nutr.* 45 (1999) 993–1002.
- [139] H.Q. Ling, G. Koch, H. Baumlein, M.W. Ganal, Map-based cloning of *chloronerva*, a gene involved in iron uptake of higher plants encoding nicotianamine synthase, *Proc. Natl. Acad. Sci. U.S.A.* 96 (1999) 7098–7103.
- [140] A. Pich, R. Manteuffel, S. Hillmer, G. Scholz, W. Schmidt, Fe homeostasis in plant cells: does nicotianamine play multiple roles in the regulation of cytoplasmic Fe concentration? *Planta* 213 (2001) 967–976.
- [141] A. Pich, G. Scholz, Translocation of copper and other micronutrients in tomato plants (*Lycopersicon esculentum* Mill.): nicotianamine-stimulated copper transport in the xylem, *J. Exp. Bot.* 47 (1996) 41–47.
- [142] U.W. Stephan, M. Grun, Physiological disorders of the nicotianamine-auxotroph tomato mutant *chloronerva* at different levels of iron nutrition. II. Iron deficiency response and heavy metal metabolism, *Biochem. Physiol. Pflanz.* 185 (1989) 189–200.
- [143] M. Takahashi, Y. Terada, I. Nakai, H. Nakanishi, E. Yoshimura, S. Mori, M. Nishizawa, Role of nicotianamine in the intracellular delivery of metals and plant reproductive development, *Plant Cell* 15 (2003) 1263–1280.
- [144] M. Klatte, M. Schuler, M. Wirtz, C. Fink-Straube, R. Hell, P. Bauer, The analysis of *Arabidopsis* nicotianamine synthase mutants reveals functions for nicotianamine in seed iron loading and iron deficiency responses, *Plant Physiol.* 150 (2009) 257–271.
- [145] L.A. Roberts, A.J. Pierson, Z. Panaviene, E.L. Walker, Yellow stripe1. Expanded roles for the maize iron-phytosiderophore transporter, *Plant Physiol.* 135 (2004) 112–120.
- [146] D. Ueno, N. Yamaji, J.F. Ma, Further characterization of ferric-phytosiderophore transporters ZmYS1 and HvYS1 in maize and barley, *J. Exp. Bot.* 60 (2009) 3513–3520.
- [147] H.H. Chu, J. Chiecko, T. Punshon, A. Lanzarotti, B. Lahner, D.E. Salt, E.L. Walker, Successful reproduction requires the function of *Arabidopsis* YELLOW STRIPE-LIKE1 and YELLOW STRIPE-LIKE3 metal-nicotianamine transporters in both vegetative and reproductive structures, *Plant Physiol.* 154 (2010) 197–210.
- [148] R.J. DiDonato Jr., L.A. Roberts, T. Sanderson, R.B. Easley, E.L. Walker, *Arabidopsis* Yellow Stripe-Like2 (YSL2): a metal-regulated gene encoding a plasma membrane transporter of nicotianamine-metal complexes, *Plant J.* 39 (2004) 403–414.
- [149] M. Le Jean, A. Schikora, S. Mari, J.F. Briat, C. Curie, A loss-of-function mutation in AtYSL1 reveals its role in iron and nicotianamine seed loading, *Plant J.* 44 (2005) 769–782.
- [150] B.M. Waters, H.H. Chu, R.J. DiDonato, L.A. Roberts, R.B. Easley, B. Lahner, D.E. Salt, E.L. Walker, Mutations in *Arabidopsis* Yellow Stripe-Like1 and Yellow Stripe-Like3 reveal their roles in metal ion homeostasis and loading of metal ions in seeds, *Plant Physiol.* 141 (2006) 1446–1458.
- [151] Y. Ishimaru, H. Masuda, K. Bashir, H. Inoue, T. Tsukamoto, M. Takahashi, H. Nakanishi, N. Aoki, T. Hirose, R. Ohsugi, N.K. Nishizawa, Rice metal-nicotianamine transporter, OsYSL2, is required for the long-distance transport of iron and manganese, *Plant J.* 62 (2010) 379–390.
- [152] M.G. Stacey, S. Koh, J. Becker, G. Stacey, AtOPT3, a member of the oligopeptide transporter family, is essential for embryo development in *Arabidopsis*, *Plant Cell* 14 (2002) 2799–2811.
- [153] M.G. Stacey, A. Patel, W.E. McClain, M. Mathieu, M. Remley, E.E. Rogers, W. Gassmann, D.G. Blevins, G. Stacey, The *Arabidopsis* AtOPT3 protein functions in metal homeostasis and movement of iron to developing seeds, *Plant Physiol.* 146 (2008) 589–601.
- [154] M.G. Stacey, H. Osawa, A. Patel, W. Gassmann, G. Stacey, Expression analyses of *Arabidopsis* oligopeptide transporters during seed germination, vegetative growth and reproduction, *Planta* 223 (2006) 291–305.
- [155] M.W. Vasconcelos, G.W. Li, M.A. Lubkowitz, M.A. Grusak, Characterization of the PT clade of oligopeptide transporters in rice, *Plant Genome* 1 (2008) 77–88.
- [156] S. Pike, A. Patel, G. Stacey, W. Gassmann, *Arabidopsis* OPT6 is an oligopeptide transporter with exceptionally broad substrate specificity, *Plant Cell Physiol.* 50 (2009) 1923–1932.
- [157] J. Gerendas, U. Schurr, Physicochemical aspects of ion relations and pH regulation in plants: a quantitative approach, *J. Exp. Bot.* 50 (1999) 1101–1114.
- [158] M.L. Gueriot, Y. Yi, Iron: nutritious, noxious, and not readily available, *Plant Physiol.* 104 (1994) 815–820.
- [159] N. von Wiren, S. Klair, S. Bansal, J. -F. Briat, H. Khodr, T. Shioiri, R.A. Leigh, R.C. Hider, Nicotianamine chelates both Fe[III] and Fe[II]. Implications for metal transport in plants, *Plant Physiol.* 119 (1999) 1107–1114.
- [160] I. Cakmak, W.H. Pfeiffer, B. McClafferty, Biofortification of durum wheat with zinc and iron, *Cereal Chem.* 87 (2010) 10–20.
- [161] U.B. Kutman, B. Yildiz, L. Ozturk, I. Cakmak, Biofortification of durum wheat with zinc through soil and foliar applications of nitrogen, *Cereal Chem.* 87 (2010) 1–9.
- [162] C. Uauy, J.C. Brevis, J. Dubcovsky, The high grain protein content gene *Gpc-B1* accelerates senescence and has pleiotropic effects on protein content in wheat, *J. Exp. Bot.* 57 (2006) 2785–2794.
- [163] C. Uauy, A. Distelfeld, T. Fahima, A. Blechl, J. Dubcovsky, A NAC gene regulating senescence improves grain protein, zinc, and iron content in wheat, *Science* 314 (2006) 1298–1301.
- [164] J. Thorne, Phloem unloading of C and N assimilates in developing seeds, *Ann. Rev. Plant Physiol.* 36 (1985) 317–343.
- [165] K. Oparka, P. Gates, Sink anatomy in relation to solute movement in rice (*Oryza sativa* L.): a summary of findings, *Plant Growth Regul.* 2 (1984) 297–307.
- [166] B. Tauris, S. Borg, P.L. Gregersen, P.B. Holm, A roadmap for zinc trafficking in the developing barley grain based on laser capture microdissection and gene expression profiling, *J. Exp. Bot.* 60 (2009) 1333–1347.
- [167] H. Roschztardt, G. Conéjéro, C. Curie, S. Mari, Identification of the endodermal vacuole as the iron storage compartment in the *Arabidopsis* embryo, *Plant Physiol.* 151 (2009) 1329–1338.

- [168] S.A. Kim, T. Punshon, A. Lanzirrotti, L.T. Li, J.M. Alonso, J.R. Ecker, J. Kaplan, M.L. Guerinot, Localization of iron in *Arabidopsis* seed requires the vacuolar membrane transporter VIT1, *Science* 314 (2006) 1295–1298.
- [169] V. Lanquar, F. Lelièvre, S. Bolte, C. Hamès, C. Alcon, D. Neumann, G. Vansuyt, C. Curie, A. Schröder, U. Krämer, H. Barbier-Brygoo, S. Thomine, Mobilization of vacuolar iron by AtNRAMP3 and AtNRAMP4 is essential for seed germination on low iron, *EMBO J.* 24 (2005) 4041–4051.
- [170] J.-F. Briat, S. Lobreaux, Iron transport and storage in plants, *Trends Plant Sci.* 2 (1997) 187–193.
- [171] S. Lobreaux, J.F. Briat, Ferritin accumulation and degradation in different organs of pea (*Pisum sativum*) during development, *Biochem. J.* 274 (1991) 601–606.
- [172] E. Marentes, M.A. Grusak, Iron transport and storage within the seed coat and embryo of developing seeds of pea (*Pisum sativum* L.), *Seed Sci. Res.* 8 (1998) 367–375.
- [173] M. Hoppler, A. Schönbachler, L. Meile, R.F. Hurrell, T. Walczyk, Ferritin-iron is released during boiling and *in vitro* gastric digestion, *J. Nutr.* 138 (2008) 878–884.
- [174] C. Cvitanič, W.J. Przybyłowicz, D.F. Urbanski, A.M. Jurkiewicz, J. Mesjasz-Przybyłowicz, M.W. Blair, C. Astudillo, E.O. Jensen, J. Stougaard, Iron and ferritin accumulate in separate cellular locations in *Phaseolus* seeds, *BMC Plant Biol.* 10 (2010) 26.
- [175] J.T. Moraghan, Accumulation and within-seed distribution of iron in common bean and soybean, *Plant Soil* 264 (2004) 287–297.
- [176] J.T. Moraghan, J. Padilla, J.D. Etchevers, K. Grafton, J.A. Acosta-Gallegos, Iron accumulation in seed of common bean, *Plant Soil* 246 (2002) 175–183.
- [177] L. Ozturk, M.A. Yazici, C. Yucel, A. Torun, C. Cekiç, A. Bağcı, H. Ozkan, H.J. Braun, Z. Sayers, I. Cakmak, Concentration and localization of zinc during seed development and germination in wheat, *Physiol. Plant* 128 (2006) 144–152.
- [178] V. Raboy, Approaches and challenges to engineering seed phytate and total phosphorus, *Plant Sci.* 177 (2009) 281–296.
- [179] T. Frank, R. Habernegg, F.J. Yuan, Q.Y. Shu, K.H. Engel, Assessment of the contents of phytic acid and divalent cations in low phytic acid (*lpa*) mutants of rice and soybean, *J. Food Compos. Anal.* 22 (2009) 278–284.
- [180] K. Liu, K.L. Peterson, V. Raboy, Comparison of the phosphorus and mineral concentrations in bran and abraded kernel fractions of a normal barley (*Hordeum vulgare*) cultivar versus four low phytic acid isolines, *J. Agric. Food Chem.* 55 (2007) 4453–4460.
- [181] M.J. Guttieri, K.M. Peterson, E.J. Souza, Mineral distributions in milling fractions of low phytic acid wheat, *Crop Sci.* 46 (2006) 2692–2698.
- [182] L. Lin, I. Ockenden, J.N.A. Lott, The concentrations and distribution of phytic acid-phosphorus and other mineral nutrients in wild-type and low phytic acid1-1 (*lpa1-1*) corn (*Zea mays* L.) grains and grain parts, *Can. J. Bot.* 83 (2005) 131–141.
- [183] J.C. Liu, I. Ockenden, M. Truax, J.N.A. Lott, Phytic acid-phosphorus and other nutritionally important mineral nutrient elements in grains of wild-type and low phytic acid (*lpa1-1*) rice, *Seed Sci. Res.* 14 (2004) 109–116.
- [184] K.M. Hambidge, J.W. Huffer, V. Raboy, G.K. Grunwald, J.L. Westcott, L. Sian, L.V. Miller, J.A. Dorsch, N.F. Krebs, Zinc absorption from low-phytate hybrids of maize and their wild-type isohybrids, *Am. J. Clin. Nutr.* 79 (2004) 1053–1059.
- [185] C.L. Adams, M. Hambidge, V. Raboy, J.A. Dorsch, L. Sian, J.L. Westcott, N.F. Krebs, Zinc absorption from a low-phytic acid maize, *Am. J. Clin. Nutr.* 76 (2002) 556–559.
- [186] I.R.G.S. Project, The map-based sequence of the rice genome, *Nature* 436 (2005) 793–800.
- [187] A.H. Paterson, J.E. Bowers, R. Bruggmann, I. Dubchak, J. Grimwood, H. Gundlach, G. Haberer, U. Hellsten, T. Mitros, A. Poljakov, J. Schmutz, M. Spannagl, H.B. Tang, X.Y. Wang, T. Wicker, A.K. Bharti, J. Chapman, F.A. Feltus, U. Gowik, I.V. Grigoriev, E. Lyons, C.A. Maher, M. Martis, A. Narechania, R.P. O'tillar, B.W. Penning, A.A. Salamov, Y. Wang, L.F. Zhang, N.C. Carpita, M. Freeling, A.R. Gingle, C.T. Hash, B. Keller, P. Klein, S. Kresovich, M.C. McCann, R. Ming, D.G. Peterson, R. Mehboobur, D. Ware, P. Westhoff, K.F.X. Mayer, J. Messing, D.S. Rokhsar, The *Sorghum bicolor* genome and the diversification of grasses, *Nature* 457 (2009) 551–556.
- [188] J. Schmutz, S.B. Cannon, J. Schlueter, J.X. Ma, T. Mitros, W. Nelson, D.L. Hyten, Q.J. Song, J.J. Thelen, J.L. Cheng, D. Xu, U. Hellsten, G.D. May, Y. Yu, T. Sakurai, T. Umezawa, M.K. Bhattacharyya, D. Sandhu, B. Valliyodan, E. Lindquist, M. Peto, D. Grant, S.Q. Shu, D. Goodstein, K. Barry, M. Futrell-Griggs, B. Abernathy, J.C. Du, Z.X. Tian, L.C. Zhu, N. Gill, T. Joshi, M. Libault, A. Sethuraman, X.C. Zhang, K. Shinozaki, H.T. Nguyen, R.A. Wing, P. Cregan, J. Specht, J. Grimwood, D. Rokhsar, G. Stacey, R.C. Shoemaker, S.A. Jackson, Genome sequence of the palaeopolyploid soybean, *Nature* 463 (2010) 178–183.
- [189] R. Lister, B.D. Gregory, J.R. Ecker, Next is now: new technologies for sequencing of genomes, transcriptomes, and beyond, *Curr. Opin. Plant Biol.* 12 (2009) 107–118.
- [190] Z. Wang, M. Gerstein, M. Snyder, RNA-Seq: a revolutionary tool for transcriptomics, *Nat. Rev. Genet.* 10 (2009) 57–63.
- [191] R.W. Doerge, Mapping and analysis of quantitative trait loci in experimental populations, *Nat. Rev. Genet.* 3 (2002) 43–52.
- [192] T.F.C. Mackay, The genetic architecture of quantitative traits, *Annu. Rev. Genet.* 35 (2001) 303–339.
- [193] S. Myles, J. Peiffer, P.J. Brown, E.S. Ersoz, Z.W. Zhang, D.E. Costich, E.S. Buckler, Association mapping: critical considerations shift from genotyping to experimental design, *Plant Cell* 21 (2009) 2194–2202.
- [194] C. Zhu, M. Gore, E.S. Buckler, J. Yu, Status and prospects of association mapping in plants, *Plant Genome* 1 (2008) 5–20.
- [195] A. Ghandilyan, N. Ilk, C. Hanhart, M. Mbengue, L. Barboza, H. Schat, M. Koornneef, M. El-Lithy, D. Vreugdenhil, M. Reymond, M.G.M. Aarts, A strong effect of growth medium and organ type on the identification of QTLs for phytate and mineral concentrations in three *Arabidopsis thaliana* RIL populations, *J. Exp. Bot.* 60 (2009) 1409–1425.
- [196] D. Vreugdenhil, M.G.M. Aarts, M. Koornneef, H. Nelissen, W.H.O. Ernst, Natural variation and QTL analysis for cationic mineral content in seeds of *Arabidopsis thaliana*, *Plant Cell Environ.* 27 (2004) 828–839.
- [197] B.M. Waters, M.A. Grusak, Quantitative trait locus mapping for seed mineral concentrations in two *Arabidopsis thaliana* recombinant inbred populations, *New Phytol.* 179 (2008) 1033–1047.
- [198] R.P. Sankaran, T. Huguet, M.A. Grusak, Identification of QTL affecting seed mineral concentrations and content in the model legume *Medicago truncatula*, *Theor. Appl. Genet.* 119 (2009) 241–253.
- [199] M.A. Klein, M.A. Grusak, Identification of nutrient and physical seed trait QTL in the model legume *Lotus japonicus*, *Genome* 52 (2009) 677–691.
- [200] M. Blair, C. Astudillo, M. Grusak, R. Graham, S. Beebe, Inheritance of seed iron and zinc concentrations in common bean (*Phaseolus vulgaris* L.), *Mol. Breed.* 23 (2009) 197–207.
- [201] K.A. Cichy, G.V. Caldas, S.S. Snapp, M.W. Blair, QTL analysis of seed iron, zinc, and phosphorus levels in an Andean bean population, *Crop Sci.* 49 (2009) 1742–1750.
- [202] S.H. Guzman-Maldonado, O. Martinez, J.A. Acosta-Gallegos, F. Guevara-Lara, O. Paredes-Lopez, Putative quantitative trait loci for physical and chemical components of common bean, *Crop Sci.* 43 (2003) 1029–1035.
- [203] G.D. Ding, M. Yang, Y.F. Hu, Y.A. Liao, L. Shi, F.S. Xu, J.L. Meng, Quantitative trait loci affecting seed mineral concentrations in *Brassica napus* grown with contrasting phosphorus supplies, *Ann. Bot.* 105 (2010) 1221–1234.
- [204] J.J. Zhao, D.C.L. Jamar, P. Lou, Y.H. Wang, J. Wu, X.W. Wang, G. Bonnema, M. Koornneef, D. Vreugdenhil, Quantitative trait loci analysis of phytate and phosphate concentrations in seeds and leaves of *Brassica rapa*, *Plant Cell Environ.* 31 (2008) 887–900.
- [205] J.C.R. Stangoulis, B.L. Huynh, R.M. Welch, E.Y. Choi, R.D. Graham, Quantitative trait loci for phytate in rice grain and their relationship with grain micronutrient content, *Euphytica* 154 (2007) 289–294.

- [206] G.J. Norton, C.M. Deacon, L.Z. Xiong, S.Y. Huang, A.A. Meharg, A.H. Price, Genetic mapping of the rice ionome in leaves and grain: identification of QTLs for 17 elements including arsenic, cadmium, iron and selenium, *Plant Soil* 329 (2010) 139–153.
- [207] A.L. Garcia-Oliveira, L.B. Tan, Y.C. Fu, C.Q. Sun, Genetic identification of quantitative trait loci for contents of mineral nutrients in rice grain, *J. Integr. Plant Biol.* 51 (2009) 84–92.
- [208] Z. Peleg, I. Cakmak, L. Ozturk, A. Yazici, Y. Jun, H. Budak, A.B. Korol, T. Fahima, Y. Saranga, Quantitative trait loci conferring grain mineral nutrient concentrations in durum wheat × wild emmer wheat RIL population, *Theor. Appl. Genet.* 119 (2009) 353–369.
- [209] R.L. Shi, H.W. Li, Y.P. Tong, R.L. Jing, F.S. Zhang, C.Q. Zou, Identification of quantitative trait locus of zinc and phosphorus density in wheat (*Triticum aestivum* L.) grain, *Plant Soil* 306 (2008) 95–104.
- [210] J. Wu, F.J. Zhao, A. Ghandilyan, B. Logoteta, M.O. Guzman, H. Schat, X.W. Wang, M.G.M. Aarts, Identification and functional analysis of two ZIP metal transporters of the hyperaccumulator *Thlaspi caerulescens*, *Plant Soil* 325 (2009) 79–95.
- [211] M. Weber, E. Harada, C. Vess, E. von Roepenack-Lahaye, S. Clemens, Comparative microarray analysis of *Arabidopsis thaliana* and *Arabidopsis halleri* roots identifies nicotianamine synthase, a ZIP transporter and other genes as potential metal hyperaccumulation factors, *Plant J.* 37 (2004) 269–281.
- [212] J.E. van de Mortel, L.A. Villanueva, H. Schat, J. Kwekkeboom, S. Coughlan, P.D. Moerland, E.V.L. van Themaat, M. Koornneef, M.G.M. Aarts, Large expression differences in genes for iron and zinc homeostasis, stress response, and lignin biosynthesis distinguish roots of *Arabidopsis thaliana* and the related metal hyperaccumulator *Thlaspi caerulescens*, *Plant Physiol.* 142 (2006) 1127–1147.
- [213] I.N. Talke, M. Hanikenne, U. Krämer, Zinc-dependent global transcriptional control, transcriptional deregulation, and higher gene copy number for genes in metal homeostasis of the hyperaccumulator *Arabidopsis halleri*, *Plant Physiol.* 142 (2006) 148–167.
- [214] V. Filatov, J. Dowdle, N. Smirnov, B. Ford-Lloyd, H.J. Newbury, M.R. Macnair, Comparison of gene expression in segregating families identifies genes and genomic regions involved in a novel adaptation, zinc hyperaccumulation, *Mol. Ecol.* 15 (2006) 3045–3059.
- [215] D.B. Dräger, A.G. Desbrosses-Fonrouge, C. Krach, A.N. Chardonens, R.C. Meyer, P. Saumitou-Laprade, U. Krämer, Two genes encoding *Arabidopsis halleri* MTP1 metal transport proteins co-segregate with zinc tolerance and account for high MTP1 transcript levels, *Plant J.* 39 (2004) 425–439.
- [216] H.C. Chiang, J.C. Lo, K.C. Yeh, Genes associated with heavy metal tolerance and accumulation in Zn/Cd hyperaccumulator *Arabidopsis halleri*: a genomic survey with cDNA microarray, *Environ. Sci. Technol.* 40 (2006) 6792–6798.
- [217] M.R. Broadley, P.J. White, J.P. Hammond, I. Zelko, A. Lux, Zinc in plants, *New Phytol.* 173 (2007) 677–702.
- [218] E. Van der Graaff, R. Schwacke, A. Schneider, M. Desimone, U.I. Flugge, R. Kunze, Transcription analysis of *Arabidopsis* membrane transporters and hormone pathways during developmental and induced leaf senescence, *Plant Physiol.* 141 (2006) 776–792.
- [219] S.A. Ramesh, S. Choimes, D.P. Schachtman, Over-expression of an *Arabidopsis* zinc transporter in *Hordeum vulgare* increases short-term zinc uptake after zinc deprivation and seed zinc content, *Plant Mol. Biol.* 54 (2004) 373–385.
- [220] Y. Ishimaru, H. Masuda, M. Suzuki, K. Bashir, M. Takahashi, H. Nakanishi, S. Mori, N.K. Nishizawa, Overexpression of the *OsZIP4* zinc transporter confers disarrangement of zinc distribution in rice plants, *J. Exp. Bot.* 58 (2007) 2909–2915.
- [221] W. Jiang, P.C. Struik, H. van Keulen, M. Zhao, L.N. Jin, T.J. Stomph, Does increased zinc uptake enhance grain zinc mass concentration in rice? *Ann. Appl. Biol.* 153 (2008) 135–147.
- [222] M. Vasconcelos, H. Eckert, V. Arahana, G. Graef, M.A. Grusak, T. Clemente, Molecular and phenotypic characterization of transgenic soybean expressing the *Arabidopsis* ferric chelate reductase gene, *FRO2*, *Planta* 224 (2006) 1116–1128.
- [223] F. Goto, T. Yoshihara, N. Shigemoto, S. Toki, F. Takaiwa, Iron fortification of rice seed by the soybean ferritin gene, *Nat. Biotechnol.* 17 (1999) 282–286.
- [224] M. Vasconcelos, K. Datta, N. Oliva, M. Khalekuzzaman, L. Torrizo, S. Krishnan, M. Oliveira, F. Goto, S.K. Datta, Enhanced iron and zinc accumulation in transgenic rice with the ferritin gene, *Plant Science* 164 (2003) 371–378.
- [225] L.Q. Qu, T. Yoshihara, A. Ooyama, F. Goto, F. Takaiwa, Iron accumulation does not parallel the high expression level of ferritin in transgenic rice seeds, *Planta* 222 (2005) 225–233.
- [226] G. Drakakaki, S. Marcel, R.P. Glahn, E.K. Lund, S. Pariagh, R. Fischer, P. Christou, E. Stoger, Endosperm-specific co-expression of recombinant soybean ferritin and *Aspergillus* phytase in maize results in significant increases in the levels of bioavailable iron, *Plant Mol. Biol.* 59 (2005) 869–880.
- [227] S. Kim, M. Takahashi, K. Higuchi, K. Tsunoda, H. Nakanishi, E. Yoshimura, S. Mori, N.K. Nishizawa, Increased nicotianamine biosynthesis confers enhanced tolerance of high levels of metals, in particular nickel, to plants, *Plant Cell Physiol.* 46 (2005) 1809–1818.
- [228] S. Lee, U.S. Jeon, S.J. Lee, Y.K. Kim, D.P. Persson, S. Husted, J.K. Schjorring, Y. Kakei, H. Masuda, N.K. Nishizawa, G. An, Iron fortification of rice seeds through activation of the nicotianamine synthase gene, *Proc. Natl. Acad. Sci. U.S.A.* 106 (2009) 22014–22019.
- [229] J. Wirth, S. Poletti, B. Aeschlimann, N. Yakandawala, B. Drosse, S. Osorio, T. Tohge, A.R. Fernie, D. Gunther, W. Gruissem, C. Sautter, Rice endosperm iron biofortification by targeted and synergistic action of nicotianamine synthase and ferritin, *Plant Biotechnol. J.* 7 (2009) 631–644.