

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USGS Staff -- Published Research

US Geological Survey

2015

Investigating Endocrine and Physiological Parameters of Captive American Kestrels Exposed by Diet to Selected Organophosphate Flame Retardants

Kimberly J. Fernie
Ecotoxicology & Wildlife Health Division

Vince Palace
Stantec Consulting Ltd.


Lisa E. Peters
Stantec Consulting Ltd.

Nil Basu
McGill University

Robert J. Letcher
Carleton University

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/usgsstaffpub>

 Part of the [Geology Commons](#), [Oceanography and Atmospheric Sciences and Meteorology Commons](#), [Other Earth Sciences Commons](#), and the [Other Environmental Sciences Commons](#)

Fernie, Kimberly J.; Palace, Vince; Peters, Lisa E.; Basu, Nil; Letcher, Robert J.; Karouna-Renier, Natalie K.; Schultz, Sandra L.; Lazarus, Rebecca S.; and Rattner, Barnett A., "Investigating Endocrine and Physiological Parameters of Captive American Kestrels Exposed by Diet to Selected Organophosphate Flame Retardants" (2015). *USGS Staff -- Published Research*. 959.
<https://digitalcommons.unl.edu/usgsstaffpub/959>

This Article is brought to you for free and open access by the US Geological Survey at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USGS Staff -- Published Research by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Kimberly J. Fernie, Vince Palace, Lisa E. Peters, Nil Basu, Robert J. Letcher, Natalie K. Karouna-Renier, Sandra L. Schultz, Rebecca S. Lazarus, and Barnett A. Rattner

Investigating Endocrine and Physiological Parameters of Captive American Kestrels Exposed by Diet to Selected Organophosphate Flame Retardants

Kimberly J. Fernie,^{*,†} Vince Palace,[‡] Lisa E. Peters,[‡] Nil Basu,[§] Robert J. Letcher,^{||} Natalie K. Karouna-Renier,[⊥] Sandra L. Schultz,[⊥] Rebecca S. Lazarus,[⊥] and Barnett A. Rattner[⊥]

[†]Ecotoxicology & Wildlife Health Division, Environment Canada, 867 Lakeshore Road, Burlington, ON, Canada L7R 4A6

[‡]Stantec Consulting Ltd., 386 Broadway Avenue, Winnipeg, MB, Canada R3C 3R6

[§]McGill University, 21,111 Lakeshore Road, Ste Anne de Bellevue, PQ, Canada H9X 3V9

^{||}Ecotoxicology & Wildlife Health Division, Environment Canada, Carleton University, 1125 Colonel By Drive, Ottawa, ON, Canada K1A 0H3

[⊥]U.S. Geological Survey, Patuxent Wildlife Research Center, BARC East Building 308, 10300 Baltimore Avenue, Beltsville, Maryland 20705, United States

S Supporting Information

ABSTRACT: Organophosphate triesters are high production volume additive flame retardants (OPFRs) and plasticizers. Shown to accumulate in abiotic and biotic environmental compartments, little is known about the risks they pose. Captive adult male American kestrels (*Falco sparverius*) were fed the same dose (22 ng OPFR/g kestrel/d) daily (21 d) of tris(2-butoxyethyl) phosphate (TBOEP), tris(2-chloroethyl) phosphate (TCEP), tris(2-chloroisopropyl) phosphate (TCIPP), or tris(1,2-dichloro-2-propyl) phosphate (TDCIPP). Concentrations were undetected in tissues (renal, hepatic), suggesting rapid metabolism. There were no changes in glutathione status, indicators of hepatic oxidative status, or the cholinergic system (i.e., cerebrum, plasma cholinesterases; cerebrum muscarinic, nicotinic receptors). Modest changes occurred in hepatocyte integrity and function (clinical chemistry). Significant effects on plasma free triiodothyronine (FT₃) concentrations occurred with exposure to TBOEP, TCEP, TCIPP, and TDCIPP; TBOEP and TCEP had additional overall effects on free thyroxine (FT₄), whereas TDCIPP also influenced total thyroxine (TT₄). Relative increases (32%–96%) in circulating FT₃, TT₃, FT₄, and/or TT₄ were variable with each OPFR at 7 d exposure, but limited thereafter, which was likely maintained through decreased thyroid gland activity and increased hepatic deiodinase activity. The observed physiological and endocrine effects occurred at environmentally relevant concentrations and suggest parent OPFRs or metabolites may have been present despite rapid degradation.



INTRODUCTION

Flame retardant (FR) compounds are used to hinder the ignition of fire. Some FRs (e.g., polybrominated diphenyl ether (PBDE) congeners in the PentaBDE and OctaBDE technical formulations) have been regulated and phased-out, resulting in the emergence and increasing usage of replacement FRs (e.g., organophosphate (OP) triester FRs).¹ OP triesters were introduced in the 1970s and are now widely used as additive flame retardants (OPFRs) (e.g., polyurethane foam, textiles) and plasticizers (e.g., floor polish, thermoplastics, lacquers). They are high production volume FRs that constituted 14% of the 1.5 million tonnes of FRs used globally in 2004,^{1,2} and 20% of European FR usage in 2006.¹ More recently (2012) in the United States, the annual production volume of tris(2-chloroisopropyl) phosphate (TCIPP) (CAS#: 13674-84-5) was reported as ~25 million kg and ~4.5–22.5 million kg for

tris(1,2-dichloro-2-propyl) phosphate (TDCIPP) (CAS#: 13674-87-8).³ Total usage quantities of TCIPP in Denmark, Finland, Norway, and Sweden were reported to range approximately from 1.1 to 2 million kg from 2007 to 2011.⁴ Some Nordic countries have also reportedly used TDCIPP⁴ and tris(2-chloroethyl) phosphate (TCEP) (CAS#: 51805-45-9) (annual usage volume in 2008: 0.2–261 tonnes¹).

OPFRs, including TCIPP, TDCIPP, TCEP, and tris(2-butoxyethyl) phosphate (TBOEP) (CAS#: 78-51-3), have been detected in abiotic environmental compartments.^{5,6} In the Great Lakes, atmospheric concentrations of summed (Σ)

Received: February 16, 2015

Revised: May 18, 2015

Accepted: May 19, 2015

Published: May 19, 2015

OPFRs exceeded those of brominated flame retardants (BFRs) by 2–3 orders of magnitude.⁷ OPFRs have also been detected in airborne particles over the Southern, Arctic, Pacific, and Indian Oceans⁸ and the North Sea,⁹ suggesting long-range transport.

Various OPFRs, including TBOEP, TCEP, TCIPP, and/or TDCIPP, have also been detected in aquatic birds in the northern hemisphere^{10–12} over the past 25 years.¹² Concentrations of 16 OPFRs varied in tissues of herring gulls (*Larus argentatus*) with TDCIPP and TCIPP dominant in egg yolk, maternal muscle, and red blood cells, TBOEP prevailing in albumen and maternal fat, and low concentrations of TCEP limited to eggs.² In the same gulls, plasma concentrations of triphenyl phosphate (TPHP) were also detected as were OPFR diester metabolites, indicating that OPFR triesters can be metabolized by birds.¹³ Feather concentrations of six OPFRs in nestling white-tailed eagles (*Haliaeetus albicilla*) were much greater than PBDEs, non-PBDE FRs, polychlorinated biphenyl 153, and dichlorodiphenyldichloroethylene (*p,p'*-DDE).¹⁴

Little is known about the potential toxic effects of OPFR triesters in humans and biota, including birds. Exposure to TDCIPP and TCIPP altered mRNA abundance of genes associated with phase I and II metabolism, thyroid hormone and lipid regulation, and growth in chicken embryonic hepatocytes (CEH).¹⁵ In chicken embryos exposed by egg injection, very high concentrations of TCIPP (9,240 and 51,600 ng/g) significantly delayed pipping and reduced growth (51,600 ng/g), and TDCIPP reduced embryo mass, gallbladder size (45,000 ng/g), and free thyroxine (FT₄) concentrations (7,640 ng/g).¹⁶ Exposure of chicken embryos to TBOEP induced hepatic CYP3A37 mRNA, and the effects of triethyl phosphate (TEP) were similar to TCIPP and/or TDCIPP: decreased pipping success, altered growth and plasma bile acids, modulated genes associated with xenobiotic and lipid metabolism and the thyroid hormone pathway, and reduced plasma FT₄ levels.¹⁷

Recently, several OPFRs, including TBOEP, TCEP, TCIPP, and TDCIPP, were listed as priority chemicals for risk assessment by Environment Canada. The risk of some chlorinated OP esters is also being evaluated by the U.S. Environmental Protection Agency. Investigations of the biological effects of specific OPFRs on avian embryos following in vitro and egg-injection exposures were recently reported.^{15–18} However, data from ecologically relevant models and other life stages are lacking. Here, the American kestrel (*Falco sparverius*) is used as a raptorial model species because it has been used extensively in avian ecotoxicological studies.^{19–25} Kestrels were found to be more sensitive to the effects of PBDEs than galliform species (e.g., chickens²⁴ or common terns (*Sterna hirundo*)).²⁵ The in vivo uptake and effects of chronic, dietary exposure to TBOEP, TCEP, TCIPP, and TDCIPP were investigated with captive kestrels, focusing on physiological systems (e.g., hepatic oxidative status, thyroid, and hepatic function) previously shown to be sensitive to OPFR exposure.^{15–18}

MATERIALS AND METHODS

Kestrel Dosing and Sampling. This study was conducted in 2011 and 2012 under guidelines approved by the Animal Care and Use Committee at the USGS Patuxent Wildlife Research Center (PWRC; Laurel, MD, USA). Captive American kestrels were housed individually in large outdoor flight pens with a shade roof, perches, food tray, and water

bowl, and shelter was available from the elements.²⁶ The birds were fed day-old dead cockerel ad libitum daily, receiving the dosed cockerel each morning and either nondosed cockerel or dead laboratory mice on alternate afternoons. We used the same dietary feeding-exposure methods of previous flame retardant studies in which captive American kestrels accumulated concentrations of PBDEs, hexabromocyclododecane (HBCDD), decabromodiphenyl ether (BDE-209), but not β -1,2-dibromo-4-(1,2-dibromoethyl)cyclohexane.^{19–23} Dosing solutions were injected into the cerebral hemispheres of the cockerel, a preferred dietary item of captive kestrels. Food consumption was monitored daily, and all dosed cockerel were consumed daily with minor exceptions.

A pilot study was conducted in 2011 using 7 adult two-year old male kestrels exposed daily by diet for 28 days to TDCIPP and 7 adult male kestrels of the same age exposed to only the safflower oil vehicle. The details and results of this pilot study are presented in the Supporting Information (SI). In 2012, a study was conducted to understand the potential uptake and effects of all four priority OPFRs. On the basis of the findings of this pilot study, a higher concentration of TDCIPP was used to characterize possible effects of elevated concentrations likely experienced by wild birds.^{12,27} In this study, a separate set of 38 one-year old adult male kestrels (7 per exposure group with 10 controls) were exposed daily by diet to TDCIPP, TBOEP, TCIPP, or TCEP for 21 days because potential effects of the higher dose were anticipated to occur within this time frame. Control male birds of the same age were exposed to the safflower oil vehicle only. Each kestrel ate one cockerel per day that had been injected with 50 μ L of a safflower oil solution containing one of each of the four OPFRs. On the basis of the mean weight (114 g) of the adult male kestrels at the initiation of the study, each kestrel received a daily dose of 2500 ng OPFR/50 μ L safflower oil/day, corresponding to 21.93 ng OPFR/g kestrel/day.

For the purposes of this study, TBOEP (>94% purity) and TCEP (>97% purity) were purchased from Sigma-Aldrich (St. Louis, MI, USA); TDCIPP (>95% purity) was purchased from TCI America (Portland, OR, USA), and TCIPP (>95% purity) was purchased from AK Scientific (Union City, CA, USA). The nominal (calculated) concentrations of OPFRs in the safflower oil dosing solutions were TBOEP at 49 μ g/ μ L, TCIPP at 52 μ g/ μ L, TCEP at 52 μ g/ μ L, and TDCIPP at 51 μ g/ μ L. The nominal concentrations were very accurate as the actual measured concentrations of each OPFR in the dosing solutions were as follows: TBOEP, 50 μ g/ μ L; TCIPP, 59 μ g/ μ L; TCEP, 41 μ g/ μ L; and TDCIPP, 53 μ g/ μ L.

The kestrels were weighed, and their general health assessed at the beginning of the exposure period and again at 7, 14, and 21 d of the exposure trial. Immediately after weighing, between 0.6 and 1.1 mL of whole blood was withdrawn from each kestrel by jugular venipuncture using a heparinized 1.0 mL syringe with a 27-gauge needle. Following their exposure, the birds were humanely euthanized by carbon dioxide. Further blood was immediately collected by cardiac venipuncture with a subsample analyzed for hematocrit and the remainder stored on ice until processing by centrifugation. Plasma was aliquoted into cryovials and stored at -80°C . For each kestrel, the thyroid gland, liver, kidney, pectoral muscle (~ 1 g), and brain were immediately collected, weighed, subsectioned, and stored in 10% buffered formalin for histological assessment or frozen at -80°C . Somatic indices (liver, kidney, brain) were

calculated as organ mass divided by body mass with the result then multiplied by 100.

Clinical Chemistry Analysis. All plasma samples were stored at -70°C and shipped on dry ice for clinical chemistry analysis by Dr. Carolyn Cray (University of Miami). Only plasma samples free of lipemia and hemolysis were analyzed for alkaline phosphatase activity (ALP), alanine aminotransferase activity (ALT), gamma-glutamyl transpeptidase (GGT) enzymatic activity, total protein (TP), prealbumin and albumin (ALB), α_1 , α_2 , β , and γ globulins (G), aspartate aminotransferase (ASAT), glutamate dehydrogenase activities (GLDH), lactate dehydrogenase isozymes (LDH), hemolysis, and lipemia indices. The A:G ratio was calculated as the sum of the prealbumin and albumin divided by the sum of all measured globulins. Heparinized plasma samples were used for biochemical determinations. Routine biochemical analyses (ALT, LDH, GGT, ALP, and TP) were conducted using an Ortho Vitros 250XR chemistry analyzer (Ortho, Rochester, NY, USA).²⁸ Further analytical details are provided in the SI.

Neurochemical Assays. Neurochemical assays were completed at the University of Michigan based on previous analytical methods.^{29,30} Homogenate and membrane preparations from the cerebrum tissues were prepared as described previously.^{29–31} Brains were homogenized in 50 mM Tris buffer (50 mM Tris HCl, 50 mM Tris Base, pH 7.4), and an aliquot of homogenate was saved for enzyme analyses. Membranes were isolated by centrifuging at 48000g for 15 min. The pellet was washed three times and resuspended in Tris buffer. Protein content of homogenate and membranes was measured using the Bradford assay, and samples were stored at -80°C . The cholinergic assays (receptor binding to muscarinic and nicotinic receptors; enzyme activity of acetylcholinesterase and butyrylcholinesterase) were performed as previously described.³⁰ A detailed description of these analytical methods is provided in the SI.

Oxidative Status. Livers collected in 2012 were analyzed for indicators of glutathione status and oxidative status. Reduced glutathione (GSH), oxidized glutathione (GSSG), and the GSSG/GSH ratio were analyzed using the DetectX Glutathione Fluorescent Detection Kit (Arbor Assays, Ann Arbor, MI, USA), total sulfhydryl (TSH) using the Measure-iT Thiol Assay Kit (Life Technologies - Molecular Probes, Inc., Eugene, OR, USA), protein bound sulfhydryl (PBSH) by calculation of the difference between TSH and GSH, and lipid peroxidation (thiobarbituric acid reactive substances, TBARS) using the QuantiChrom TBARS Assay Kit (Bioassay Systems, Hayward, CA, USA). All assays were validated prior to sample analysis to determine the detection limit and assess accuracy, precision, and dilutional parallelism and to identify any matrix effects (none). Each sample was analyzed in duplicate with a mean coefficient of variation (CV; mean \pm standard deviation) of $2.01 \pm 1.47\%$, $4.26 \pm 2.09\%$, and $1.25 \pm 0.68\%$ / $2.44 \pm 1.53\%$ for TBARS, TSH, and reduced/total GSH, respectively. Duplicate samples for which CVs exceeded 10% were reanalyzed. A detailed description of the analytical methods used for the neurochemical assays and oxidative stress assays are provided in the SI.

Analysis of Thyroid Function. A three-pronged assessment of thyroid function was performed by determining circulating triiodothyronine (T_3) and T_4 concentrations, hepatic T_4 -outer ring deiodinase activity (T_4 -ORD), and histological assessments of the structure of the thyroid gland. Circulating concentrations of free and total T_3 (FT₃, TT₃) and

T_4 (FT₄, TT₄) were determined according to the manufacturer's instructions using commercially available radioimmunoassays (MP Biomedical, Santa Ana, CA, USA) and based on linear response versus concentration calibration curves (R^2 : $T_3 = 0.997$, $T_4 = 0.953$) across the physiological range of hormone concentrations ($T_3 = 0.5\text{--}8\text{ ng/mL}$, $T_4 = 20\text{--}200\text{ ng/mL}$). Hepatic deiodinase activity was determined as previously described with American kestrels.³² The thyroid gland of each kestrel was assessed for structural changes in epithelial cell height (ECH) and colloid parameters as previously described.³³ Full details for assessments of hepatic deiodinase activity and thyroid gland histology are provided in the SI.

Chemical Analysis. In the Organic Contaminants Research Laboratory (OCRL)/Letcher Laboratories at Environment Canada, concentrations of TBOEP, TCEP, TCIPP, and TDCIPP were determined in liver and kidney tissue samples in isolated fractions.^{2,13,34} Liquid chromatography-electrospray-tandem quadrupole-mass spectrometry (LC-ESI(+)-MS/MS) was used to determine the OPFRs in the isolated fractions using the following MRM transitions: TCEP (m/z 284 > 63), TCIPP (m/z 329.1 > 99), TDCIPP (m/z 430.9 > 99), TBOEP (m/z 294.3 > 102). Mean recovery (± 1 SD) of the internal standard d_{27} -TBP was $85 \pm 15\%$ for all liver and kidney analyses. An internal standard method was used for quantification and concentrations of target compounds were recovery-corrected. In 2012, the OCRL/Letcher Laboratories participated in the first international NORMAN PFR (OPFR) ILS and our OPFR results for this QA/QC exercise complied within 10% of the NORMAN PFR certified data.

A pork liver homogenate (Ottawa market), previously determined to be absent of any OPFRs, was fortified with each target OPFR and analyzed with the batch of samples ($n = 12$, liver or kidney). On the basis of fortified pork liver, the estimated method limit of quantification (MLOQ) and detection (MLOD) of the OPFRs was defined as the minimum amount of analyte that produced a peak with a calculated signal-to-noise ratio of 10 and 3, respectively. As a result, the MLOQ and MLOD (ng/g ww) for each OPFR are TBOEP (0.27, 0.09), TCEP (0.1, 0.03), TCIPP (0.31, 0.1), and TDCIPP (0.14, 0.04), respectively.

The method blank consisted of internal standard d_{27} -TBP spiked diatomaceous earth (DE) that was spiked with 200 μL chicken egg albumin (also previously determined to be absent of any OPFRs and used to maintain desirable IS recovery through volatility) and was used throughout the entire extraction, concentration, and analysis procedure. This was conducted in triplicate with the batch of 12 samples to demonstrate freedom from cross-contamination and contaminants that would interfere with analysis and to determine the levels of contamination associated with the processing and analysis of samples. The method blank concentration ranges for each of the four OPFRs were as follows: 1.7–4.1 ng/g ww for TCEP, 2.6–6.5 ng/g ww for TCIPP, 0.9–1.5 ng/g ww for TDCIPP, and 1.4–5.5 ng/g ww for TBOEP. OPFR blank subtraction was performed as follows: the highest level of contamination determined in the method blank triplicate was selected for sample correction, as well as the subtraction of one standard deviation of the three blanks.

Statistical Methods. End points were compared for each treatment to vehicle-treated controls by parametric analysis of variance (ANOVA) (e.g., hepatic T_4 -ORD), repeated measures (RM) ANOVA (i.e., thyroid hormones), or nonparametric ANOVA and RM ANOVA on ranked data (i.e., clinical

chemistry) using SAS 9.3. Residuals were tested for homogeneity of variance and normality. Only the concentrations of circulating thyroid hormones were successfully log transformed. Because each histological parameter of the thyroid gland required multiple measurements per bird, nested ANOVAs on ranked data were performed with treatment as the factor and the individual bird as the nested factor; Scheffé's *posthoc* tests followed. Hepatic T₄-ORD was examined with an ANOVA followed by least-squares means *posthoc* *t*-tests of all pairwise comparisons without adjustments. RM ANOVAs initially identified overall significant treatment effects on thyroid hormones or clinical chemistry for each OPFR compared to controls, followed by 1-way ANOVA to identify significant treatment differences at individual time periods when overall significance was identified. Spearman's correlation analysis was used to determine statistical associations between hepatic lipid and TBOEP concentrations and plasma concentrations of albumin and thyroid hormones. Means \pm SEM are presented and statistical significance occurred when $p \leq 0.05$. Only significant comparisons relative to the controls are reported unless noted.

RESULTS AND DISCUSSION

Kestrel Exposure and Uptake of OPFRs. After method blank-correction, hepatic concentrations of TCEP, TCIPP, and TDCIPP and all renal concentrations of the four OPFRs were less than MLOD (≤ 0.1 ng/g ww). Hepatic concentrations of TBOEP were quantifiable in TBOEP-exposed (5.0 ± 0.6 ; 3.2 – 7.9 ng/g ww) and control birds (4.0 ± 0.9 ; 0.3 – 9.4 ng/g ww), suggesting potential cross-contamination of the latter, but were not associated with lipid concentrations ($p = 0.10$). Although these results suggest the potential for the OPFRs failing to be absorbed by the kestrels during digestion, the same dietary exposure methodology used in the current study had previously resulted in kestrels accumulating PBDEs,^{19,23} HBCDD,²⁰ and BDE-209²¹ in their tissues or eggs. Furthermore, hepatic concentrations of these and other OPFRs were not detected in maternal herring gulls,² suggesting that OPFRs do not accumulate in hepatic tissues of birds. In a previous study, concentrations of TBOEP and TEP were negligible in pipped chickens exposed by egg injection,¹⁷ but TCIPP and TDCIPP increasingly accumulated in various tissues (e.g., liver) in a dose-dependent manner in pipped chickens exposed to much higher dosing concentrations¹⁶ than used in the current study.

The lack of detectable OPFR concentrations in the current kestrels is consistent, although not directly comparable, with the undetectable concentrations of these OPFRs in the tissues of some wild birds, including the eggs of common terns,³⁵ and the liver and brain of adult herring gulls.² However, low ppb concentrations of several OPFRs were reported in the plasma¹³ and eggs of herring gulls from the Great Lakes.^{2,12,34} Concurrently, nestling peregrine falcons (*Falco peregrinus*) had low plasma concentrations of TBOEP (1.0 – 7.5 ng/g ww), TCEP (0.2 – 2.0 ng/g ww), TCIPP (0.9 – 5.5 ng/g ww), and/or TDCIPP (0.3 – 1.0 ng/g ww) (Fernie et al., Environment Canada, unpublished data). Similar low ppb concentrations of these and other OPFRs were found in the feathers of white-tailed eagles¹⁴ and livers of kittiwakes (*Rissa tridactyla*) and common eiders (*Somateria mollissima*).¹¹ Together, the results of these studies suggest that accumulation of chlorinated and nonhalogenated OPFRs occurs in (wild) birds, presumably exposed continuously for longer periods or to higher

concentrations than experienced by the captive kestrels in the current study.

The undetectable residues of TCEP, TCIPP, TDCIPP, and to a lesser extent TBOEP, in the captive kestrels likely reflects rapid metabolism through enzyme-mediated processes. OP diesters, including BDCIPP and BCIPP derived from TDCIPP and TCIPP, respectively, were recently reported in humans,³⁶ suggesting the potential for *in vivo* metabolism of the four OPFRs used in the current study. In an *in vitro* rat liver microsomal assay, more than 95% of TDCIPP was rapidly metabolized (<5 min) to BDCIPP.³⁷ Similarly, TPHP was metabolized *in vitro* by CEH and mainly to the glucuronide conjugate of *para*-OH-TPHP.²⁷ Genes associated with xenobiotic phase I and II metabolism and lipid regulation, specifically CYP3A37, CYP2H1, and UGT1A9, were up-regulated in CEH and chicken neuronal cells exposed to TDCIPP or TCIPP.¹⁵ CYP3A37 and CYP2H1 are phase I cytochrome P450 metabolizing enzymes involved in oxidation of lipophilic compounds, including xenobiotics, and UGT1A9 is a phase II conjugation enzyme that joins small lipophilic molecules with larger polar glucuronic acids into excretable metabolites.^{15,18} These *in vitro* results were substantiated *in vivo* with pipped chicks that exhibited induction of hepatic CYP3A37 mRNA and type I deiodinase by TCIPP, CYP3A37, and CYP2H1 by TDCIPP¹⁶ and hepatic CYP3A37 mRNA by TBOEP.¹⁷

Biological Effects. The exposure to TBOEP, TCEP, TCIPP, or TDCIPP had no significant effect on the body mass or temporal weight gain of the American kestrels (p -values ≥ 0.22) and reflect similar results when adult male kestrels were exposed to DE-71^{19,38} but not a technical mixture of HBCDD (HBCDD-TM).³² However, the current OPFR-kestrel findings contrast with the reduced body mass of pipped chickens following *in vivo* developmental exposure to a much higher dose of TBOEP ($45,400$ ng/g).¹⁷

Traditional clinical chemistry measures of avian organ function are sensitive but nonspecific indicators of the effects of various persistent organic pollutants.^{39,40} Increases in circulating A:G ratios were reportedly associated with increasing plasma concentrations of Σ PBDE and Σ OH-PBDE metabolites in wild peregrine falcon nestlings.⁴⁰ The kestrels exposed to TCIPP or TDCIPP demonstrated significant overall changes in plasma A:G ratios (ranked RM ANOVA: $p \leq 0.01$) with significantly suppressed ratios at 21 d exposure to TDCIPP (ANOVA: $F_{1,16} = 5.71$, $p = 0.03$). Exposure to TBOEP had a significant overall effect on plasma albumin (RM ANOVA: $F_{3,13} = 6.95$, $p = 0.005$) with increased concentrations evident at 7 and 21 d exposure (ranked ANOVAs: $p \leq 0.05$). Changes in circulating ALP and ALT in some raptor species,^{39,40} but not all,⁴¹ have been associated with mirex, chlordanes, or PBDE concentrations in their tissues and were significantly affected in the current kestrels exposed to TCEP, TBOEP, or TDCIPP (ALT only) (ranked RM ANOVA: $p \leq 0.02$). Although increased plasma bile acids,¹⁷ liver,¹⁸ and gall bladder¹⁶ size were reported in pipped chickens following exposure to TEP and TCP, there were no significant effects on bile acids or any somatic organ indices of the adult kestrels exposed to TBOEP, TCEP, TCIPP, or TDCIPP. Although significant, the changes in A:G, albumin, ALP, and ALT in the kestrels were generally within the reported normal range of other raptors,^{40–43} albeit not kestrels, and thus may represent marginal changes in hepatocyte function and integrity from the OPFR exposure conditions in the current study.

OPFRs were characterized as having similar or more potent *in vitro* neurotoxicity as known or suspected neurotoxins.⁴⁴ Exposure of PC12 cells to TDCIPP promoted differentiation into dopaminergic and cholinergic neuronal phenotypes, and TCEP and TCIPP promoted the cholinergic phenotype.⁴⁴ In another study, TCIPP and TDCIPP were sequestered in the cerebral hemisphere of chicks, but there was no assessment of potential neurotoxic effects.¹⁶ A range of cholinergic markers were studied in both brain (cerebrum) and plasma of the kestrels (Table S1, SI). In the pilot study, the exposure to TDCIPP resulted in a statistically significant but small (5%) suppression of cerebrum cholinesterase activity (see SI) that was likely to be of little biological relevance. Yet the kestrels' exposure to the higher concentration of TDCIPP, or to TBOEP, TCEP, or TCIPP at the concentrations used in the current study, did not elicit significant changes in AChE or BChE enzyme activity in plasma or brain, or nicotinic or muscarinic cholinergic receptor levels in the brain (ANOVAs: all $p \geq 0.13$) (Table S1, SI).

Multiple indicators of oxidative status have been altered in response to *in vitro* and *in vivo* exposures to various flame retardants. The *in vitro* exposure of PC12 cells to TDCIPP increased lipid peroxidation, but exposure to TCIPP or TCEP had no effect on this indicator.⁴⁴ Exposure to PBDE congeners has elicited changes in oxidative status in raptors⁴⁵ but not passerines.^{46,47} Hepatic glutathione ratio and lipid peroxidation were induced in juvenile American kestrels exposed embryonically and as nestlings to a mixture of BDE-47, -99, -100, and -154.⁴⁵ Although a similar antioxidant-rich diet involving cockerel was fed to the adult male kestrels in the current study, the exposure to TBOEP, TCEP, TCIPP, or TDCIPP did not alter glutathione status or indicators of hepatic oxidative status (all $p \geq 0.11$).

Thyroid Function. Exposure to some OPFRs has elicited endocrine changes in fish,⁴⁸ and notably altered thyroid hormones in fish⁴⁹ and pipped chickens,^{16,17} with modeling demonstrating an association between TDCIPP exposure and FT_4 in humans.⁵⁰ Generally, circulating thyroid hormone concentrations are closely regulated in vertebrates through feedback mechanisms involving the hypothalamus, pituitary and thyroid glands, and peripheral tissues. Circulating TT_3 and TT_4 are bound to transthyretin (TTR) and albumin with varying importance depending on the class of vertebrate being considered. However, biological effects are generally elicited by circulating unbound FT_3 (predominantly) and FT_4 , and these unbound THs are also thought to exert the greatest negative feedback influence on the thyroid gland.

The exposure of the adult kestrels to each of the four OPFRs had significant overall effects on circulating FT_3 and to TBOEP on plasma TT_3 concentrations (all RM ANOVA $p \leq 0.05$). Circulating FT_4 concentrations (TBOEP, TCEP) and TT_4 (TDCIPP) in the adult kestrels were also significantly altered (all RM ANOVA $p \leq 0.05$), as were temporal changes in FT_4 and FT_3 plasma concentrations (see SI). Consistent significant increases in the circulating thyroid hormones were observed specifically at 7 d exposure to each OPFR (Figure 1a and b and Figure S1a and b in the SI). At 7 d, the exposure to TBOEP significantly increased the FT_3 concentrations by 32% ($F_{1,15} = 5.41$, $p = 0.03$) and TT_3 concentrations by 96% ($F_{1,15} = 20.23$, $p = 0.0005$) (Figure 1a and Figure S1a in the SI), whereas the TCEP treatment significantly increased the concentrations of FT_3 (57%) ($F_{1,16} = 19.27$, $p = 0.0005$) (Figure 1a) and FT_4 (96%) ($F_{1,14} = 7.42$, $p = 0.02$) (Figure 1b). At 7 d exposure to




Figure 1. Circulating concentrations of (a) FT_3 and (b) FT_4 thyroid hormones at 7 d in adult male American kestrels exposed daily to TBOEP, TCEP, TCIPP, or TDCIPP (22 ng OPFR/g kestrel/d) for 21 d. All comparisons are made to the adult male control kestrels. Concentrations of TT_3 and TT_4 are reported in Figure S1 a and b in the SI. Means \pm standard errors about the means (SEMs) are presented. Sample sizes for each thyroid hormone measured consisted of 10 control kestrels and 7 kestrels per OPFR treatment group, with the exception of the TBOEP ($n = 6$) and TDCIPP ($n = 6$) treatment groups, when assessing FT_4 concentrations. Statistically significant p -values are indicated as follows: $p \leq 0.0001$ (***), $p \leq 0.01$ (**), and $p \leq 0.05$ (*).

TDCIPP, FT_3 (32%) (Figure 1a) and TT_4 (31%) (Figure S1b, SI) concentrations were significantly increased; in comparison, the limited suppression of TT_4 (13%) at 21 d may be of little biological importance (all $p \leq 0.05$).

The changes reported here in the circulating thyroid hormones of the adult kestrels contribute to our understanding of the potential and varying effects of OPFRs^{16–18} and other FRs^{32,45,46,51} on thyroid function in birds. In pipped chickens, developmental exposure to TDCIPP or TEP, but not TBOEP or TCIPP, modulated genes associated with the thyroid hormone pathway and decreased plasma FT_4 but not FT_3 levels or glandular TT_4 concentrations.^{16,17} However, FT_4 and glandular TT_4 were unaltered following exposure of chicken embryos to tris(methylphenyl) phosphate (TMPP; formerly TCP).¹⁸ The reported changes in circulating thyroid hormones in several bird species exposed to various OPFRs builds on existing knowledge of thyroid sensitivity to multiple FRs, including HBCD,³² and PBDEs in some^{45,46} but not all^{38,51} avian studies.

Changes in circulating thyroid hormones may reflect changes in hormone transport to target tissues by transthyretin (TTR) and/or albumin. Although TTR was downregulated following in vitro exposure of CEH to TCIPP but not TDCIPP,¹⁵ it was unaltered in pipped chickens exposed in vivo to either OPFR.¹⁶ In the current kestrels, circulating levels of albumin were significantly and positively correlated with FT₃, FT₄, and TT₄ at 7 d and with TT₄ and TT₃ at 14 and 21 d ($n = 38$, Spearman's R -values ≥ 0.42 , $p \leq 0.01$). Significant changes, mostly increases, in the concentrations of FT₃, FT₄, and TT₄, but generally not in albumin, were evident at these time points, suggesting complex changes and other mechanisms were involved in eliciting effects on circulating thyroid hormones in the kestrels.

Thyroid hormonal changes can be transient either temporally or in a dose-dependent manner under the influence of contaminant stress. Changes in thyroid gland structure and circulating hormone concentrations, as well as deiodinase enzymes, are sensitive indicators of overall changes in thyroid function and gland activation. In pipped chickens, glandular T₄ was unaltered following TBOEP or TEP exposure.¹⁷ Histological assessments of the thyroid glands of the current adult kestrels revealed that exposure to the four OPFRs significantly altered ECH:colloid (Figure 2a) (nested nonparametric ANOVA: $F_{30,699} = 9.50$, $p < 0.0001$) as well as ECH and colloid size (Figure S2a and b, SI) compared to controls. The ECH:colloid ratio, an integrated measure of thyroid gland activity,⁵² was significantly increased in the kestrels exposed to TCEP, TCIPP, or TDCIPP for 21 d (*posthoc* $p \leq 0.008$) (Figure 2a), suggesting reduced activity of their thyroid glands.⁵² Compared to the controls, TDCIPP exposure significantly reduced ECH and increased colloid area (*posthoc* $p \leq 0.004$). TCEP exposure significantly reduced ECH (*posthoc* $p < 0.0001$) (Figure S2a and b, SI).

Thyroid hormone activation (i.e., conversion from T₄ to T₃) and degradation (conversion to reverse T₃ or T₂ and further deiodinated products) are mediated by various deiodinase enzymes primarily in the liver, but also in the kidney, small intestine, and brain. In vitro changes were evident in deiodinase genes from chickens following embryonic exposure to TCIPP but not TDCIPP, whereby Type I deiodinase enzymes were induced in hepatocytes.¹⁶ In the current in vivo study with adult kestrels, exposure to the four OPFRs significantly affected hepatic T₄-ORD activity (ANOVA: $F_{4,34} = 3.24$, $p = 0.025$) (Figure 2b) with exposure to TBOEP, TCIPP, and TDCIPP significantly increasing hepatic T₄-ORD activity involved in the conversion of T₄ to T₃ (*posthoc* $p \leq 0.04$). Although activity of other deiodinase enzymes was not assessed, the pattern of the results suggests that the OPFR exposure altered thyroid function of the kestrels in vivo. We hypothesize that the increased hepatic T₄-ORD activity in combination with the decreased gland activity was potentially sufficient to maintain circulating thyroid hormone concentrations at 21 d of exposure. Additional research is required to test this hypothesis.

In the current study, the exposure of captive kestrels to low concentrations of TBOEP, TCEP, TCIPP, or TDCIPP is likely similar to or lower than that encountered by wild birds^{2,13,34} because no measurable concentrations were detected in the kestrels. Some clinical chemistry measures suggested moderate alterations in hepatocyte integrity and function. Exposure to each of the four OPFRs elicited overall changes in circulating thyroid hormones, particularly in FT₃, with relative increases of 32–96% at 7 d of exposure; thereafter, the only change in




Figure 2. Significant changes in (a) thyroid gland histology (ECH:colloid perimeter) and (b) hepatic T₄-outer ring deiodinase (T₄-ORD) were observed in adult male kestrels at 21 d exposure to TBOEP, TCEP, TCIPP, or TDCIPP when compared to control kestrels. The ECH:colloid ratio represents an integrated measure of thyroid gland activity (Bocian-Sobkowwska et al. 1997). Significant changes in epithelial cell height and colloid area are reported in Figure S2a and b in the SI. Means \pm SEMs are presented. Sample sizes consist of 7 birds per treatment group, including controls. Statistically significant p -values are indicated as follows: $p \leq 0.0001$ (***), $p \leq 0.01$ (**), and $p \leq 0.05$ (*).

circulating thyroid hormone concentrations was minor, and the thyroid hormones were likely maintained during this time through reduced activity of the thyroid gland and/or the increased activity of hepatic T₄-ORD enzymes. The results of the current in vivo study with adult kestrels and the previous studies with pipped chickens^{16,17} demonstrate that exposure to these particular OPFRs has the potential to elicit effects in both young and adult birds. Further research is required, however, to determine potential endocrine and physiological effects of OPFRs on wild birds given their chronic, longer-term exposure to OPFRs compared to that experienced by the American kestrels in the current study.

■ ASSOCIATED CONTENT

● Supporting Information

This section describes the pilot study conducted with TDCIPP, and the detailed methodology and results, appropriate figures, and table of neurochemistry and clinical chemistry values for American kestrels from the subsequent study involving the exposure of the captive birds to TBOEP, TCEP, TCIPP, or TDCIPP. The Supporting Information is available free of

charge on the ACS Publications website at DOI: 10.1021/acs.est.5b00857.

AUTHOR INFORMATION

Corresponding Author

*Phone: +1-905-336-4843; email: kim.fernie@ec.gc.ca.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank Environment Canada (Chemicals Management Plan, Ecotoxicology and Wildlife Health Division) and the United States Geological Survey; Dr. Carolyn Cray (U. Miami); Dr. John French, Wayne Bauer, Mary Maxey (all USGS Patuxent Wildlife Research Center). Thanks also go to Lewis Gauthier in the OCRL/Letcher Laboratories at Environment Canada (NWRC, Ottawa, ON, Canada) for the OPFR analysis of the kidney and liver tissues and to Dr. Jennifer Rutiewicz for performing the neurochemical biomarker work. Funding for this project was provided by the Chemicals Management Plan and the Ecotoxicology and Wildlife Health Division of Environment Canada. The authors declare no competing financial interest. We thank Pamela Martin (Environment Canada) and Dr. Miguel Moira (USGS) for their reviews of an earlier draft of this manuscript.

REFERENCES

- (1) Van der Veen, L.; de Boer, J. Phosphorous flame retardants: properties, production, environmental occurrence, toxicity and analysis. *Chemosphere* **2012**, *88*, 1119–1153.
- (2) Greaves, A.; Letcher, R. J. Comparative body compartment composition and *in ovo* transfer of organophosphate flame retardants in North American Great Lakes herring gulls. *Environ. Sci. Technol.* **2014**, *48*, 7942–7950.
- (3) United States Environmental Protection Agency. 2012. Chemical Data Access Tool (CDAT). Available from: http://java.epa.gov/oppt_chemical_search (accessed 14 April 2015).
- (4) Substances in Preparations in Nordic Countries [SPIN] (database on the Internet). 2013. Copenhagen (DK): Nordic Council of Ministers [cited 2013 October]. Available from: <http://195.215.202.233/DotNetNuke/default.aspx>.
- (5) Stapleton, H. M.; Kosterhaus, S.; Eagle, S.; Fuh, J.; Meeker, J. D.; Blum, A.; Webster, T. F. Detection of organophosphate flame retardants in furniture foam and U.S. house dust. *Environ. Sci. Technol.* **2009**, *43*, 7490–7495.
- (6) Sundkvist, A. M.; Olofsson, U.; Haglund, P. Organophosphorus flame retardants and plasticizers in marine and fresh water biota and in human milk. *J. Environ. Monit.* **2010**, *12*, 943–951.
- (7) Salamova, A.; Ma, Y.; Venier, M.; Hites, R. High levels of organophosphate flame retardants in the Great Lakes atmosphere. *Environ. Sci. Technol.* **2014**, *1*, 8–14.
- (8) Möller, A.; Xie, Z.; Caba, A.; Sturm, R.; Ebinghaus, R. Organophosphate flame retardants and plasticizers in the atmosphere of the North Sea. *Environ. Pollut.* **2011**, *159*, 3660–3665.
- (9) Möller, A.; Sturm, R.; Xie, Z.; Cai, M.; He, J.; Ebinghaus, R. Organophosphate flame retardants and plasticizers in airborne particles over the Northern Pacific and Indian Ocean toward the polar regions: Evidence for global occurrence. *Environ. Sci. Technol.* **2012**, *46*, 3127–3134.
- (10) Leonards, P.; Stendal, E. H.; van der Veen, I.; Berg, V.; Bustnes, J. O.; van Leeuwen, S. Screening of organophosphorous flame retardants 2010. SPFO-Report 1091/2011. TA-2786/2011.
- (11) Evensen, A.; Leknes, H.; Christensen, G. N.; Warner, N.; Remberger, M.; Gabrielsen, G. W. Screening of new contaminants in samples from Norwegian Arctic. NIVA Report 4351-1. SPFO-Report 1049/2009. TA-2510/2009978-82-449-0065-2, 2009.
- (12) Letcher, R. J.; Chen, D.; Chu, S. G.; Gauthier, L. T. Organophosphate triester flame retardant and phosphoric acid diester metabolite analysis and spatial and temporal trends in herring gulls from the North American Great Lakes. *Organohalogen Compd.* **2011**, *73*, 1257–1260.
- (13) Su, G.; Greaves, A. K.; Gauthier, L. T.; Letcher, R. J. Liquid chromatography-electrospray-tandem mass spectrometry method for determination of organophosphate diesters in biotic samples including Great Lakes herring gull plasma. *J. Chromatogr. A* **2014**, *1374*, 85–92.
- (14) Eulaers, I.; Jaspers, V. L. B.; Halley, D. J.; Lepoint, G.; Nygård; Pinxten, R.; Covaci, A.; Eens, M. Brominated and phosphorous flame retardants in white-tailed eagle *Haliaeetus albicilla* nestlings: Bioaccumulation and association with dietary proxies ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\delta^{34}\text{S}$). *Sci. Total Environ.* **2014**, *478*, 48–57.
- (15) Crump, D.; Chiu, S.; Kennedy, S. W. Effects of tris(1,3-dichloro-2-propyl) phosphate and tris(1-chloropropyl) phosphate on cytotoxicity and mRNA expression in primary cultures of avian hepatocytes and neuronal cells. *Toxicol. Sci.* **2012**, *126*, 140–148.
- (16) Farhat, A.; Crump, D.; Chiu, S.; Williams, K. L.; Letcher, R. J.; Gauthier, L. T.; Kennedy, S. W. *in ovo* effects of two organophosphate flame retardants—TCPP and TDCPP—on pipping success, development, mRNA expression, and thyroid hormone levels in chicken embryos. *Toxicol. Sci.* **2013**, *134*, 92–102.
- (17) Egloff, C.; Crump, D.; Porter, E.; Williams, K. L.; Letcher, R. J.; Gauthier, L. T.; Kennedy, S. W. Tris(2-butoxyethyl)phosphate and triethyl phosphate alter embryonic development, hepatic mRNA expression, thyroid hormone levels, and circulating bile acid concentrations in chicken embryos. *Toxicol. Appl. Pharmacol.* **2014**, *279*, 303–310.
- (18) Crump, D.; Porter, E.; Egloff, C.; Williams, K. L.; Letcher, R. J.; Gauthier, L. T.; Kennedy, S. W. 1,2-Dibromo-4-(1,2-dibromoethyl)-cyclohexane and tris(methylphenyl) phosphate cause significant effects on development, mRNA expression, and circulating bile acid concentrations in chicken embryos. *Toxicol. Appl. Pharmacol.* **2014**, *277*, 279–287.
- (19) Fernie, K. J.; Shutt, J. L.; Letcher, R. J.; Ritchie, J. I.; Sullivan, K.; Bird, D. M. Environmentally relevant concentrations of DE-71 and HBCD alter eggshell thickness and reproductive success of American kestrels. *Environ. Sci. Technol.* **2009**, *43*, 2124–2130.
- (20) Fernie, K. J.; Martenson, S. C.; Bird, D. M.; Ritchie, I. J.; Letcher, R. J. Reproductive changes in American kestrels (*Falco sparverius*) in relation to exposure to technical hexabromocyclododecane flame retardant. *Environ. Toxicol. Chem.* **2011**, *30*, 2570–2575.
- (21) Letcher, R. J.; Martenson, S. C.; Fernie, K. J. Dietary exposure of male American kestrels (*Falco Sparverius*) to decabromodiphenyl ether (BDE-209) flame retardant: Uptake, distribution, debromination and enzyme induction. *Environ. Int.* **2014**, *63*, 182–190.
- (22) Martenson, S. C.; Letcher, R. J.; Graham, L.; Kimmins, S.; Tomy, G.; Palace, V. P.; Ritchie, I. J.; Gauthier, L.; Bird, D. M.; Fernie, K. J. The Flame Retardant β -1,2-Dibromo-4-(1,2-dibromoethyl)-cyclohexane (β -TBECH): Fate, Fertility and Reproductive Success in American Kestrels (*Falco sparverius*). *Environ. Sci. Technol.* **2012**, *46*, 8440–8447.
- (23) Sullivan, K. M.; Martenson, S. C.; Letcher, R. J.; Bird, D. M.; Ritchie, I. J.; Shutt, J. L.; Fernie, K. J. Changes in the incubation by American kestrels (*Falco sparverius*) during exposure to the polybrominated diphenyl ether (PBDE) mixture DE-71. *J. Toxicol. Environ. Health, Part A* **2013**, *76*, 978–989.
- (24) McKernan, M. A.; Rattner, B. A.; Hale, R. C.; Ottinger, M. A. Toxicity of polybrominated diphenyl ethers (DE-71) in chicken (*Gallus gallus*), mallard (*Anas platyrhynchos*), and American kestrel (*Falco sparverius*) embryos and hatchlings. *Environ. Toxicol. Chem.* **2009**, *28*, 1007–17.
- (25) Rattner, B. A.; Lazarus, R. S.; Heinz, G. H.; Hale, R. C.; Karouna-Renier, N. K.; Schultz, S. L. Apparent tolerance of common tern (*Sterna hirundo*) embryos to a pentabrominated diphenyl ether mixture (DE-71). *Chemosphere* **2012**, *93*, 441–447.
- (26) Porter, R. D.; Wiemeyer, S. N. Propagation of American kestrels. *J. Wildl. Manage.* **1970**, *34*, 594–604.

- (27) Su, G.; Crump, D.; Letcher, R. J.; Kennedy, K. W. Rapid in vitro metabolism of the flame retardant triphenyl phosphate and effects on cytotoxicity and mRNA expression in chicken embryonic hepatocytes. *Environ. Sci. Technol.* **2014**, *48*, 13511–13519.
- (28) Tang, F.; Messinger, S.; Cray, C. Use of an indirect sampling method to produce reference intervals for hematologic and biochemical analyses in *Psittaciform* species. *Journal of Avian Medicine and Surgery* **2013**, *27*, 194–203.
- (29) Basu, N.; Scheuhammer, A. M.; Sonne, C.; Letcher, R. J.; Born, E. W.; Dietz, R. Is dietary mercury of neurotoxicological concern to polar bears (*Ursus maritimus*)? *Environ. Toxicol. Chem.* **2009**, *28*, 133–140.
- (30) Rutkiewicz, J.; Crump, D.; Scheuhammer, A. M.; Jagla, M.; Basu, N. Investigation of spatial trends and neurochemical impacts of mercury in herring gulls across the Laurentian Great Lakes. *Environ. Pollut.* **2010**, *158*, 2733–2737.
- (31) Rutkiewicz, J.; Bradley, M.; Mittal, K.; Basu, N. Methylmercury egg injections: Part 2-Pathology, neurochemistry, and behavior in the avian embryo and hatchling. *Ecotoxicol. Environ. Saf.* **2013**, *93*, 77–86.
- (32) Marteinson, S. C.; Kimmins, S.; Letcher, R. J.; Palace, V. P.; Bird, D. M.; Ritchie, I. J.; Fernie, K. J. Diet exposure to technical hexabromocyclododecane (HBCD) affects testes and circulating testosterone and thyroxine levels in American kestrels (*Falco sparverius*). *Environ. Res.* **2011**, *111*, 1116–1123.
- (33) Park, B. J.; Palace, V. P.; Wautier, K.; Gemmill, B.; Tomy, G. Thyroid axis disruption in juvenile Brown trout (*Salmo trutta*) exposed to the flame retardant β -tetrabromocyclohexane (β -TBECH) via the diet. *Environ. Sci. Technol.* **2011**, *45*, 7923–7927.
- (34) Chen, D.; Letcher, R. J.; Chu, S. G. Determination of non-halogenated, chlorinated and brominated organophosphate flame retardants in herring gull eggs based on liquid chromatography-tandem quadrupole mass spectrometry. *J. Chromatogr. A* **2012**, *1220*, 169–174.
- (35) Brandsma, S. H.; Leonards, P. E. G.; Leslie, H. A.; de Boer, J. Tracing organophosphorous and brominated flame retardants and plasticizers in an estuarine food web. *Sci. Total Environ.* **2015**, *505*, 22–31.
- (36) van den Eede, N.; Heffernan, A. L.; Aylward, L. L.; Hobson, P.; Neels, H.; Mueller, J. F.; Covaci, A. Age as a determinant of phosphate flame retardant exposure of the Australian population and identification of novel urinary PFR metabolites. *Environ. Int.* **2015**, *74*, 1–8.
- (37) Chu, S. G.; Chen, D.; Letcher, R. J. Dicationic ion-pairing of phosphoric acid diesters post-liquid chromatography and subsequent determination by electrospray positive ionization-tandem mass spectrometry. *J. Chromatogr. A* **2011**, *1218*, 8083–8088.
- (38) Marteinson, S. C.; Kimmins, S.; Bird, D. M.; Shutt, J. L.; Letcher, R. J.; Ritchie, I. J.; Fernie, K. J. Embryonic exposure to the polybrominated diphenyl ether mixture, DE-71, affects testes and circulating testosterone concentrations in adult American kestrels (*Falco sparverius*). *Toxicol. Sci.* **2011**, *121*, 168–176.
- (39) Sonne, C.; Bustnes, J. O.; Herzke, D.; Jaspers, V. L. B.; Covaci, A.; Halley, D. J.; Moum, T.; Eulaers, I.; Eens, M.; Ims, R. A.; Hanssen, S. A.; Erikstad, K. E.; Johnsen, T. V.; Schnug, L.; Rigét, F. F.; Jensen, A. L. Relationships between organohalogen contaminants and blood plasma clinical-chemistry parameters in chicks of three raptor species from northern Norway. *Ecotoxicol. Environ. Saf.* **2010**, *73*, 7–17.
- (40) Sonne, C.; Bustnes, J. O.; Herzke, D.; Jaspers, V. L. B.; Covaci, A.; Eulaers, I.; Halley, D. J.; Moum, T.; Ballesteros, M.; Eens, M.; Ims, R. A.; Hanssen, S. A.; Erikstad, K. E.; Johnsen, T. V.; Rigét, F. F.; Jensen, A. L.; Kjelgaard-Hansen, M. Blood plasma clinical-chemistry parameters as biomarker endpoints for organohalogen contaminant exposure in Norwegian raptor nestlings. *Ecotoxicol. Environ. Saf.* **2012**, *80*, 76–83.
- (41) Smits, J. E. G.; Fernie, K. J. Avian Wildlife as Sentinels of Ecosystem Health. *CIMID* **2013**, *36*, 333–342.
- (42) Lumeij, J. T.; Remple, J. D. Plasma urea, creatinine and uric acid concentrations in relation to feeding in peregrine falcons (*Falco peregrinus*). *Avian Pathology* **1991**, *20*, 79–83.
- (43) Lanzarot, M.; Montesinos, A.; San Andrés, M. I.; Rodriguez, C.; Barahona, M. V. Hematological, protein electrophoresis and cholinesterase values of free-living nestling peregrine falcons in Spain. *J. Wildl. Dis.* **2001**, *37*, 172–177.
- (44) Dishaw, L. V.; Powers, C. M.; Ryde, I. T.; Roberts, S. C.; Seidler, F. J.; Slotkin, T. A.; Stapleton, H. M. Is the PentaBDE replacement, tris(1,3-dichloro-2-propyl)phosphate (TDCPP), a developmental neurotoxicant? Studies in PC12 cells. *Toxicol. Appl. Pharmacol.* **2011**, *256*, 281–289.
- (45) Fernie, K. J.; Shutt, L.; Mayne, G.; Hoffman, D.; Letcher, R. J.; Drouillard, K.; Ritchie, I. Exposure to polybrominated diphenyl ethers (PBDEs): changes in thyroid, vitamin A, glutathione homeostasis and oxidative stress in American kestrels (*Falco sparverius*). *Toxicol. Sci.* **2005**, *88*, 375–383.
- (46) Eng, M.; Elliott, J. E.; Williams, T. D. An assessment of the developmental toxicity of BDE-99 in the European starling using an integrated laboratory and field approach. *Ecotoxicology* **2014**, *23*, 1505–1516.
- (47) Eng, M.; Williams, T. D.; Elliott, J. E. Developmental exposure to a brominated flame retardant: an assessment of effects on physiology, growth, and reproduction in a songbird, the zebra finch. *Environ. Pollut.* **2013**, *178*, 343–349.
- (48) Liu, X.; Ji, K.; Jo, A.; Moon, H.-B.; Choi, K. Effects of TDCPP or TPP on gene transcriptions and hormones of HPG axis, and their consequences on reproduction in adult zebrafish (*Danio rerio*). *Aquat. Toxicol.* **2013**, *134–135*, 104–111.
- (49) Wang, Q.; Lian, K.; Liu, J.; Yang, L.; Guo, Y.; Liu, C.; Zhou, C. Exposure of zebrafish embryos/larvae to TDCPP alters concentrations of thyroid hormones and transcriptions of genes involved in the hypothalamic-pituitary-thyroid axis. *Aquat. Toxicol.* **2013**, *126*, 207–213.
- (50) Meeker, J. D.; Stapleton, H. M. House dust concentrations of organophosphate flame retardants in relation to hormone levels and semen quality parameters. *Environ. Health Perspect.* **2010**, *118*, 318–323.
- (51) Winter, V.; Williams, T. D.; Elliott, J. E. A three-generational study of in ovo exposure to PBDE-99 in the zebra finch. *Environ. Toxicol. Chem.* **2013**, *32*, 345–352.
- (52) Bocian-Sobkowska, J.; Wozniak, W.; Malendowicz, L. K. Morphometric studies on the development of the human thyroid gland. II. The late fetal life. *Histol. Histopathol.* **1997**, *12*, 79–84.

SUPPLEMENTARY INFORMATION: COVER SHEET

Investigating Endocrine and Physiological Parameters of Captive American kestrels Exposed by Diet to Selected Organophosphate Flame Retardants

Kimberly J. Fernie^{1*}; Vince Palace²; Lisa E. Peters²; Nil Basu³; Robert J. Letcher⁴; Natalie K. Karouna-Renier⁵; Sandra L. Schultz⁵; Rebecca S. Lazarus⁵; B. A. Rattner⁵.

¹ Ecotoxicology & Wildlife Health Division, Environment Canada, 867 Lakeshore Road, Burlington, ON, Canada L7R 4A6. +1-905-336-4843; kim.fernie@ec.gc.ca

² Stantec Consulting Ltd., 386 Broadway Avenue, Winnipeg, MB, Canada R3C 3R6

³ McGill University, 21,111 Lakeshore Road, Ste Anne de Bellevue, PQ, Canada H9X 3V9

⁴ Ecotoxicology & Wildlife Health Division, Environment Canada, Carleton University, 1125 Colonel By Drive, Ottawa, ON, Canada K1A 0H3

⁵ U.S. Geological Survey, Patuxent Wildlife Research Center, BARC East Building 308, 10300 Baltimore Avenue, Beltsville MD 20705, USA

Number of pages in SI (including this page): 17

Number of figures: 4 in total although labelled as Figs S1a, S1b, S2a, S2b

Number of tables: 1

SUPPLEMENTARY INFORMATION

Investigating Endocrine and Physiological Parameters of Captive American kestrels Exposed by Diet to Selected Organophosphate Flame Retardants

Fernie, Kimberly J^{1*}; Palace, Vince²; Peters, Lisa²; Basu, Nilandri³; Letcher, Robert J⁴; Karouna-
Renier, Natalie K⁵; Schultz, Sandra L⁵; Lazarus, Rebecca S⁵; Rattner, Barnett A⁵.

¹ Ecotoxicology & Wildlife Health Division, Environment Canada, 867 Lakeshore Road,
Burlington, ON, Canada L7R 4A6. +1-905-336-4843; kim.fern timer@ec.gc.ca

² Stantec Consulting Ltd., 386 Broadway Avenue, Winnipeg, MB, Canada R3C 3R6

³ McGill University, 21,111 Lakeshore Road, Ste Anne de Bellevue, PQ, Canada H9X 3V9

⁴ Ecotoxicology & Wildlife Health Division, Environment Canada, Carleton University, 1125
Colonel By Drive, Ottawa, ON, Canada K1A 0H3

⁵ U.S. Geological Survey, Patuxent Wildlife Research Center, BARC East Building 308, 10300
Baltimore Avenue, Beltsville MD 20705, USA

Introduction

In the Supplemental Information section, the methodology and results of a pilot study conducted with TDCIPP in 2011 are presented. In addition, the detailed methodology and results, including appropriate figures and table, from the main study conducted in 2012, are presented for the neurochemistry, oxidative status, thyroid function, and clinical chemistry assessments of the captive adult male American kestrels exposed to TBOEP, TCEP, TCIPP or TDCIPP.

Materials and Methods

Pilot Study

A pilot study was conducted in 2011 using 7 adult two-year old male American kestrels exposed daily by diet for 28 days to TDCIPP and 7 adult male kestrels of the same age exposed to only the safflower oil vehicle. The initial exposure dose in the pilot study was 226 ng TDCIPP /d; each kestrel ate one cockerel/d that had been injected with 50 μ L of a safflower oil solution containing TDCIPP at 4.52 ng/ μ L. Based on the mean weight (114 g) of the adult male kestrels at the initiation of the study, each kestrel received a daily dose of 1.98 ng TDCIPP/g kestrel/d. This dosing concentration was 10x less than the TDCIPP exposure concentration used in the 2012 main study. The kestrels were weighed and their general health assessed at the beginning of the exposure period, and again at 7, 14, 21, and 28 d of the pilot exposure trial. All housing, handling, blood sampling and dissecting procedures were the same as those used in the main study conducted in 2012, and are described in Fernie et al. (submitted with this SI section).

Neurochemical Assays

Neurochemical assays were completed for the 2011 pilot study at the National Wildlife Research Center (Environment Canada, Ottawa, ON, Canada). Brains were dissected from the

cranium at the time of necropsy, fixed in formalin and stored in a scintillation vial until processing. Neurochemical analysis was conducted at the National Wildlife Research Center (Environment Canada, Ottawa, ON) in 2011 and at the University of Michigan (Basu Laboratory) in 2012. In 2011, the whole brain was homogenized in buffer/Triton-X-100 at a ratio of 1 part brain and 3 part of 1% Triton solution to a final concentration of 250 mg/mL. The homogenate was centrifuged to eliminate interferences during spectrophotometric analysis and 10 μ L of the supernatant homogenate was analyzed in duplicate. Cholinesterase (ChE) analysis was completed as described in MET-BMK-CHE-01J, as modified by Hill and Fleming (1982). The cuvette contained 3 mL of DTNB solution (5,5'-dithiobis-nitrobenzoic acid in 2.5×10^{-4} M in phosphate buffer 0.05M, pH 7.9), 10 μ L of brain homogenate and 100 μ L of acetylthiocholine iodide 0.0312 M (for a final concentration of 1.0 mM). ChE activity was determined with a spectrophotometer (Hewlett Packard Diode Array HP8453, S/N 02502465) at 25°C. The change in absorbance at 405 nm over one minute was recorded, with readings every 15 seconds, and a wait time of 60 seconds to ensure readings were made during the linear segment of the reaction. ChE analysis was conducted in duplicate for each bird.

In 2012, neurochemical assays were completed at the University of Michigan (NB) based on previous analytical methods (Basu et al. 2009; Rutkiewicz et al. 2010, 2013). Cellular membranes were prepared from brain tissues using protocols described elsewhere (Basu et al., 2009). Binding to the muscarinic (mACh) and nicotinic nACh) receptors, was performed in Na/K buffer (50 mM NaH_2PO_4 , 5 mM KCl, 120 mM NaCl, pH 7.4). Thirty μ g of membrane preparation was re-suspended in buffer and added to microplate wells containing a 1.0 μ M GF/B glass filter (Millipore, Boston, MA, USA). For muscarinic and nicotinic receptor binding, samples were incubated with 1 nM [^3H]-QNB (42 Ci/mmol; NEN/Perkin Elmer, Boston, MA,

USA) and 1 nM [^3H]-Cytisine (26.8 C/mmol; NEN/Perkin Elmer), respectively, for 60 min. All assays were carried out with gentle shaking at room temperature and binding reactions were terminated by vacuum filtration. The filters were rinsed three times with buffer and then allowed to soak for 96 hrs in 25 μL of OptiPhase Supermix Cocktail (Perkin Elmer). Radioactivity retained by the filter was quantified by liquid scintillation counting in a microplate detector (Wallac Microbeta, Perkin Elmer) having a counting efficiency of approximately 50%. Specific binding to both receptors was defined as the difference in radioligand bound in the presence and absence of 100 μM unlabeled atropine. Binding was reported as fmol of radioisotope bound per mg of membrane protein (fmol/mg). All samples were assayed in quadruplicate for total and non-specific binding.

The enzymatic activities of cholinesterase (acetylcholinesterase in plasma and brain tissue, and butyrylcholinesterase in plasma) were completed according to published procedures with slight modifications. Tissues were sonicated for 30 s in cold Na/K buffer that included 0.5 % (v/v) Triton X-100. Following a 10 min centrifugation (15,000 $\times g$, 4 $^{\circ}\text{C}$) the supernatant was removed. For cholinesterase activity, 0.5 μg of supernatant protein was mixed with 100 μM 10-acetyl-3, 7-dihydroxyphenoxazine (Amplex Red, Molecular Probes), 200 mU horseradish peroxidase, 20 mU choline oxidase, and 100 μM acetylcholine. For acetylcholinesterase, iso-OMA was used to inhibit butyrylcholinesterase activity. For butyrylcholinesterase, galanthamine was used to inhibit acetylcholinesterase activity. Following a 30 min incubation period for both assays, the reaction end-product (resorufin: $\lambda_{\text{ex}} = 540$, $\lambda_{\text{em}} = 590$) was monitored between 30 and 60 min (CytoFluor 2350, Millipore). Specific activities were expressed as nmol of resorufin formed per min per unit (μg or mg) protein. Each sample was assayed in triplicate.

Oxidative Status

In 2012, a portion of each liver was analyzed for seven indicators of oxidative stress: reduced glutathione (GSH), oxidized glutathione (GSSG), GSSG:GSH, total glutathione (tGSH), total sulfhydryl (TSH), protein bound sulfhydryl (PBSH), and thiobarbituric acid reactive substances (TBARS). Frozen liver samples were thawed and homogenized on ice in 2X PBS (pH 7.4; Fisher BioReagents, Waltham, MA) at 200 µg/µl. The homogenate was centrifuged at 14,000 rpm for 10 minutes at 4°C and 60 µl aliquots of the supernatant were transferred to 0.2 ml tubes and frozen at -80°C until analysis. For GSH, GSSG, GSSG/GSH, and tGSH analysis, the liver supernatant was thawed and diluted to 12.5 µg/µl in 1X PBS and analyzed using the DetectX® Glutathione Fluorescent Detection Kit (Arbor Assays, Ann Arbor, MI) following the manufacturer's protocol. For determination of TBARS, the sample supernatant was thawed on ice and diluted to 100 µg/µl in 1X PBS. TBARS levels were determined using the QuantiChrom™ TBARS Assay Kit (Bioassay Systems, Hayward, CA) following the manufacturer's instructions. TSH was determined in thawed supernatant that had been diluted to 6.25 µg/µl in 1X PBS using the Measure-iT™ Thiol Assay Kit (Life Technologies - Molecular Probes, Inc., Eugene, OR) following the manufacturer's instructions. Concentrations of PBSH were calculated as the difference between TSH and GSH concentrations.

Thyroid Function: Deiodinase Activity and Histology of the Thyroid Gland

In 2012, a three-pronged assessment of thyroid function was performed (LP, VP) by determining circulating triiodothyronine (T₃) and T₄ concentrations, hepatic T₄-outer ring deiodinase activity (T₄-ORD), and histological assessments of the structure of the thyroid gland. For determination of deiodinase activity, a 100-200 mg sample of each liver was homogenized in ice cold 0.1 M sodium phosphate buffer (pH=7.4) containing 1 mM ethylenediaminetetraacetic acid (EDTA) and 20 mM dithiothriitol (DTT), centrifuged at 9,000 X g and 4 °C for 15 minutes,

and the resulting S9 supernatant recovered. The S9 homogenate was frozen and stored at -80°C until analysis. Protein concentration was determined in a separate aliquot of S9 homogenate using the Bradford (1976) method. To determine deiodinase activity, 10 μl of S9 homogenates were incubated at 37°C with 900 μl of sodium phosphate buffer. At timed intervals, 5 μl of dimethylsulfoxide containing 3000 ng of T_4 were added to each sample to begin the reactions. Reactions were terminated 90 minutes later by adding 1 ml of ice cold methanol. Two types of blanks were included with each analytical run: one with all assay constituents except S9 and another containing all constituents except T_4 . To determine T_3 produced by the reactions, 100 μl of the final reaction volume was used in commercially available radioimmunoassays (MP Biomedical, Santa Ana CA, USA). Deiodinase activity was expressed as pmol of T_3 liberated per minute per mg of protein, with blank correction included as previously described for American kestrels (Martinson et al. 2011).

Formalin-fixed tissues were processed and embedded in paraffin using standard procedures (Luna 1968). Tissues were sectioned at approximately 7 μm , mounted on slides, and stained with hematoxylin and eosin (H & E). Each histological slide was examined using a Carl Zeiss EMS stereoscope with an APOLumar SI.2Y objective, and an AxioCam HRC digital camera at 26 milliseconds (ms) exposure to capture images of the thyroid tissues for measurement. Digital images of two separate fields of view for each slide were taken at 40x magnification. For each thyroid gland, ten complete follicles per image (for a total of 20 follicles) were selected and analyzed as described by Park et al. (2011). Briefly, epithelial cell heights (ECH) were measured (μm) at 4 locations per follicle, approximately 90 degrees from each other, and then averaged to calculate a mean ECH per follicle. Area (μm^2) of the colloid inside the same 20 thyroid follicles was also measured. The data were expressed as mean

thyroid ECH and mean colloid area per individual bird. The mean ratio of colloid perimeter:ECH was also calculated as an indicator of thyroid gland activation and the potential for thyroid hormone production (Bocian-Sobkowwska et al. 1997). All analyses were performed using Zen Lite 2012 (2012) software.

Clinical Chemistry

The Ortho Vitros 250XR chemistry analyzer (Ortho, Rochester, NY, USA) was maintained by the recommended manufacturer and internal quality assurance programs. Standard QA/QC measures were run to assure analyzer performance. The following methods were used: AST- aspartate + alpha ketoglutarate method; ALT – alanine + alpha ketoglutarate method; LDH – pyruvate + NADH method; GTT – L-gammaglutamyl p-nitroanilide method; ALK PHOS – p-nitrophenyl phosphate method; Total protein – biuret method. GLDH was analyzed on the Randox Daytona analyzer (Kearneysville, WV, USA) and the reagents were also purchased from and implemented according to Randox guidelines. QA/QC for the GLDH involved a 2-tier control, and employed the alpha oxoglutarate + NADH method. Hemolysis, lipemia and bile acids were analyzed according to Cray and Andreopoulos (2003), and analysis of bile acids utilized the MP Biomedical radioimmunoassay kit. Electrophoresis was used to determine prealbumin, albumin, and globulins, and run using the Helena system according to Cray et al. (2011).

Statistical Methods

As described in the main study (Ferne et al., submitted), endpoints were compared for each treatment to vehicle-treated controls only, by parametric analysis of variance (ANOVAs) (e.g., hepatic T4-ORD) or repeated measures (RM) ANOVAs (i.e., thyroid hormones), or non-parametric ANOVAs and RM ANOVAs on ranked data (i.e., clinical chemistry), using SAS

9.3®. Residuals were tested for homogeneity of variance and normality. Only the concentrations of circulating thyroid hormones were successfully log transformed. Since each histological parameter of the thyroid gland required multiple measurements per bird, nested ANOVAs on ranked data were performed with treatment as the factor and the individual bird as the nested factor; Scheffe's *post-hoc* tests were then applied. Hepatic T4-ORD was examined with an ANOVA followed by least square means *post-hoc t*-tests of all pairwise comparisons without adjustments. RM ANOVAs initially identified overall significant treatment effects on thyroid hormones or clinical chemistry for each OPFR compared to controls, followed by 1-way ANOVAs to identify significant treatment differences at individual time periods when overall significance was identified. Spearman's correlation analysis was used to determine statistical associations between hepatic lipid and TBOEP concentrations, and plasma concentrations of albumin and thyroid hormones. Means \pm SEM are presented and statistical significance is considered when $p \leq 0.05$.

Results

The Pilot Study, 2011

In the pilot study, the exposure of the kestrels to the lower concentration of TDCIPP had no effect on circulating TT₃ and TT₄ levels (P-values ≥ 0.10) or plasma cholinesterase activity (KW Chi-square P = 0.65), but resulted in lower brain cholinesterase activity (41.9 ± 0.8 $\mu\text{mol/min/g}$) compared to controls (44.4 ± 0.6 $\mu\text{mol/min/g}$) (KW Chi-square = 4.45 df = 1 P = 0.03). This represented a 5% reduction in the brain cholinesterase activity of the TDCIPP-exposed kestrels compared to the control kestrels. Based on numerous studies with cholinesterase-inhibiting pesticides in birds, this degree of suppression of cholinesterase activity

in brain does not seem to be associated with physiological or behavioral deficits (Grue et al. 1991, 1997).

Cholinesterase activity, clinical chemistry, and plasma thyroid hormones: 2012

For the adult male American kestrels involved in the main study conducted with the four OPFRs in 2012, cholinesterase activity and clinical chemistry values are presented in Table S1. To the best of our knowledge, this is the first report of such information captured in one publication.

In 2012, there were significant effects identified for each chemical in terms of changes in concentrations of one or both thyroid hormones in the male kestrels. All comparisons are made to control male kestrels only. For the kestrels exposed to TBOEP, there was a significant overall effect on plasma FT₄ (RM ANOVA: Trt: $F_{3,10} = 3.60$ $P = 0.05$) and TT₃ (RM ANOVA: Trt: $F_{3,12} = 5.87$ $P = 0.01$), as well as the temporal changes in FT₄ (RM ANOVA: Trt*Time: $F_{2,11} = 4.96$ $P = 0.03$). In addition, the exposure to TBOEP for 7 d also significantly increased FT₃ concentrations ($F_{1,15} = 5.41$ $P = 0.03$) and TT₃ concentrations ($F_{1,15} = 20.23$ $P = 0.0005$).

When the captive kestrels were exposed to TCEP, there was a significant overall effect on FT₄ (RM ANOVA: Trt: $F_{3,11} = 6.16$ $P = 0.01$) and FT₃ concentrations (RM ANOVA: Trt: $F_{3,13} = 6.83$ $P = 0.005$), and the temporal changes in FT₄ (RM ANOVA: Trt*Time: $F_{2,12} = 8.28$ $P = 0.006$) and FT₃ concentrations (RM ANOVA: Trt*Time: $F_{2,14} = 6.81$ $P = 0.009$). The exposure to TCEP for 7 d resulted in increased concentrations of both FT₃ ($F_{1,16} = 19.27$ $P = 0.0005$) and FT₄ ($F_{1,14} = 7.42$ $P = 0.02$).

The exposure of the male kestrels to TCIPP had a significant effect overall on their plasma FT₃ concentrations (RM ANOVA: Trt: $F_{3,13} = 4.04$ $P = 0.03$) and the temporal changes in FT₃ (RM ANOVA: Trt*Time: $F_{2,14} = 6.03$ $P = 0.01$), with a significant suppression of FT₃

concentrations at 14 d exposure ($F_{1,16} = 4.91$ $P = 0.04$). There were no significant effects of the TCIPP exposure on plasma FT₄ or TT₄ concentrations in the kestrels.

The exposure to the higher TDCIPP concentration used in 2012, had a significant overall effect on the plasma FT₃ concentrations (RM ANOVA: Trt: $F_{3,13} = 3.48$ $P = 0.05$) and the temporal changes in plasma FT₃ (RM ANOVA: Trt*Time: $F_{2,14} = 5.49$ $P = 0.02$), with a significant elevation of FT₃ concentrations at 7 d exposure ($F_{1,16} = 5.28$ $P = 0.04$). The exposure to TDCIPP also significantly increased plasma concentrations of TT₄ at 7 d ($F_{1,16} = 11.02$ $P = 0.005$) and suppressed them at 21 d ($F_{1,16} = 4.37$ $P = 0.05$), contributing to significant overall effects on plasma TT₄ levels (RM ANOVA: Trt: $F_{3,13} = 7.25$ $P = 0.004$) and temporal changes in TT₄ concentrations (RM ANOVA: Trt*Time: $F_{2,14} = 11.31$ $P = 0.001$). There was also a significant overall effect of TDCIPP on the temporal changes of FT₄ (RM ANOVA: Trt*Time: $F_{2,11} = 4.03$ $P = 0.05$).

Table S1. Cholinesterase activity and plasma clinical chemistry concentrations measured in captive adult male American kestrels.

Variable	Unit of Measure	N	Minimum	Maximum
<i>Cholinesterase Activity</i>				
Brain Acetylcholinesterase Activity	OD/min/mg	35	5	8
Brain Butrylcholinesterase Activity	OD/min/mg	35	0	0
Brain Muscarinic Cholinergic Receptor	fmol/mg	35	269	843
Brain Nicotinic Cholinergic Receptor	fmol/mg	35	14	40
Plasma Acetylcholinesterase Activity	OD/min/mg	36	1	5
Plasma Butrylcholinesterase Activity	OD/min/mg	34	0.0	0.2
<i>Clinical Chemistry</i>				
Albumin:Globulin Ratio		38	0.5	2
Alanine aminotransferase activity (ALT)	U/L	38	34	297
Gamma-glutamyl transpeptidase (GGT)	U/L	38	4	7
Albumin (%)	%	38	21	55
Albumin	g/dL	38	1	3
Bile acids	umol/L	33	1	161
Total Proteins (TP)	g/dL	38	3	6
Aspartate aminotransferase (ASAT)	U/L	38	37	366
Lactate dehydrogenase isozymes (LDH)	U/L	38	1708	10632
Alkaline Phosphotase (ALP)	U/L	38	51	842
Glutamate dehydrogenase activities (GLDH)	U/L	38	0.1	17

Fig. S1. Circulating concentrations of TT₃ and TT₄ at 7 d exposure in adult male American kestrels that were exposed daily to TBOEP, TCEP, TCIPP or TDCIPP (22 ng OPFR/g kestrel/d) for a total of 21 d. Means \pm standard errors about the means (SEMs) are presented. Statistically significant p-values are indicated as follows: $p \leq 0.0001$ (***); $p \leq 0.01$ (**); $p \leq 0.05$ (*).

S1a. Circulating TT₃ concentrations.


S1b. Circulating TT₄ concentrations.

Fig. S2. Significant changes in thyroid gland histology measures, notably epithelial cell height and volume of colloid, were observed in adult male kestrels at 21 d exposure to TBOEP, TCEP, TCIPP or TDCIPP. Means \pm SEMs are presented. Statistically significant p-values are indicated as follows: $p \leq 0.0001$ (***); $p \leq 0.01$ (**); $p \leq 0.05$ (*).

S2a. Thyroid gland: epithelial cell height (ECH).

S2b. Thyroid gland: colloid area.

S1a. Circulating TT₃ concentrations.



S1b. Circulating TT₄ concentrations.






Fig. S2a. Thyroid gland: epithelial cell height (ECH).



S2b. Thyroid gland: colloid area (data for perimeter of colloid not shown).



Literature Cited

- Basu, N., Scheuhammer, A. M., Sonne, C., Letcher, R. J., Born, E. W., Dietz, R. Is dietary mercury of neurotoxicological concern to polar bears (*Ursus maritimus*)? *Environ. Toxicol. Chem.* **2009**, 28, 133-140.
- Bocian-Sobkowska, J., Wozniak, W., Malendowicz, L. K. Morphometric studies on the development of the human thyroid gland. II. The late fetal life. *Histol. Histopath.* **1997**, 12, 79-84.
- Cray, C., Andreopoulos, A. Comparison of two methods to determine plasma bile acid concentrations in healthy birds. *J. Avian Med. Surg.* **2003**, 17, 11-15.
- Cray, C., King, E., Rodriguez, M., Decker, L.S., Arheart, K.L. Differences in protein fractions of avian plasma among three commercial electrophoresis systems. *J Avian Med. Surg.* **2011**, 25, 102-110.
- Grue, C.E., Hart, A.D.M., Mineau P. Biological consequences of depressed brain cholinesterase activity in wildlife. In *Cholinesterase-inhibiting insecticides: their impact on wildlife and the environment*; Mineau, P., Ed. Elsevier: New York 1991; pp 151-209.
- Grue, C.E. Gibert P.L., Seeley, M.E. Neurophysiological and behavioral changes in non-target wildlife exposed to organophosphate and carbamate pesticides: thermoregulation, food consumption, and reproduction. *Amer. Zool.* **1997**, 369-388.
- Hill, E.F., Fleming, W. J. Anticholinesterase poisoning of birds: field monitoring and diagnosis of acute poisoning. *Environ. Toxicol. Chem.* **1982**, 1, 27-38.
- Luna, L. G. Manual of histological staining methods of the Armed Forces Institute of Pathology. 3rd edn. McGraw Hill, New York. 1968.

- Martinson, S. C., Kimmins, S., Letcher, R. J., Palace, V. P., Bird, D. M., Ritchie, I. J., Fernie, K. J. Diet exposure to technical hexabromocyclododecane (HBCD) affects testes and circulating testosterone and thyroxine levels in American kestrels (*Falco sparverius*). *Environ. Res.* **2011**, *111*, 1116-1123.
- Park, B. J., Palace, V. P., Wautier, K., Gemmill, B., Tomy, G. Thyroid axis disruption in juvenile Brown trout (*Salmo trutta*) exposed to the flame retardant β -tetrabromocyclohexane (β -TBECH) via the diet. *Environ. Sci. Technol.* **2011**, *45*, 7923-7927.
- Rutkiewicz, J., Crump, D., Scheuhammer, A. M., Jagla, M., Basu, N. Investigation of spatial trends and neurochemical impacts of mercury in herring gulls across the Laurentian Great Lakes. *Environ. Pollut.* **2010**, *158*: 2733-2737.
- Rutkiewicz, J., Bradley, M., Mittal, K., Basu, N. Methylmercury egg injections: Part 2- Pathology, neurochemistry, and behavior in the avian embryo and hatchling. *Ecotox. Environ. Saf.* **2013**, *93*, 77-86