

2-6-2015

New distributional records for pleasing lacewings (Neuroptera: Dilaridae, *Nallachius* spp.) in the Americas

David E. Bowles

Missouri State University, davidbowles@missouristate.edu

Atilano Contreras-Ramos

Instituto de Biología, acontreras@ib.unam.mx

Mariza A. Sarmiento-Cordero

Instituto de Biología, mariza.sarmiento@ibunam2.ibiologia.unam.mx

Michael L. Ferro

Louisiana State Arthropod Museum, spongymesophyll@gmail.com

Follow this and additional works at: <http://digitalcommons.unl.edu/insectamundi>

Bowles, David E.; Contreras-Ramos, Atilano; Sarmiento-Cordero, Mariza A.; and Ferro, Michael L., "New distributional records for pleasing lacewings (Neuroptera: Dilaridae, *Nallachius* spp.) in the Americas" (2015). *Insecta Mundi*. 915.
<http://digitalcommons.unl.edu/insectamundi/915>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0406

New distributional records for pleasing lacewings
(Neuroptera: Dilaridae, *Nallachus* spp.) in the Americas

David E. Bowles
National Park Service
Heartland Inventory and Monitoring Network
c/o Department of Biology
Missouri State University
901 South National Ave., Springfield, MO 65897 U.S.A.

Atilano Contreras-Ramos and Mariza A. Sarmiento-Cordero
Instituto de Biología, UNAM, Depto. de Zoología, Apdo.
Postal 70-153, 04510 México, D.F., Mexico

Michael L. Ferro
Louisiana State Arthropod Museum
Department of Entomology
Louisiana State University Agricultural Center
404 Life Sciences Building
Baton Rouge, LA 70803 U.S.A.

Date of Issue: February 6, 2015

David E. Bowles, Atilano Contreras-Ramos, Mariza A. Sarmiento-Cordero, and
Michael L. Ferro
New distributional records for pleasing lacewings (Neuroptera: Dilaridae,
Nallachus spp.) in the Americas
Insecta Mundi 0406: 1–10

ZooBank Registered: urn:lsid:zoobank.org:pub:8A9BBBCD-D289-44FC-9219-3297F73BC7C2

Published in 2015 by

Center for Systematic Entomology, Inc.

P. O. Box 141874

Gainesville, FL 32614-1874 USA

<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Chief Editor: Paul E. Skelley, e-mail: insectamundi@gmail.com

Head Layout Editor: Eugenio H. Nearn

Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas

Review Editors: Listed on the **Insecta Mundi** webpage

Manuscript Preparation Guidelines and Submission Requirements available on the **Insecta Mundi** webpage at: <http://centerforsystematicentomology.org/insectamundi/>

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and Agrifood Canada, Ottawa, ON, Canada

The Natural History Museum, London, UK

Muzeum i Instytut Zoologii PAN, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Eugenio H. Nearn

New distributional records for pleasing lacewings (Neuroptera: Dilaridae, *Nallachius* spp.) in the Americas

David E. Bowles
National Park Service
Heartland Inventory and Monitoring Network
c/o Department of Biology
Missouri State University
901 South National Ave., Springfield, MO 65897 U.S.A.
David_Bowles@nps.gov

Atilano Contreras-Ramos and Mariza A. Sarmiento-Cordero
Instituto de Biología, UNAM, Depto. de Zoología, Apdo.
Postal 70-153, 04510 México, D.F., Mexico
acontreras@ib.unam.mx, mariza.sarmiento@ibunam2.ibiologia.unam.mx

Michael L. Ferro
Louisiana State Arthropod Museum
Department of Entomology
Louisiana State University Agricultural Center
404 Life Sciences Building
Baton Rouge, LA 70803 U.S.A.
spongymesophyll@gmail.com

Abstract. We report on collections of seven species of pleasing lacewings (Neuroptera: Dilaridae) from the Americas. New country distributional records are reported for *Nallachius pulchellus* (Banks) from Honduras and Trinidad, and *Nallachius phantomellus* Adams from Ecuador. *Nallachius ovalis* Adams and *Nallachius prestoni* (McLachlan) are reported from Brazil representing the second reported collections of those species. Additional state records and in-country distributional information are presented for the other species. Two species could not be identified with certainty.

Key Words. Neotropics, biological diversity

Introduction

Pleasing lacewings (Neuroptera: Dilaridae) are a small, poorly known, and rarely collected family consisting of four genera and about 70 species (Oswald 1998; Machado and Rafael 2010). Male dilarids have prominent pectinate antennae (see Machado and Rafael 2010) and females have a long ovipositor, which recurves over the abdomen. The biology of this family is poorly described, due in part to their rarity. Larvae are known to inhabit decayed wood or under the bark of dead trees (MacLeod and Spiegler 1961; Penny 1981; Oswald 1998). Penny (1981) indicated that dilarids appear to be predators that have been successfully reared on soft-bodied insect larvae and eggs. The Dilaridae are geographically widespread (Oswald 1998), but only 20 species are known from the Western Hemisphere and all belong to the genus *Nallachius* Navás in the subfamily Nallachiinae. Relatively few previously published distributional records exist for the known species in the Americas (Adams 1970; Penny 1981, 1994, 2002; Hoffman 1990; Maes and Flint 1994; Kuhar 1995; Monserrat 2005; Machado and Rafael 2010). Distributional records for dilarids from Mexico and Central and South America are particularly scarce (Table 1). Additional distributional and phenological information for pleasing lacewings helps us better understand the diversity of this rare group of insects. Here we report on the distributions of seven species of *Nallachius* from the Americas. We also provide updated distributional records for all species of *Nallachius* in the Americas (Table 1).

Methods

Genitalia were dissected from specimens, cleared in room temperature saturated sodium hydroxide, and examined using stereo- and compound microscopes. Cleared genitalia were stored in glycerin in a genitalia vial placed on the pin below the specimen or placed in the vial. Collection acronyms are as follows: Colección Nacional de Insectos del Instituto de Biología, **Universidad** Nacional Autónoma de México [CNIN-UNAM]; Florida State Collection of Arthropods [FSCA]; Louisiana State University, Louisiana State Arthropod Museum [LSAM]; University of Missouri, Columbia, W. R. Enns Entomology Museum [UMRM]; North Carolina State University Insect Museum [NCSU]; Oklahoma Baptist University [OBU]; Oklahoma State University, K.C. Emerson Entomology Museum [OSEC]. The emergence chamber used to collect some specimens is described in Ferro and Carlton (2011). Adams (1970) provided a key to the species of *Nallachius*, which includes the species reported here. Information in brackets was added by the authors.

While not a substitution for examination of physical specimens, photographic records are an important and valid means to learn more about the distribution and phenology of species. The Internet website BugGuide.net (2014) is a crowd-sourced portal for the aggregation of photographs of insects and other arthropods found within the United States and Canada. Submitted photos are identified to the lowest level possible and placed within the online “guide”. Photographs not retained by BugGuide.net (“frass”) are removed after 30 days. All other photographs are retained for the duration of the existence of the site; however, photographs and accompanying information can be deleted or modified by the submitter at any time. Archival web services, such as the WayBack Machine (archive.org/web/), also maintain copies of BugGuide.net taken at multiple instances in time.

BugGuide.net shows several photographs of *N. americanus* and *N. pulchellus* that were taken at numerous locations in the U.S. The photographed specimens have not been physically examined, may not have been deposited in museums, and/or may no longer exist. However, the photographs unambiguously depict *N. americanus* and *N. pulchellus* and are available for future scrutiny. Herein, we reference those photographs as supplemental records for the species. The URL for each “page” with an image begins with <bugguide.net/node/view/> and is designated with a unique multi-digit “page number”. For example, <bugguide.net/node/view/686584/> returns an image taken on 7 August 2012, submitted by L. Williams. For each record below the unique “page number”, 686584 in the example above, is provided. When multiple photos of an individual specimen are presented all “page numbers” are given. Specimen level collection data based on photographs depicted on BugGuide.net follow.

Results and Discussion

We report distributional records for seven species of *Nallachius* from Argentina, Ecuador, Honduras, Mexico, Trinidad, and the United States (Table 1). New country distribution records for *Nallachius pulchellus* Banks are reported from Honduras and Trinidad, and *Nallachius phantomellus* Adams is newly reported from Ecuador. *Nallachius prestoni* (McLachlan) is reported for only the second time from Brazil, and a new state record for *Nallachius ovalis* Adams is reported for that country as well. Several new state and parish distribution records are reported *Nallachius americanus* (McLachlan) and *Nallachius pulchellus* Banks from Mexico and the United States (Fig. 1). Two series of *Nallachius* collected from Argentina could not be reliably identified to species. Country-level species distribution of Dilaridae in the Western Hemisphere is depicted in Figure 2.

Nallachius americanus (McLachlan)

Nallachius americanus has been previously reported from the United States, Puerto Rico, and Venezuela (Carpenter 1940, 1947; Steyskal 1944; Gurney 1947; Stange 1961; MacLeod and Spiegler 1961; Adams 1970; Sperka 1972; Lawson and McCafferty 1984; Hoffman 1990; Kuhar 1995). Previously published records for the United States include the District of Columbia, Florida, Georgia, Indiana, Kentucky, Maryland, Michigan, Pennsylvania, South Carolina, Texas, and Virginia (Gurney 1947; Ma-

cLeod and Spiegler 1961; Adams 1970; Lawson and McCafferty 1984; Hoffman 1990; Kuhar 1995). New state records are reported here for *N. americanus* for the United States including Arkansas, Louisiana, Missouri, North Carolina, Oklahoma, and Tennessee.

Several of the reported localities from BugGuide.net shown here in supplemental records represent potential new state records for *N. americanus* including Kansas and New Jersey.

UNITED STATES: **Arkansas:** Conway Co., Petit Jean St. Pk.; 7 Hollows Tr., 35° 06' 43" N, 92° 56' 59" W, 05.viii.2008; R.L. Brown, male [NCSU]; Newton Co., Buffalo Nat. River, Fitton Cave Trail, 2.5 km NW Erbie, Campground, 36° 05' 35" N, 93° 14' 10" W, Malaise Trap, 24-28.vi.1994, C. E. Carlton, 1 male (in alcohol) (LSAM). **Louisiana:** Tangipahoa Par., 26.viii.1978, LSAM0019234, Fluker, 1 male (LSAM); St. Tammany Par., 4.2 mi. NE Abita, Springs, sec. 24, T6S, R12E, 7.ix.1990, V. A. Brou, UV, 1 male (LSAM); same data, 09.ix.2006, V. A. Brou, 1 male (LSAM); West Feliciana Par., Feliciana Preserve, 30°47.68' N, 91°15.24' W, 27.viii.2005 UV/MV light, coll. M. Gimmel, 1 male (LSAM); same data, M. Ferro, 7 males (LSAM); same data, but Feliciana Presrv., 30°47.760' N, 91° 15.356' W, 25.iii-22. iv.2007, LN34B -CWD Rear 1, M. Ferro, LSAM 0173457, [collected with emergence chamber from hardwood, decay class 2], 1 male (LSAM); same data, but 7.5 mi. W. of St. Francisville, 30° 47.705' N, 91°15.255' W, 02.x.2004, M. Ferro, blacklight, 1 male (LSAM), 2 males (1 in alcohol) (UMRM). **Mis-**




Figure 1. State-level distribution of *Nallachius americanus* (diamonds) and *Nallachius pulchellus* (circles) in the United States and Mexico. B = record from BugGuide.net.

souri: [Boone Co.], Ashland, 13.vii.1989, D. Utt, Malaise trap, male [collection information is held at Enns Entomology Museum, University of Missouri, Columbia, Missouri, but actual specimen is held by D. Utt, OBU]. **North Carolina:** Wake Co., Raleigh, 1402 Lorimer Rd., 35.7786° N, 78.7035° W, 7.viii.2010, P.L. Mullins, A.F. Ernst, MV light trap, male [NCSU]. **Oklahoma:** [Logan Co.], Guthrie, 5 mi. S. on Sooner, 22.vii.1998, R. Grantham 1 male (in alcohol) (OSEC); same, but 28.vii.1998, 1 male (in alcohol) (OSEC). **Tennessee:** Blount Co. GSMNP, Tremont, 35°37.308' N, 83°40.447' W, 18.v–24.vi.2006, SN25B -CWD Rear 1, M. Ferro, [collected with emergence chamber from hardwood decay class 3–4], 1 larva (in alcohol) (LSAM); Sevier Co. GSMNP, Sugarlands QW, 35°39.826' N, 83°31.509' W, 24.vi–15.vii.2006, SN35C -CWD Rear 1, M. Ferro, [collected with emergence chamber from hardwood decay class 3–4], 1 larva (in alcohol) (LSAM); Greenbrier Field Station – UT, 35° 44.315' N, 83° 25.417' W, 10.vi.2010, Col. M. Ferro, Mercury vapor and blacklight, 1 male (LSAM).

Supplemental Records. UNITED STATES: Florida: Desoto Co.: Arcadia, 3.iii.2012, J. Evoy, (617904); Leon Co.: Tallahassee, 23.v.2014, P. Homann (925454). Okaloosa Co.: Niceville, 25.iv. 2012, M. Friedman, (634057). **Georgia:** Chatham Co.: Wilmington Island, Savannah, 18.v.2006, J. Barrett, at light, (62630, 62631); Clarke Co.: Athens/Sandy Creek Park, 22.iv.2012, C. Champagne, underside of leaf, (636440); Douglas Co.: Winston, Moccasin Lake, 3.viii.2008, J. Armstrong, at light, (218752, 218753); same data, 6 August 2008, (250882); Henry Co.: McDonough, 12.ix.2011, Doug, (577121). McIntosh Co.: Sapelo Island, 23.v.2014, M. Zhang (940064). **Kansas:** Johnson Co.: Lenexa, 23.viii.2012, B. Betros, “Came to MVL in a woodland/prairie area.” (694362, 694363). **Louisiana:** Caddo Par.: ~6 mi E of Vivian, 2.ix.2013, R. M. Taylor Jr. (835027). **Missouri:** Clay Co.: Liberty, 7.viii.2012, L. Williams, at porch light at night, (686584). **New Jersey:** Camden Co.: Atco, 21.vi.2011, Y. Alexander, on oak leaf, (532757). **North Carolina:** Halifax Co., 31.vii.2010, J. M. Lynch, at light sheet, (437274); Orange Co.: Duke Forest off of Hwy 751, 25.viii.2007, P. Coin “Cotinis”, light sheet, (140444); Warren Co., 25.viii.2011, P. Scharf, (591945). **Oklahoma:** Mayes Co.: Pryor, 3.ix.2012, B. Webster, attracted to porch light, (699608, 699609). **Pennsylvania:** Berks Co.: West Reading, 3.vii.2011, J. Eckert (878292). **South Carolina:** Aiken Co.: Spiderweb, 17.vi.2010, J. C. Jones, at light, (412715, 412716); same data, 5.viii.2010, (437999); same data, 19.vii.2011, (548773); same data, 24.v.2011, (548781). Dorchester Co.: McAlhany Nature Preserve, 24.v.2014, J. Carpenter (934550). **Texas:** Fayette Co.: Schulenburg, 13.ix.2013, A. Hendrickson (842745, 842746).

Nallachius pulchellus Banks

The type specimen of *Nallachius pulchellus* is from Cuba, but the species is widely distributed throughout the Americas and, south of the United States, its distribution broadly overlaps with that of *N. americanus*. Oswald et al. (2002) noted that there were no published records of dilarids from Mexico, but that specimens of *Nallachius pulchellus* (Banks) were known from north of Mazatlán in Sinaloa and from Madre María Island off the coast of Nayarit. Later, Monserrat (2005) reported *N. pulchellus* from Jalisco, Mexico. In addition to those records, previously published occurrences for *N. pulchellus* are Arizona (Carpenter 1940, Adams 1970), Costa Rica (Hoffman 1990), and Dominican Republic (Monserrat 2005). We recently examined museum specimens from CNIN-UNAM and FSCA that substantially increase the range of *N. pulchellus* in Mexico, Central America and the Caribbean. These include records from the states of Chiapas, Colima, Morelos, Puebla, Sinaloa, Sonora, and Veracruz, Mexico, representing the first published records for these states. *Nallichus pulchellus* previously had been recorded only from tropical dry forest in Mexico, but these recent collection records show the species inhabits a much broader range of habitats. We also report the occurrence of *N. pulchellus* from Honduras and Trinidad marking the first time the species has been recorded from those countries.

HONDURAS: Olancho: Sierra de Agalta, 8 km N.E. Catacamas, 30.iv.1993, 3000', L. A. Stange, R. Miller, 2 males (FSCA). **MEXICO: Chiapas:** Ixtapa, 11.iv.1962, F. D. Parker, La. A. Stange, 1 male (FSCA); **Colima:** 5.8 Km. NW, Ixtlahuacán, 19° 01' 15.7" N, 103° 46' 37.8" W, 25.iv.2006, S. Zaragoza, F. A. Noguera, E. González, E. Ramírez, L. Salas, blacklight trap, elevation 345 m, 1 male (CNIN-UNAM); **Jalisco:** El Limón, San Buenaventura, 19° 47.614' N, 104° 03.324' W, 9.ii.1997, S. Zaragoza,




Figure 2. Country-level species distribution of Dilaridae in the Western Hemisphere. *Nallachius adamsi* Penny: 1; *Nallachius americanus* (McLachlan): 2; *Nallachius bruchi* Navás: 3; *Nallachius championi* (Navás): 4; *Nallachius bicolor* Adams: 5; *Nallachius furcatus* Machado: 6; *Nallachius hermosus* (Banks): 7; *Nallachius infuscatus* Penny: 8; *Nallachius limai* Adams: 9; *Nallachius loxanus* Navás: 10; *Nallachius maculatus* Penny: 11; *Nallachius martosi* Monserrat: 12; *Nallachius ovalis* Adams: 13; *Nallachius parkeri* Penny: 14; *Nallachius phantomellus* Adams: 15; *Nallachius potiguar* Macado and Rafael: 16; *Nallachius prestoni* (McLachlan): 17; *Nallachius pulchellus* (Banks): 18; *Nallachius pupillus* Navás: 19; *Nallachius reductus* Carpenter: 20.

F. A. Noguera, E. González, E. Ramírez, blacklight trap, elevation 720 m, 1 male (CNIN-UNAM); same data, 6.xi.1996, 1 male (CNIN-UNAM); entre [between] San Buenaventura y [and] Los Yesos, 4 Km. SW San Buenaventura, 19° 45.722' N, 104° 02.875' W, 6.ii.1997, S. Zaragoza, F. A. Noguera, E. González, E. Ramírez, blacklight trap, elevation 740 m, 1 male (CNIN-UNAM); **Morelos**: Tlaquiltenango, Presa Lorenzo Vázquez W Huautla, Estación CEAMISH, 18° 27' 53.59" N, 99° 02' 21.32" W, 14.ii.1996, S. Zaragoza, F. A. Noguera, E. González, E. Ramírez, blacklight trap, elevation 940 m, 1 male (CNIN-UNAM); same data, 18° 27' 47.51" N, 99° 02' 07.9" W, 15.ii.1996, 1 male (CNIN-UNAM); same data, 13.iv.1996, 2 males (CNIN-UNAM; all CNIN-UNAM specimens in 80% ethyl alcohol); **Puebla**: 3 mi. E. Ixtapa de Matamoros, 24.IV.1962, F. D. Parker, L. A. Stange, 1 male (FSCA); **Sinaloa**: 16 mi. S. Guamuchil, 20.v.1962, F. D. Parker, L. A. Stange, 2 female (FSCA); **Sonora**: La Aduana, 22.v.1962, F. D. Parker, L. A. Stange, 1 male (FSCA); **Veracruz**: Sumidero (near Fortín de las Flores), Planta de la Cervecería Moctezuma (power plant in canyon), residence of Ing. Daniel Rábago, 19.v.1965, H. V. Weems, Jr., blacklight trap near stream, elevation 2500–3000 feet [762–1067 m], 1 male (in alcohol) (FSCA). **TRINIDAD**: Simla [William Beebe Tropical Research Station, 10°41'1"N, 61°17'W] 2-13.v.77, P. Feinsinger, Malaise trap, 2 males (in alcohol) (FSCA). **UNITED STATES**: **Arizona**: Cochise Co., Coronado NF, Chiricahua Mts., 31.86864° N, 109.18845° W, 31.vii.2010, Southfork FC, blacklight MV/UV (57), M. Ferro, 1 female (LSAM); same data, 04.viii.2012, 1 male (LSAM); Coronado NF, Chiricahua Mts., 31° 50.431' N, 109° 07.752' W, 04.viii.2010, Sulphur Canyon, blacklight MV/UV, M. Ferro, 1 male (LSAM).

Supplemental records. UNITED STATES: **Arizona**: Pima Co.: Gardner Canyon, 24.vii.2013, CreationsFinest, at light (877253); Santa Cruz Co.: Madera Canyon, upper parking lot, 4.ix.2013, M. Brummermann, (837080, 837138).

Nallachius ovalis Adams

Adams (1970) described this species from Nova Teutônia, state of Santa Catarina, Brazil. It was previously only known from the type series, which includes the holotype, and male and female paratypes. We examined a large series of *Nallachius ovalis* collected from the type locality. To our knowledge, this represents only the second collection of this species, the first being the type series. It is not clear why these additional specimens were not included in Adams (1970), although they were clearly collected about the same time as the type series. Machado and Rafael (2010) reported on a collection of *Nallachius dicolor* Adams, which also is known from Nova Teutônia, state of Santa Catarina, Brazil. These two species are closely related, and their genitalia are largely indistinguishable. They can be easily separated on the basis of their unique wing venation. The costal area of *N. ovalis* is wider than that of *N. dicolor*, and while *N. ovalis* has most of the distal costal veinlets forked, the veinlets are simple in *N. dicolor*. The three specimens from São Paulo, Brazil were in alcohol and slightly faded, and the genitalia would not completely clear. Nonetheless, the observable characters indicate this species is *N. ovalis*, and this locality represents a new state record for the species in Brazil.

BRAZIL: **Santa Catarina**, Nova Teutônia, 300 m, 1-12-x.1962, F. Plauman, 27°11' S, 52°23' W, 13 males, 16 females (FSCA). **São Paulo**, Sierra do "Barati", xi.1969, M. Alvarenga, 3 males (in alcohol) (FSCA).

Nallachius phantomellus Adams

We examined a single male specimen of this species collected in Ecuador. It agreed in most aspects with the description and illustrations presented in Adams (1970) and Machado and Rafael (2010). The distinctive wing maculation, and configuration of the gonocoxites and internal genital armature matched those described by Adams. The specimen we examined differed in having the tip of gonocoxite entire rather than serrate as in *N. phantomellus*. The type material described by Adams (1970) was collected from the Rio Caragualà, Brazil. The specimen we examined from Ecuador was from some 3,000 km to the northwest of the type locality. This collection suggests that *N. phantomellus* has a more widespread

distribution in South America than previously reported, or the specimen from Ecuador represents an undescribed and closely related species. Until additional material becomes available we are considering this species to be *N. phantomellus*. *Nallachius loxanus* Navás, a member of the *americanus* species group, is the only other dilarid species known from Ecuador.

ECUADOR: Orellana Province, Yasuni NP, Yasuni Res. Sta., 00°40.4"S, 76°23.86"W, 01 Jul 2008, M. Ferro, MV Blacklight, 1 male (LSAM).

Nallachius prestoni (McLachlan)

We examined a single male of this species that was collected near the type locality in Rio de Janeiro, Brazil. It is similar to *N. reductus* Carpenter and *N. dicolor* Adams, but differs in wing venation and shape (Adams 1970).

BRAZIL: Rio de Janeiro, Serra dos Órgãos, 29.i.1969, J. Lotti, L. Stange 1 male (in alcohol) (FSCA).

Nallachius spp.

The specimens indicated here were labelled as *Nallachius bruchi* Navás by another researcher. The female lectotype, and only known specimen, of this species is from Alta Garcia, Córdoba Province, Argentina. Navás (1923) did not illustrate *N. bruchi* and the description is considered inadequate (Adams 1970). Since Córdoba Province is located south of Tucumán Province and separated by Santiago del Estero Province, we presume the identifier made the assumption that the specimens shown here must be *N. bruchi*. Based on the characters presented for this species by Adams (1970) and the redescription of the female lectotype by Monserrat (2005), the specimens we examined do not appear to be *N. bruchi*. The specimens in alcohol were badly faded, but wing venation could be discerned. In the forewing, MP2 and CuA were not fused. Nearly all coastal veinlets were unforked except for one specimen, which had a couple of forked veinlets. The male antennae had 10–11 elongate processes. One of the pinned males was missing genitalia, and the specimens in alcohol would not clear entirely. The male genitalia are most similar to that of *Nallachius dicolor* Adams, but that species currently is known only from Brazil (Adams 1970, Machado and Rafael 2010). However, *N. dicolor* has been collected from Santa Catarina, Brazil, which shares a border with Argentina suggesting that this species also may occur there. While we cannot entirely rule out the previous identifier's assessment that these specimens represent *N. bruchi*, we do not feel that a species name can be assigned to this material in its present condition.

ARGENTINA: Tucumán [Prov.], 430 m, 1.x.1966, W. Weyrauch, light, 1 male (FSCA); same, but Rio (E) Cajon, 11.vi.1972., Acacia [rest of hand-written label illegible], 1 male (FSCA); same, but Raco, 30-xii-1973, L. Stange, 1 female (FSCA); same, but 11 km W. Las Cejas, 7.iii.1967, L. Stange, 2 males (in alcohol) (FSCA); same, but 9-iii-11.iv.1968, Malaise Trap, 1 male (in alcohol) (FSCA); same, but El Soledad, 6-20.x.1966, L. Stange, Malaise trap, 3 males, 1 female (in alcohol) (FSCA).

We also examined a single female specimen of *Nallachius* collected from a different location in Argentina. In addition to the primary locality label, a second label on the specimen pin placed above the location label was printed on red paper with the hand-written words "emergido [emerged] 14.i.2000, árbol seco en pie" [standing dry tree], *Celtis tala* Gill[et]. The former date also suggests the year on the locality label was 3.xi.1999. This is a relatively large specimen with a forewing length of 5.5 mm. In the forewing, the MP2 vein is not fused with the CuA vein, and there are six large brown spots in the coastal area and smaller spots scattered over the remainder of the wing. Most of the distal costal veinlets are simple. These characteristics are consistent with those of *N. dicolor*. Given that this specimen was collected in Northwestern Argentina, it either represents a range extension of *N. dicolor*, or possibly an undescribed species.

ARGENTINA: La Rioja [Prov.], Los Molinos, 3.xi.19[?]9, DiLorio O. leg., 1 female (FSCA).

Acknowledgments

We thank Victor Monserrat (Universidad Complutense, Spain) and Renato Machado (Texas A&M University) for reviewing and providing constructive criticism of an earlier version of this paper. We also thank Paul Skelley, Florida State Collection of Arthropods; Chris Carlton, Louisiana State Arthropod Museum; Dale Utt, Oklahoma Baptist University; Richard Grantham, Oklahoma State University, and Bob Blinn, North Carolina State University Insect Museum loaning us specimens examined for this paper or granting us permission to use label data.

Literature Cited

- Adams, P. A. 1970. A review of the New World Dilaridae. *Postilla* 148: 1–30.
- BugGuide.net. 2014. Iowa State University, Ames. Last accessed 26 August 2014.
- Carpenter, F. M. 1940. A revision of the Nearctic Hemerobiidae, Berothidae, Sisyridae, Polystoechotidae and Dilaridae (Neuroptera). *Proceedings of the American Academy of Arts and Sciences* 74: 193–280.
- Carpenter, F. M. 1947. Taxonomic notes on the Dilaridae (Neuroptera). *Psyche* 54: 100–109.
- Gurney, A. B. 1947. Notes on Dilaridae and Berothidae, with special reference to the immature stages of the Nearctic genera (Neuroptera). *Psyche* 54: 145–169.
- Ferro, M. L., and C. E. Carlton. 2011. A practical emergence chamber for collecting Coleoptera from rotting wood, with a review of emergence chamber designs to collect saproxylic insects. *The Coleopterists Bulletin* 65: 115–124.
- Hoffman, K. M. 1990. Distributional notes on North and Central American Dilaridae (Neuroptera). *Entomological News* 101: 155–157.
- Kuhar, T. P. 1995. A pleasing lacewing, *Nallachius americanus* (McLachan), from southwestern Virginia (Neuroptera: Dilaridae). *Banister* 5: 32–33.
- Lawson, H. R., and W. P. McCafferty. 1984. A checklist of Megaloptera and Neuroptera (Planipennia) of Indiana. *Great Lakes Entomologist* 17: 129–131.
- Maes, J. M., and O. S. Flint, Jr. 1994. Dilaridae, Berothidae y Sisyridae (Planipennia): tres familias nuevas para la fauna de Nicaragua. *Revista Nicaragüense de Entomología* 29: 3–5.
- Machado, R. J. P., and J. A. Rafael. 2010. Two new species of Dilaridae (Insecta: Neuroptera) with additional notes on Brazilian species. *Zootaxa*. 2421: 61–68.
- MacLeod, E. G., and P. E. Spiegler. 1961. Notes on the larval habitat and developmental peculiarities of *Nallachius americanus* (McLachan) (Neuroptera: Dilaridae). *Proceedings of the Entomological Society of Washington* 63: 281–286.
- Monserrat, V. J. 2005. Nuevos datos sobre algunas pequeñas familias de neurópteros (Insecta: Neuroptera: Nevrothidae, Osmylidae, Sisyridae, Dilaridae). *Heteropterus: Revista de Entomología* 5: 1–26.
- Navás, L. 1914. Family Dilaridae. *Neuroptera. Genera Insectorum* 156: 1–14.
- Navás, L., 1923. Estudios sobre Neuropteros (Insectes). *Arxius [Arxivs] de l'Institute [d'Estudis Catalans, Seccio] de Ciencias, Barcelona* 7: 179–203.
- Oswald, J. D. 1998. Annotated catalogue of the Dilaridae (Insecta: Neuroptera) of the World. *Tijdschrift voor Entomologie* 141: 115–128.
- Oswald, J. D., A. Contreras-Ramos, and N. D. Penny. 2002. Neuroptera (Neuropterida). p. 559–581. *In*: J. L. Bosquets, and J. J. Morrone (eds.). *Biodiversidad, taxonomía y biogeografía de Artrópodos de México: hacia una síntesis de su conocimiento*. Vol. 3. Universidad Nacional Autónoma de México, Distro Federal, México. 690 p.
- Penny, N. D. 1981. Neuroptera of the Amazon Basin. Part 2. Dilaridae. *Acta Amazonica* 11: 383–390.
- Penny, N. D. 1994. A new species of *Nallachius* (Neuroptera: Dilaridae) from Costa Rica. *Pan-Pacific Entomologist* 70: 309–312.

- Penny, N. D. 2002.** Family Dilaridae. p. 236–237 (text), 394–397 (figures). In: Penny, N. D. (ed.). A Guide to the Lacewings (Neuroptera) of Costa Rica. Proceedings of the California Academy of Sciences 53: 161–457.
- Sperka, C. L. 1972.** *Nallachius americanus* (McLachlan) (Neuroptera: Dilaridae) in Kentucky. Entomological News 83: 282–283.
- Stange, L. A. 1961.** New records for the genus *Nallachius* Navás (Neuroptera: Dilaridae). Pan-Pacific Entomologist 37: 144.
- Steyskal, G. C. 1944.** Notes on *Nallachius americanus* (McL.) (Dilaridae, Neuroptera). Psyche 51: 183–184.

Received January 12, 2015; Accepted January 20, 2015.

Review Editor Andy Rasmussen.

Table 1. The species of *Nallachius* known from the Americas with distributional information updated from Oswald (1998). Type localities are for holotype males unless otherwise indicated.

Species	Type Locality	Distribution	References
<i>Nallachius adamsi</i> Penny	Brazil: Amazonas, Reserva Ducke	Brazil	Penny 1981; Oswald 1998; Machado and Rafael 2010
<i>Nallachius americanus</i> (McLachlan)	USA: Bee Spring, Kentucky	Costa Rica, Puerto Rico, USA (eastern), Venezuela	McLachlan 1880; Steyskal 1944; Gurney 1947; Adams 1970; Hoffman 1990; Oswald 1998; Penny 2002; this study
<i>Nallachius bruchi</i> Navás	Argentina: Alta Gracia, Córdoba	Argentina	Navás 1923; Carpenter 1947; Oswald 1998; Monserrat 2005
<i>Nallachius championi</i> (Navás)	Guatemala: Cerro Zunil [lectotype, male]	Guatemala	Navás 1914; Adams 1970; Oswald 1998
<i>Nallachius bicolor</i> Adams	Brazil: Nova Teutônia, Santa Catarina	Brazil	Adams 1970; Oswald 1998; Machado and Rafael 2010
<i>Nallachius furcatus</i> Machado and Rafael	Brazil: Paraíba, Cabaceiras (Fazenda Bravo)	Brazil	Machado and Rafael 2010
<i>Nallachius hermosus</i> (Banks)	Colombia: Pacho, E. Cordilleras [female]	Columbia	Banks 1913; Carpenter 1947; Adams 1970; Oswald 1998
<i>Nallachius infuscatus</i> Penny	Brazil: Amazonas, Reserva Ducke	Brazil	Penny 1981
<i>Nallachius limai</i> Adams	Brazil, Nova Teutônia, Santa Catarina	Brazil	Adams 1970; Oswald 1998; Machado and Rafael 2010
<i>Nallachius loxanus</i> Navás	[Ecuador]: Loja, Equateur	Ecuador	Navás 1911; Carpenter 1947; Adams 1970; Oswald 1998
<i>Nallachius maculatus</i> Penny	Brazil: Rondônia, 48 km east of Porto Velho	Brazil	Penny 1981
<i>Nallachius martosi</i> Monserrat	Chile: Talca, Altos de Vilches	Chile	Monserrat 2005
<i>Nallachius ovalis</i> Adams	Brazil: Nova Teutônia, Santa Catarina	Brazil	Adams 1970; Oswald 1998; this study
<i>Nallachius parkeri</i> Penny	Costa Rica: Guanacaste, 3 km SE of Rio Naranjo	Costa Rica	Penny 1994
<i>Nallachius phantomellus</i> Adams	Brazil: Rio Caragualá, Mato Grosso do Sul	Brazil, Ecuador	Adams 1970; Oswald 1998; this study
<i>Nallachius potiguar</i> Machado and Rafael	Brazil: Rio Grande do Norte: Baía Formosa, Mata Estrela	Brazil	Machado and Rafael 2010
<i>Nallachius prestoni</i> (McLachlan)	Brazil: Rio de Janeiro	Brazil	McLachlan 1880; Carpenter 1947; Adams 1970; Oswald 1998; this study
<i>Nallachius pulchellus</i> (Banks)	Cuba: Central Soledad (near Cienfuegos)	Costa Rica, Cuba, Dominican Republic, Honduras, Mexico, Trinidad, USA (western)	Banks 1938; Carpenter 1947; Adams 1970; Hoffman 1990; Oswald 1998; Monserrat 2005; this study
<i>Nallachius pupillus</i> Navás	Paraguay: San Bernardino	Paraguay	Navás 1930; Adams 1970; Oswald 1998
<i>Nallachius reductus</i> Carpenter	Paraguay: Ilalyria [Ualyaia]	Paraguay	Carpenter 1947; Adams 1970; Oswald 1998